neuropathologist. The book is concerned with the mechanism of eye movement and particularly "goal-directed behaviour." The translation reads well, and there are about 300 references of which a fifth are taken from the Russian literature. The chapters deal with structure and function of eye muscles, fixation, tracking, saccadic movements, convergence, and the higher cortical functions.

Despite the conventional arrangement of the book it cannot be regarded as a satisfactory guide to modern knowledge of control of eye movements. Much of the text is concerned with the author's own observations. There are descriptions of disturbed eye movement in patients with aneurysm and other circulatory disorders which are attributed to local alterations in blood flow. Measurements of cerebral blood flow are correlated with "functional loads" such as optokinetic nystagmus and arithmetic. His observations on single motor unit discharge in the rabbit are given in some detail. There are various recordings of rapid eye tremor which do not accord with other publications.

The book is well produced and the index is satisfactory. Its main value is in providing information about work in Russia and the attempts to develop a unifying concept to explain eye movement.

BRYAN ASHWORTH

Circulation of the Blood Edited by D. Geraint James. (Pp. 495; illustrated; £25.00.) Pitman Medical: Tunbridge Wells. 1978.

This handsome volume contains a wide ranging series of papers on various facets of the circulation collected together as a memorial to William Harvey on the occasion of the quarto-centenary of his birth. In some instances the contributions are incisive and authoritative—for example, Professor Sherlock's review of the portal circulation and Professor Browse's incisive coverage of the problem of thrombo-embolism—but most workers in the circulatory field will find something of interest in the volume although the very generality of its coverage and its price, may inhibit distribution of a rather handsome memorial.

LINDSAY SYMON

Atlas of Neonatal Electroencephalography By Sarah S. Werner, Janet E. Stockard and Reginald G. Bickford. (Pp. xii+211; illustrated; price not stated.) Raven Press: New York. 1977. This is a most welcome Atlas, beautifully illustrated and clearly printed. Much attention is given to the recognition of various types of artefacts. The illustrations are selected very carefully, probably from a fairly large material. The so-called "10-20" electrode placement has been modified very sensibly and in addition to the EEG respiratory excursions, electromyography eye movements are displayed polygraphically. Both normal and abnormal features in the EEGs of premature babies from 24 weeks gestation onwards are presented as well as the normal and abnormal features of babies born at term. In addition to ordinary EEG techniques, the last chapter of the book includes a series of "Spectral Monitoring" as applied to the newborn as an alternative type of EEG display.

This Atlas will be most useful not only to clinical neurophysiologists but also to neonatologists. Its format, though convenient for a large desk, may be rather awkward for most bookcases. Both authors and publishers should be congratulated for such an elegant presentation.

G. PAMPIGLIONE

The Neurological Examination of the Full-Term Newborn Infant Second edition. By Heinz F. R. Prechtl. (Pp. 68; illustrated; price not stated.) Spastics International Medical Publications and William Heinemann Medical: London. 1977.

The revision of this well-known monograph will add to its admirers. As number 63 of *Clinics in Developmental Medicine*, it needs no introduction to paediatricians, but the adult neurologist would find it profitable to read the first 10 pages and to consider how the concepts of optimal performance might apply to older subjects. Dr Prechtl's book is not a guide to the localisation of brain disease, but an application of system theory to assess the neonatal nervous system as an information processing system. The method is now quantitative but the user must not extrapolate the normal values given to later age groups.

J. A. SIMPSON

Letters

Human muscle afferent responses to tendon taps

SIR,—The two recent papers by Murthy *et al.* (1978a, b) are of some interest as they describe a relatively simple method for recording the afferent responses elicited by tendon percussion in intact human subjects. Unfortunately criticisms can be made, both of experimental method and assumed facts such that the results presented and conclusions reached cannot be accepted as valid.

1. Their "indifferent" electrode situated "over the radius," presumably close to the superficial branch of the radial nerve which supplies the skin at the site of percussion. Since they stress the advantages of differential recording they should ensure that their indifferent electrode is truly indifferent (cf. Fig. 4a and its legend on page 228).

2. Percussion of the muscle was standardised, and they made measurements of percussion force. It is difficult to determine the effect of manoeuvre when it is not known whether the stimulus remained constant.

3. With the hand placed as in Fig. 4a on page 221, even light percussion would have stimulated receptors in other intrinsic muscles of the hand and in the skin over the ulnar aspect of the hypothenar eminence. Such receptors would contribute to the evoked ulnar nerve potential, and their contribution would not be abolished by anaesthetising the skin over the percussion site. In Fig. 4a on page 223, a small deflection of some 25%, occurring at the appropriate latency, is recorded on tapping skin near the muscle tendon rather than the tendon itself. This potential could be of muscle or skin origin (or both).

4. They state that the "phasic tendon tap is known to be a selective and adequate stimulus to the primary sensory endings of the muscle spindle" (page 220). This statement is incorrect. As any of the workers who have recorded from single muscle afferents in man can attest, light percussion with a finger can excite secondary spindle endings, Golgi tendon organs, and pacinian corpuscles in relaxed muscles even if it excites primary endings better.

The evidence they quote in support of this statement is from experiments in the cat using brief rapid muscle stretch. Under the rigorous conditions that can be imposed in animal experiments, the parameters of stretch can be controlled to allow selective activation of primary endings, but this cannot be done in man.

5. The authors assume that changes in the evoked ulnar nerve potential produced by reinforcement or eye closure result from changes in fusimotor bias. However, they have not eliminated a change in muscle length. Even if they had immobilised the hand (which they did not do), it is not possible to immobilise human subjects completely. Only an accurate strain gauge will detect the small changes in muscle length that can still occur in an immobilised human hand.

6. On page 229 the authors state that, of the factors contributing to the reinforcement of tendon jerks, "the fusimotor effects are more dominant." However, nerve block studies have shown that reflex reinforcement still occurs when the muscle spindle contribution has been abolished by local anaesthetics (Clare and Landau, 1964) or by pressure/ischaemia (Bussel *et al.*, 1978). In addition, Hagbarth *et al.* (1975) have monitored the spindle input during reinforcement manoeuvres and have shown that tendon jerks are potentiated in manoeuvres which did not affect the input from spindles. These studies do not exclude that fusimotor activation contributes to the reflex potentiation seen in the clinical setting, but they do demonstrate that the Jendrassik manoeuvre does not exert its effects predominantly by facilitating fusimotor neurones. This conclusion could also have been reached by Murthy *et al.* by examining their Fig. 4b and c on page 223. The afferent response in Fig. 4e is smaller and more dispersed than in Fig. 4b (although its onset is earlier), but a reflex response is seen in Fig. 4e (although its onset is earlier), but a reflex response is seen in Fig. 4e and not in Fig. 4b. Assuming the experimental conditions were identical (and that is an assumption), some form of central reflex enhancement must have occurred.

7. In single unit recordings Burg *et al.* (1974) demonstrated what they considered to be a fusimotor effect on spindle discharge during reinforcement

manoeuvres, but their methods did not exclude changes in length of the percussed muscle and, assuming that the effects were of fusimotor origin, they did not establish that the effects were due to selective activation of the fusimotor system. When these studies were repeated with appropriate controls, a selective effect of reinforcement manoeuvres on the fusimotor system could not be demonstrated (Hagbarth *et al.*, 1975).

For too long has the neurological literature been plagued by speculations about the role of the fusimotor system in normal and diseased states, based on techniques which are not adequate to implicate or exclude fusimotor involvement conclusively. It would be of great interest if the promising technique described by Murthy *et al.* were applied with more rigorous experimental controls.

DAVID BURKE

*Unit of Clinical Neurophysiology,
The Prince Henry Hospital,
Little Bay, NSW 2036,
Australia.*

References

- Burg, D., Szumski, A. J., Struppler, A., and Velho, F. (1974). Assessment of fusimotor contribution to reflex reinforcement in humans. *Journal of Neurology, Neurosurgery, and Psychiatry*, **37**, 1012-1021.
- Bussel, B., Morin, C., and Pierrot-Deseilligny, E. (1978). Mechanism of monosynaptic reflex reinforcement during Jendrassik manoeuvre in man. *Journal of Neurology, Neurosurgery, and Psychiatry*, **41**, 40-44.
- Clare, M. H., and Landau, W. M. (1964). Fusimotor function. Part V. Reflex reinforcement under fusimotor block in normal subjects. *Archives of Neurology (Chicago)*, **10**, 123-127.
- Hagbarth, K.-E., Wallin, G., Burke, D., and Löfstedt, L. (1975). Effects of the Jendrassik manoeuvre on muscle spindle activity in man. *Journal of Neurology, Neurosurgery, and Psychiatry*, **38**, 1143-1153.
- Murthy, K. S. K., Gildenberg, P. L., and Seeliger-Petersen, W. (1978a). Human muscle afferent responses to tendon taps. 1. Characteristics of the waveform recorded with transcutaneous electrodes. *Journal of Neurology, Neurosurgery, and Psychiatry*, **41**, 220-225.
- Murthy, K. S. K., Gildenberg, P. L., and Seeliger-Petersen, W. (1978b). Human muscle afferent responses to tendon taps. 2. Effects of variations in fusimotor bias. *Journal of Neurology, Neurosurgery, and Psychiatry*, **41**, 226-231.

SIR,—Thank you for providing an opportunity to reply to Dr Burke's criticism of our papers (Murthy *et al.*, 1978a, b). His points are answered in the same sequence.

1. There was no doubt that the contribution to the afferent waveform was mainly from the ulnar nerve with the indifferent electrode over the radius picking up no detectable activity from the radial nerve. Any such activity in the radial nerve would have resulted in a negative phase of the recorded waveform as observed in Fig. 3a (page 228) which was included to show such a possibility.

2. The qualitative evaluation of the recorded waveforms does not suffer from want of more rigorous experimental controls. The variability in the force of the taps would have affected the size of the response quantitatively rather than alter its qualitative features (latency and duration). For more quantitative evaluation modifications can be made in the experimental method, especially in positioning and immobilising the hand, by using an automated percussion device to eliminate the variability in the force of the taps and by using sensitive transducers to record changes in muscle length.

3. As to the contributions from receptors in other intrinsic muscles of the hand and in the skin over the ulnar aspect of the hypothenar eminence, it should be borne in mind that the mechanical vibration travels very slowly in the muscles and also should decay in some fashion with distance from the place where the taps are applied. Controls as illustrated in Fig. 4a (page 233 in Murthy *et al.*, 1978a) permitted us to conclude that contributions from other intrinsic muscles would be insignificant.

4. We would like to question the connection between the qualitative observations of other muscle afferents responding to a local prod on the belly of the muscle and a percussion applied to the muscle tendon in our study. Dr Burke