Nurse F. Smith and Deputy Sister P. Walsh cheerfully assisted at every stage, and Dr. B. Blackwell made publication possible by detailed and constructive criticism of the manuscript. Sir Aubrey Lewis gave support and encouragement. To these and many others unmentioned, the author wishes to express his sincere gratitude.

REFERENCES

Benaim, S. (1957). Group psychotherapy within a geriatric unit: an

experiment. Int. J. soc. Psychiat., 3, 123-128.
Burchard, E. M. L., Michaels, J. J., and Kotkov, B. (1948). Criteria for the evaluation of group therapy. Psychsom. Med., 10, 257-274.

Linden, M. E. (1953). Group psychotherapy with institutionalised senile women: study in gerontologic human relations. Int. J. Gr. Psychother., 3, 150-170.

Rechteschaffen, A. (1959). Psychotherapy with geriatric patients: a review of the literature. J. Geront., 14, 73-84.

Schwartz, E. D., and Goodman, J. I. (1952). Group therapy of obesity in elderly diabetics. Geriatrics, 7, 280-283.

Silver, A. (1950). Group psychotherapy with senile psychotic patients. Geriatrics, 5, 147-150.

Taylor, F. K. (1958). A history of group and administrative therapy in Great Britain, Brit. J. med. Psychol, 31, 153-173.

Thomas, G. W. (1943). Group psychotherapy. A review of the recent literature. Psychosom. Med., 5, 166-180.

Wolff, K. (1957). Group psychotherapy with geriatric patients in a

mental hospital. J. Amer. Geriat. Soc., 5, 13-19.
World Health Organization. (1953). Third Report of the Expert Committee on Mental Health. Technical Report Survey No. 17. Geneva.

Appendix

The nurses' notes were rated for their opinions of the patients under two broad headings: (1) behaviour disturbance and (2) withdrawal. One point was assigned to each word or phrase indicating disturbance or with-

drawal if it appeared on the lists given below. These lists were devised by a previous inspection of the records to be measured and are obviously unique to this experiment.

(1) DISTURBANCE

accusing aggressive agitated annoying antisocial anxious attention seeking

banging behaving badly

blacked out careful handling required

childish collapsing turns complains confused

difficult discharging herself disgruntled disorientated dissatisfied

distressed disturbed disturbing others domineering

dressing required emotional

feeding required frightened frightening

hostile hysterical

insulting irritable manipulative

noisy (being a) nuisance

offensive

paces up and down paranoid

(seeks) reassurance refuses treatment, food, drink, etc resentful restless

sleepless smashes spits stubborn

rude

suspicious tense threatening

troublesome (having) turns

uncooperative unpleasant unpredictable

wandering about

(2) WITHDRAWAL

(in) bed before 8 p.m.

friendless

keeps to himself

laughs to herself lonesome

(poorly) occupied (stays) in own room

pre-occupied (requires) prompting

(very) quiet ·

refuses to mix, O.T., to get up, etc

silent (very) slow solitary (not) talking unoccupied

unsociable vacant

withdrawn

CORRECTION

In the paper by Richard L. Hewer and Norman Robinson, 'Diabetes mellitus in Friedreich's ataxia' (J. Neurol. Neurosurg. Psychiat., 1968, 31, No. 3) the sentence in the summary (page 230) which reads: 'Twelve further cases showed abnormalities of glucose tolerance amounting to diabetes mellitus' should read 'Twelve further cases showed abnormalities of glucose tolerance not amounting to diabetes mellitus'.