

<sup>1</sup>Oxford University Department of Clinical Neurology, John

Radcliffe Hospital, Oxford, UK

Cambridge, Massachusetts,

Neurodegeneration Research,

of Psychiatry, London, UK

Correspondence to

King's College London, Institute

Dr Martin Turner, Department of

Clinical Neurology, West Wing

Level 3, John Radcliffe Hospital,

Oxford OX3 9DU, UK; martin.

turner@clneuro.ox.ac.uk

Received 4 February 2010

Revised 6 March 2010

Accepted 12 April 2010

Published Online First

18 June 2010

<sup>2</sup>PatientsLikeMe Inc.,

<sup>3</sup>MRC Centre for

USA

## Concordance between site of onset and limb dominance in amyotrophic lateral sclerosis

M R Turner,<sup>1</sup> P Wicks,<sup>2</sup> C A Brownstein,<sup>2</sup> M P Massagli,<sup>2</sup> M Toronjo,<sup>3</sup> K Talbot,<sup>1</sup> A Al-Chalabi<sup>3</sup>

#### ABSTRACT

**Background** Focality of onset of amyotrophic lateral sclerosis (ALS) is not understood. Attempts to implicate physical exercise in the aetiology of ALS have provided inconsistent results. If physical use of a limb were important in defining the site of onset, then handedness might be expected to influence the side of upper limb-onset disease and footedness likewise in lower limb-onset ALS. **Methods** ALS patients registered with an internet-based

support site were invited to complete an online questionnaire concerning site of onset of symptoms and their dominant hand and foot. A binomial test of proportions was used to investigate the null hypothesis that handedness and footedness do not influence side of onset in upper and lower limb-onset ALS, respectively.

**Results** 343 ALS patients with limb-onset disease were studied. For upper limb-onset patients, there was concordance for side of onset and handedness (64%; p<0.0006). For lower limb-onset patients, concordance for side of onset and footedness was absent. The frequency of left handedness was commensurate with that found in the general population.

Interpretation These results are potentially consistent with the hypothesis that exercise influences pathogenesis in ALS since routine physical demands on the upper limb are heavily influenced by limb dominance, whereas in the lower limbs the commonest function is standing or locomotion, which uses both legs equally. However, there may also be an inherent cortical vulnerability underlying upper limb-onset laterality, possibly influenced by changes in neuronal connectivity and cortical excitability in relation to handedness and reflected by the "split hand" phenomenon consistently observed in ALS.

The site of onset of loss of function in the neurodegenerative disorder amyotrophic lateral sclerosis (ALS) is approximately divided equally between upper limb, lower limb and bulbar territories,<sup>1</sup> with only a very small proportion ( $\sim$ 3%) beginning in the respiratory or axial musculature.<sup>2</sup>

Although there are a number of active themes in the quest to understand pathogenesis in ALS,<sup>3</sup> very little is known about why the onset is focal, though it has been shown that upper (UMN) and lower motor neuron (LMN) signs (the hallmark of ALS) are maximal within this territory and show contiguous spread.<sup>4</sup>

The inconsistent observations of relative athleticism among people with  $ALS^{5-7}$  have led to continued debate about the possible role of physical exercise in the pathogenesis of ALS,<sup>8</sup> and putative direct effects via excitotoxicity and mitochondrial dysfunction.<sup>3</sup> A more indirect association, whereby the genetic profiles that underlie ALS susceptibility and athleticism have common components,<sup>9</sup> currently seems equally, if not more, likely.

If physical exercise is a direct determinant of motor neuron damage, then it might be expected that the site onset of symptoms in limb-onset ALS would be biased in favour of the dominant limb through relatively greater use. We attempted to confirm the null hypothesis that the side of initial symptoms in limbonset ALS is unrelated to limb dominance.

#### **METHODS**

Patientslikeme.com is an internet-based information and support resource for patients with a variety of diseases. ALS patients who had selfregistered on the site were invited to complete an anonymous on-line questionnaire to establish the site of onset of their symptoms and their dominance for upper limb (UL) use (handedness) and lower limb (LL) use (footedness). The specific hypothesis was not mentioned to reduce bias, and respondents were only able to submit one questionnaire.

Independence of variables was tested using  $\chi^2$  for categorical variables and linear regression for continuous variables.

A binomial test of proportions was used to establish the degree of concordance between the side of onset and handedness or footedness for upper and lower limb-onset ALS patients, respectively. Under the null hypothesis, side of onset and limb dominance should be independent and, therefore, equally likely.

All calculations were performed using SPSS (V.14, SPSS Inc).

#### RESULTS Overall group

A total of 502 patients responded (estimated to be 75% of the total possible). Seventy-two patients were either unable to specify the site of symptom onset (n=13) or entered more than one site for initial symptoms (n=59) and were removed from the subsequent analysis. Of the remaining 430 patients, 55% were male. The mean age of respondents was 56 years (SD 10, 26-82); there were no significant differences in age with respect to sex. The site of initial symptoms was bulbar in 18%, UL 37% (right 23%, left 14%), LL 43% (right 24%, left 19%) and respiratory 2%. There was no significant difference in the proportion of left-handedness or leftfootedness for males versus females, nor for bulbar versus limb-onset patients. The male:female ratio was significantly reversed for bulbar (0.7:1) versus limb-onset patients (1.3:1; p=0.006) but with no significant difference in mean age.

#### Limb-onset sub-group

A total of 343 limb-onset patients were available for analysis. For handedness, 87% reported right dominance, 10% left and 3% equal. For footedness, 86% reported right dominance, 10% left and 4% equal. Ten patients (3%) reported discordance for upper and lower limb dominance.

#### Concordance of handedness to side of UL onset

Excluding those reporting equal dominance for handedness (n=11), for upper limb-onset patients, there was concordance for side of onset and handedness: 97 of 151 (64%) overall (right 90 of 139 and left 7 of 12); p<0.0006.

#### **Concordance of footedness to side of LL onset**

Excluding those reporting equal dominance for footedness (n=13), for the lower limb-onset patients, concordance for side of onset and footedness was absent: 99 of 181 (55%) overall (right 91 of 162 and left 8 of 19); p=0.234.

#### DISCUSSION

This study demonstrated high concordance for handedness and side of upper limb-onset ALS but no similar relationship in relation to footedness and side of lower limb onset.

For the upper limb-onset patients, our finding is potentially consistent with the hypothesis that exercise influences pathogenesis in ALS, since routine physical demands on the upper limb are heavily influenced by limb dominance. For the lower limb-onset patients, the lack of an effect of footedness is also consistent with this hypothesis since the commonest function is standing or locomotion, which uses both legs equally.

However, another potential reason for a difference in the result between upper and lower limb onset is some inherent vulnerability in upper limb motor neuronal organisation and laterality. There is evidence for greater connectivity in the dominant motor cortex with respect to handedness<sup>10</sup>; specifically, it has been noted that the production of movements in dominant and non-dominant hands involves distinct cerebral networks.<sup>11</sup> Evolutionary aspects of neocortical function might have relevance to ALS pathogenesis, in that humans have evolved uniquely complex levels of corticomotoneuronal organisation and connectivity as part of the switch to bipedal locomotion and particularly the development of fine hand control through an opposable thumb.<sup>12</sup> Such complexity may have come at a price and harbour a vulnerability in the ageing motor system unmasked by ALS. The demonstration of a cortical correlate of the "split hand" phenomenon (preferential wasting of the lateral thenar aspect of the hand) in ALS patients<sup>13</sup> suggests that it is at least plausible that upper limb-onset ALS favours the dominant hand through increased cortical connectivity. Furthermore, in light of the demonstration of cortical hyperexcitability as a feature of early sporadic ALS<sup>14</sup> and as a preceding event in familial ALS,<sup>15</sup> it is noteworthy that a transcranial magnetic stimulation study in relation to handedness in healthy participants demonstrated a shorter cortical silent period (reflecting reduced inhibitory function) in the dominant hand.<sup>16</sup>

A study of 21 cases of monomelic atrophy (a purely LMN syndrome) showed no relationship to handedness.<sup>17</sup> Although not ascertained in our study, it may be pertinent to explore handedness in relation to upper limb-onset ALS further with respect to the degree of UMN involvement clinically.

Finally, with reference to the proposal that ALS may be associated with previous athleticism, although left handedness has been noted to be higher among the sporting elite, reaching 50% in some disciplines,<sup>18</sup> we did not observe any increase in left handedness in ALS patients compared with the general population.

#### Limitations

The use of a questionnaire-based survey is potentially subject to responder and recall bias. In respect of responder bias, however, the demographics, the distribution of the frequencies of the sites of symptom onset and the sex ratio differences between bulbar and limb onset suggest that our responders are representative of the general ALS population. With respect to recall bias, it is likely that weakness will be noticed sooner in a dominant limb if both limbs are equally weak because of the greater awareness and higher use. While this cannot be completely excluded as an explanation for our findings, it would not be consistent with our overall observation of focal onset of disease. There is also a potential discrepancy between questionnaire and performance-based measures of footedness that may have confounded the results in relation to this.<sup>19</sup>

Prospective, clinic-based studies of handedness in relation to upper limb-onset ALS are needed to confirm our findings, and further research aimed specifically at understanding more about corticomotoneuronal connectivity might be revealing.

**Acknowledgements** We thank the Medical Research Council, Motor Neurone Disease Association of Great Britain and Ireland, the NIHR for Biomedical Research and those patients registered with http://Patientslikeme.com who took the time to return the questionnaire.

**Funding** MRT is supported by the MRC/MNDA Lady Edith Wolfson Clinician Scientist Fellowship. PW, CAB, and MPM are employees of PatientsLikeMe and own stock options in the company. PatientsLikeMe receives research support from UCB, Novartis, and Avanir.

#### Competing interests None.

Provenance and peer review Not commissioned; externally peer reviewed.

#### REFERENCES

- Haverkamp LJ, Appel V, Appel SH. Natural history of amyotrophic lateral sclerosis in a database population. Validation of a scoring system and a model for survival prediction. *Brain* 1995;118(Pt 3):707–19.
- Shoesmith CL, Findlater K, Rowe A, et al. Prognosis of amyotrophic lateral sclerosis with respiratory onset. J Neurol Neurosurg Psychiatry 2007;78:629–31.
- Rothstein JD. Current hypotheses for the underlying biology of amyotrophic lateral sclerosis. Ann Neurol 2009;65(Suppl 1):S3-9.
- Ravits J, Paul P, Jorg C. Focality of upper and lower motor neuron degeneration at the clinical onset of ALS. *Neurology* 2007;68:1571–5.
- Felmus MT, Patten BM, Swanke L. Antecedent events in amyotrophic lateral sclerosis. *Neurology* 1976;26:167–72.
- Kurtzke JF, Beebe GW. Epidemiology of amyotrophic lateral sclerosis: 1. A casecontrol comparison based on ALS deaths. *Neurology* 1980;30:453–62.
- Chio A, Benzi G, Dossena M, et al. Severely increased risk of amyotrophic lateral sclerosis among Italian professional football players. *Brain* 2005;128:472–6.
- Harwood CA, McDermott CJ, Shaw PJ. Physical activity as an exogenous risk factor in motor neuron disease (MND): a review of the evidence. *Amyotroph Lateral Scler* 2009;10:191–204.
- Scarmeas N, Shih T, Stern Y, et al. Premorbid weight, body mass, and varsity athletics in ALS. *Neurology* 2002;59:773–5.
- Hammond G. Correlates of human handedness in primary motor cortex: a review and hypothesis. *Neurosci Biobehav Rev* 2002;26:285–92.
- 11. Walsh RR, Small SL, Chen EE, et al. Network activation during bimanual movements in humans. *Neuroimage* 2008;43:540–53.
- 12. Eisen A. Amyotrophic lateral sclerosis-evolutionary and other perspectives. *Muscle Nerve* 2009;40:297–304.
- 13. Weber M, Eisen A, Stewart H, *et al*. The split hand in ALS has a cortical basis. *J Neurol Sci* 2000;**180**:66–70.
- Prout AJ, Eisen AA. The cortical silent period and amyotrophic lateral sclerosis. *Muscle Nerve* 1994;17:217–23.
- Vucic S, Nicholson GA, Kiernan MC. Cortical hyperexcitability may precede the onset of familial amyotrophic lateral sclerosis. *Brain* 2008;131:1540–50.
- Priori A, Oliviero A, Donati E, et al. Human handedness and asymmetry of the motor cortical silent period. Exp Brain Res 1999;128:390–6.
- 17. Gourie-Devi M, Gururaj G, Vasisth S, et al. Risk factors in monomelic amyotrophy: a case control study. NIMHANS J 1993;11:79–87.
- Raymond M, Pontier D, Dufour AB, et al. Frequency-dependent maintenance of left handedness in humans. Proc Biol Sci 1996;263:1627–33.
- Grouios G, Hatzitaki V, Kollias N, et al. Investigating the stabilising and mobilising features of footedness. *Laterality* 2009;14:362–80.

### Information for patients from JNNP

# Handedness may play a part in nerve disease onset

Researchers have discovered that you are more likely to first get symptoms of a type of motor neurone disease in the hand or arm you use more often.

### What do we know already?

Amyotrophic lateral sclerosis is a disease that kills off nerve cells. Eventually it can lead to paralysis. We don't know what causes it (although some cases are inherited), but scientists have speculated that it is linked in some way to intensive exercise in people with a genetic vulnerability.

Symptoms of ALS, such as muscle weakness and twitching, often start in the hands or feet. Researchers thought that, if ALS was linked to exercise, then symptoms would be more likely to start in the right hand if someone was right-handed and the left hand if they were left-handed.

So they surveyed a group of more than 500 people from a patient-support website, to see whether this was the case.

### What does the new study say?

People whose symptoms started in their hands or arms were likely to have had the symptoms start on the same side as their dominant hand (i.e., the right side for right-handed people and left side for left-handed). About 64 in 100 people reported this to have been the case.

When symptoms started in the feet or legs, there was no particular concordance. The researchers say this might be expected, as people tend to use both their lower limbs equally, for walking or standing, but their handedness affects how much their right or left arms or hands get used.

### How reliable are the findings?

Quite a few people in the study couldn't remember which side their symptoms started in, or gave more than one site for the beginning of the symptoms. The results rely on people remembering accurately where their symptoms began. It might also be the case that people were more likely to notice or remember when their dominant hand was affected – for example, you might not notice muscle weakness in your left hand, if you do most things with your right hand. Only when your right hand is affected might you realise there is a problem.

### Where does the study come from?

The study was done by researchers at the John Radcliffe Hospital in Oxford and Kings College Hospital, London.

### What does this mean for me?

If you have ALS, you might be curious to think back to where your symptoms started, and whether they were in your dominant or weaker side. But although these findings might help researchers work out the cause of ALS, they're unlikely to make much practical difference.

Turner MR, Wicks P, Brownstein CA, et al. Concordance between site of onset and limb dominance in amyotrophic lateral sclerosis. *J Neurol Neurosurg Psychiatry* 2011;**82**:853–4. http://jnnp.bmj.com/content/82/8/853.full

This summary was prepared by the staff of Best Health, BMJ Group's patient information service. This information does not replace medical advice. If you have a medical problem please see your doctor.

