Letters

Familial association of Von Hippel-Lindau complex and δ - β thalassaemia

SIR,—It is well known that the Von Hippel-Lindau complex in about 20% of the cases has a familial incidence and that it is inherited.³ Thalassaemia is also a genetically determined haematological disorder characterised by underproduction of at least one of the normal globin peptide chains of haemoglobin. We report the association of Von Hippel-Lindau complex and δ - β thalassaemia in a family composed of eight members, six of whom were affected by one or both diseases.

In September 1980, a 27-year-old man was admitted for treatment of a cerebellar haemangioblastoma. His mother was successfully treated by us for the same disease 10 years before. The tumour was totally removed and the patient discharged 10 days later

free of symptoms. The patient had moderate erythrocytosis, a low mean corpuscular volume and mean corpuscular haemoglobin, with normal mean corpuscular haemoglobin concentration. Haemoglobin studies showed a normal concentration of A₂ haemoglobin and moderate increase of F haemoglobin. δ - β thalassaemia was suspected and an haematologic familiar study confirmed the diagnosis. The incidence of both diseases in the different members of the family were studied (figure). The mother had both δ - β thalassaemia and cerebellar haemangioblastoma. The father was free of both diseases. One sister and one brother of the patient had only δ - β thalassaemia, one brother had a retinal angioma and the other brother had no disease. The only member of the next generation, the son of the brother with retinal angioma, also had a retinal angioma.

Both diseases in this family are inherited independently, so it is possible that the responsible genes are not contained in the chromosomal structures. As far as we know this is the first description of an association of Von Hippel-Lindau complex and δ - β thalassaemia in members of the same family.^{1,2} We think that the erythrocytosis observed in these patients was probably due not to abnormal secretion of erythropoietin but to the presence of abnormal haemoglobin.

References

- 1 Grossman M, Melmon K. Von Hippel-Lindau disease. In: Vinken PJ, ed. *Handbook of Clinical Neurology*. Amsterdam: North Holland Publishing; 14:241-259.
- 2 Jeffreys R. Pathological and haematological aspects of posterior fossa haemangioblastomata. *J Neurol Neurosurg Psychiatry* 1975; 38:112-9.
- 3 Russell D, Rubinstein L. Pathology of tumours of the nervous system. 4th ed. London: Edward Arnold Publisher 1977: 46, 117.

R MARTÍNEZ, J CABEZUDO,
J VAQUERO, E ARETIO
Department of Neurosurgery
L BARBOLLA
Department of Haematology

*Clinica Puerta de Hierro,
Autonomous University,
C/San Martín de Porres 4,
Madrid 35, Spain*

Figure
Pedigree of the family.