Matters arising

Aqueduct stenosis and schizophrenia

The following letter was inadvertently omitted from *J Neurol Neurosurg Psychiatry* 1983;46:1158. Part of the reply by Reveley and Reveley is reprinted.

Sir: Reveley and Reveley, in their recent paper in this Journal¹ state that "we could find no reports linking schizophrenia specifically with hydrocephalus." In fact Lying-Tunell² reported two cases of normal pressure hydrocephalus that had presented for decades as schizophrenia. Remarkably, both patients' psychosis was permanently alleviated by shunting. Price and Tucker,³ in a review of the psychiatric manifestations of normal pressure hydrocephalus, reported psychotic symptoms with fluctuating cognitive deficits as characteristic clinical features.

Reveley and Reveley also state that typically, hydrocephalic patients show better verbal performance than expected and that these patients may also be excessively talkative and emotionally labile ("the cocktail party syndrome"). It may be useful to add that normal pressure hydrocephalus is usually marked by impaired verbal memory⁴ and frank apathy.⁵ And finally, the authors suggest that in "some cases the ventricular enlargement seen in schizophrenia differs from hydrocephalus only in degree." Nyback *et al*⁶ have also postulated a similar, normal pressure hydrocephalus-like CSF disturbance in schizophrenics. However, the available evidence argues against these hypotheses. Hydrocephalus is almost always a progressive disease process.⁷ Weinberger and Wyatt⁸ found that the ventriculomegaly in schizophrenia was non-progressive, had a normal CSF pressure, and showed no evidence of a flow disturbance on isotope cisternography.

MANTOSH J DEWAN

PETER BICK

State University of New York,
Upstate Medical Center,
750 East Adams Street,
Syracuse, NY 13210, USA

References

- 1 Reveley AM, Reveley MA. Aqueduct stenosis and schizophrenia. *J Neurol Neurosurg Psychiatry*

Psychiatry 1983;46:18-22.

- 2 Lying-Tunell U. Psychotic symptoms in normal pressure hydrocephalus. *Acta Psychiatr Scand* 1979;59:415-9.
- 3 Price TR, Tucker GJ. Psychiatric and behavioral manifestations of normal pressure hydrocephalus. A case report and brief review. *J Nerv Ment Dis* 1977;164:51-5.
- 4 Berglund M, Gustafson L, Hagberg B. Amnesic-confabulatory syndrome in hydrocephalic dementia and Korsakoffs psychosis in alcoholism. *Acta Psychiatr Scand* 1979;60:323-33.
- 5 Fisher CM. The clinical picture in occult hydrocephalus. *Clin Neurosurg* 1976;24:270-84.
- 6 Nyback H, Weisel FA, Berggren FM, *et al*.: Computed tomography of the brain in patients with acute psychosis and in healthy volunteers. *Acta Psychiatr Scand* 1982;65:403-14.
- 7 Fitz CR, Rao K. The ventricular and subarachnoid spaces in children. In: Lee SH, Rao K, eds. *Cranial Computed Tomography*. New York: McGraw-Hill, 1982.
- 8 Weinberger DR, Wyatt RJ. Brain Morphology in Schizophrenia: In vivo studies. In: Henn F, Nasrallah H, eds. *Schizophrenia As a Brain Disease*. New York: Oxford University Press, 1982.

Reveley and Reveley reply

We are grateful to Dewan and Bick for bringing the reports of Lying-Tunell¹ and Price and Tucker² to our attention. In the light of these reports, we should revise our statement to say "we could find no reports linking schizophrenia specifically with aqueduct stenosis and hydrocephalus." For it is clear from the radiological and clinical description that Lying-Tunell does not report cases of aqueduct stenosis, just as we do not report cases of normal pressure hydrocephalus (NPH). For example, Case 1 of Lying-Tunell actually had "marked dilatation of Sylvian aqueduct" on pneumoencephalogram, while Case 2 had a "highly dilated ventricular system". Price and Tucker gave no details of the pneumoencephalogram, other than to say that it is "diagnostic for NPH". The size of the fourth ventricle was not specifically noted.

NPH characteristically develops in adulthood³ and is associated with demen-

tia, ataxia, and incontinence, as well as other symptoms less characteristic, as noted in the cases of Lying-Tunell and Price and Tucker. In contrast our cases 2 and 3 had onset of hydrocephalus in infancy, within a few weeks of birth, and developed skull enlargement. Case 1 had onset in adolescence, but did not develop any clinical signs of NPH. She had a difficult forceps delivery, which may have contributed to the development of aqueduct stenosis.

The questions of CSF flow disturbance and progression of ventricular enlargement in schizophrenia have yet to be fully explored. Weinberger and Wyatt's⁴ finding of non-progression was based on only five cases after 2-3 year follow up. Disturbances in CSF flow in schizophrenics have been found by Sedvall and associates⁵ using isotope cisternography.

Hydrocephalus is a condition with multiple aetiologies. It is unknown if any of these processes are linked to the ventricular enlargement occasionally occurring in functional psychoses, and this should be the subject of further research.

MICHAEL A REVELEY
ADRIAN M REVELEY
Genetics Section
Institute of Psychiatry
De Crespigny Park
London SE5 8AF, UK

References

- 1 Lying-Tunell U. Psychotic symptoms in normal-pressure hydrocephalus. *Acta Psychiatr Scand* 1979;59:415-9.
- 2 Price TR, Tucker GJ. Psychiatric and behavioural manifestations of normal pressure hydrocephalus. A case report and brief review. *J Nerv Ment Dis* 1977;164:51-5.
- 3 Fisher CM. A clinical picture in occult hydrocephalus. *Clin Neurosurg* 1977;24:270-84.
- 4 Weinberger DR, Wyatt RJ. Brain morphology in schizophrenia. In: Henn F, Nasrallah H, eds. *Schizophrenia as a Brain Disease*. New York: Oxford University Press, 1982:148-175.
- 5 Sedvall G. Evidence for alterations of brain morphology and function in subgroups of schizophrenic patients. *Abstracts of the 7th World Congress of Psychiatry*. 1983: p. 12.