cholinergic marker enzymes. *Neurosci Lett* 1982;**28**:217-22.

- ⁸ Henke H, Lang W. Cholinergic enzymes in neocortex, hippocampus and basal forebrain of non-neurological and senile dementia of Alzheimer-type patients. *Brain Res* 1983; **267**:281-91.
- ⁹ White P, Hiley CR, Goodhart MJ, et al. Neocortical cholinergic neurons in elderly people. *Lancet* 1977;*i*:668-71.
- ¹⁰ Davies P, Verth AH. Regional distribution of muscarinic acetylcholine receptors in normal and Alzheimer's type dementia brains. *Brain Res* 1978;**138**:385-92.
- ¹¹ Bowen DM, Spillane JA, Curzon G, et al. Accelerated ageing or selective neuronal loss as an important cause of dementia? *Lancet* 1979;*i*:11-14.
- ¹² Reisine TD, Yamamura HI, Bird ED, Spokes E, Enna SJ. Pre- and post-synaptic neurochemical alterations in Alzheimer's disease. *Brain Res* 1978;**159**:477-80.
- ¹³ Reisine TD, Fields JZ, Yamamura HI, et al. Neurotransmitter receptor alterations in Parkinson's disease. *Life Sci* 1977;**21**:335-44.
- ¹⁴ Rinne UK, Koskinen V, Lönnberg P. Neurotransmitter receptors in the parkinsonian brain. In: Rinne UK, Klingler M, Stamm G, eds. *Parkinson's Disease—Current Progress, Problems and Management*. Amsterdam: Elsevier, 1980:93-107.
- ¹⁵ Drachman DA, Leavitt J. Human memory and the cholinergic system. A relationship to ageing. *Arch Neurol* 1974;**30**:113-21.
- ¹⁶ Mishkin M. Memory in monkeys severely impaired by combined but not by separate removal of amygdala and hippocampus. *Nature* 1978;**273**:297-8.

Accepted 16 December 1983

Metastatic carcinoid tumour compressing the cauda equina

Sir: We report a patient with a progressive syndrome of cauda equina compression due to an osteogenic metastasis from an intestinal carcinoid tumour.

A 64-year-old female presented with a one year history of low back pain irradiating to both legs for three months. Prolonged sitting caused numbness in both thighs. Six months before she had been examined for these complaints at another hospital, and radiographs of the lumbar spine were reported normal. She had a ten year history of abdominal pain and intermittent diarrhoea and two years previously had presented with acute intestinal obstruction. A small bowel carcinoid was removed at operation. There were multiple metastases to the mesentery and abdominal lymph nodes. Physical examination showed bilateral grade 4 quadriceps weakness; the knee and ankle jerks on the right

side were reduced. Straight leg raising was normal on both sides but there was bilateral femoral nerve stretch pain. Cauda equina compression at the L3 level was presumed. Plain radiographs of the lumbar spine showed sclerosis of the body and the right pedicle of L3. Metrizamide myelography demonstrated a complete block at L3. CSF analysis including a smear of the sediment was unremarkable. At surgery a highly vascular tumour was found in the epidural space opposite L3. The tumour was clearly defined and did not seem to arise from bone, but a biopsy of the vertebral arch and body demonstrated carcinoid tissue in the spongiosa. The lumbar spine was irradiated with 6000 rads. Signs and symptoms disappeared and the patient remained well for 12 months. Postoperatively she complained of periods of flushing which reacted well to methysergide. She died 15 months later following multiple abdominal complications included intestinal obstructions.

Although carcinoid tumours were initially considered as non-infiltrating and non-metastasising,¹ subsequent case reports illustrated that carcinoid tumours may metastasise to the lymph nodes,^{2,3} and very rarely to the nervous system.^{4,5} To our knowledge only one well-documented case of cervical spinal cord compression has been published,⁶ and no cases with lumbar involvement producing cauda compression could be traced. When our patient presented with low back pain and inconstant radicular symptoms, the presence of degenerative bone disease or disc pathology was much more probable⁷⁻⁹ than metastatic carcinoid tumour. Plain radiographs of the spine six months previously were normal but at a later stage there were discrete abnormalities on plain radiographs of the lumbar spine. As osteoblastic metastases are well recognised complications of malignant carcinoid tumours, further examinations were performed and showed a metastasis at L3. This seems a very rare complication of carcinoid tumour. A neurofibroma had also to be excluded, because the combination of carcinoid and von Recklinghausen's neurofibromatosis had been stressed in two previous reports.^{10,11}

In conclusion, the patient illustrates that in presence of persisting spinal pain, radiculopathy or myelopathy, a vertebral metastasis should be considered even when the primary tumour is of carcinoid origin. As in the case of other bone metastases a bone isotope scan has to be performed even when plain films of the spine are nor-

mal¹² while also an immuno-assay technique with specific antibodies to serotonin may be a very sensitive method of carcinoid metastasis detection.¹³ These procedures may detect the lesion earlier, resulting in temporary pain relief by röntgen irradiation. Our patient remained pain free for one year.

BJJ ANSINK

HCE VAN LAMBALGEN

Department of Neurology

G DAVIES

Department of Neuroradiology,

Sint Lucas Ziekenhuis,

Jan Tooropstraat 164,

1061 AE AMSTERDAM, The Netherlands

References

- ¹ Oberndorfer SS. Karzinoide Tumor des Dünndarms. *Frankfurter Zeits Pathologie* 1907;**1**:426-37.
- ² Gosset A, Masson P. Tumeurs carcinoides et métastases mésentériques. *Press Méd* 1914;**22**:237-9.
- ³ Paevy PW, Rogers JV, Clement JL, Burns JB. Unusual osteoblastic metastases from carcinoid tumors. *Radiology* 1973;**107**:327-30.
- ⁴ Wagner PK. Magenkarzinoid mit Hirnmetastase. *Med Klin* 1979;**74**:1439-43.
- ⁵ Fishman ML, Rosenthal S. Optic nerve metastasis from a mediastinal carcinoid tumor. *Br J Ophthalmol* 1976;**60**:583-8.
- ⁶ Manoli RS, Barteley CR. Osteolytic and osteoblastic metastases due to carcinoid tumors. *Cl Nucl Med* 1980;**5**:102-5.
- ⁷ Gathier JC. Radicular disorders due to lumbar discopathy. In: Vinken PJ, Bruyn GW eds. *Handbook of Clinical Neurology*, vol. 29: 573-604. Amsterdam: North Holland, 1976.
- ⁸ Guidetti B, Fortuna A. Differential diagnosis of intramedullary and extramedullary tumours. In: Vinken PJ, Bruyn GW eds. *Handbook of Clinical Neurology*, vol. 19: 51-75. Amsterdam: North Holland, 1975.
- ⁹ Levitt P, Ranshoff J, Speilholz N. The differential diagnosis of tumors in the conus medullaris and cauda equina. In: Vinken PJ, Bruyn GW eds. *Handbook of Clinical Neurology*, vol. 19: 77-90. Amsterdam: North Holland, 1975.
- ¹⁰ Trouillas P, Robert JM. Maladie de Recklinghausen et phacomatoses de dysplasie néoplasiantes frappant le système APUD. *Nouv Press Méd* 1974;**3**:2399-401.
- ¹¹ Alamowitch Ch, Larousse C. Association de la maladie de Recklinghausen et carinoide de l'ampoule de Vater. *Nouv Press. Méd* 1982;**11**:3255-7.
- ¹² Toomey B, Felson B. Osteoblastic bone metastases in gastrointestinal and bronchial carcinoids. *Am Roentgen Ray Soc (Cincinnati, Ohio)* 1959;22-5.
- ¹³ Cuello AC. Serotonin immunoreactivity in carcinoid tumours demonstrated by a monoclonal antibody. *Lancet* 1982;*i*:771-83.

Accepted 22 December 1983