References

- ¹ Earl CJ, Fullerton PM, Wakefield GS, Schutta HS. Hereditary Neuropathy with liability to pressure palsies. *Q J Med* 1964;33:481-98.
- ² Oh SJ, Sarala PK, Kuba T, Elmore RS. Tarsal tunnel syndrome: electrophysiological study. *Ann Neurol* 1979;5:327-30.
- ³ Fu R, Delisa JA, Kraft GH. Motor nerve latencies through the tarsal tunnel in normal adult subjects: standard determinations corrected for temperature and distance. *Arch Phys Med Rehab* 1980;61:243-8.
- ⁴ Dekel S, Coates R. Primary carpal stenosis as a cause of "idiopathic" carpal tunnel syndrome. *Lancet* 1979;i:1024.
- ⁵ Lam SJS. Tarsal tunnel syndrome. *J Bone Jt Surg* 1967;49B:87-92.

Accepted 2 December 1985

Alteration of the visual blink reflex in patients with dementia

Sir: The clinical diagnosis of senile dementia of the Alzheimer type and multi-infarct dementia can be very difficult.^{1,2} Neurophysiological tests are not specific. In a recent study, the EEG and visual evoked potentials were reported to be normal in respectively 34% and 76% of patients with Alzheimer type dementia.³ Therefore, the development of additional sensitive tests which could discriminate dementia from other disorders, would be of great value.

We have found a simple neurophysiological test, the visual blink reflex, to be abnormal in patients with Alzheimer type dementia or multi-infarct dementia. The visual blink reflex consists of a reflex contraction of the eyelids in response to a bright light which is flashed in front of the eyes of the subject. Normal subjects usually show a visual blink reflex with a constant latency of approximately 50 ms, although the visual blink reflex can be absent in up to 12% of normals.⁴ The visual blink reflex is a subcortical reflex but the exact pathway in the brainstem is unknown.^{5,6}

Fifteen patients with senile dementia (Alzheimer type dementia or multi-infarct dementia) (mean age 77.2 ± 7.7 yr) were examined according to the method described by Malin.⁴ The EMG activity of the orbicularis oculi muscles was recorded with surface skin electrodes. Patients were not informed that the blink reflex was measured in order to avoid voluntary blinking. In each subject we determined the average latency time of the visual blink reflex from 10 trials. The controls were 13 normal aged persons (mean age 70.5 ± 11 yr).

Twelve out of 15 patients showed a symmetrical visual blink reflex with a mean latency for the group of 104.3 ± 30.0 ms (range 65-158 ms). In three patients no visual blink reflex could be elicited. The visual blink reflex of the group of normal subjects had a mean latency of 49.7 ± 2.2 ms (range 47-53.3 ms). The visual blink reflex was absent in one control. The mean latencies of the patient and the control group differed significantly ($p < 0.001$).

Our study demonstrates that the latency of the visual blink reflex of patients with senile dementia is markedly increased. Possibly, the alteration of the visual blink reflex in senile dementia is due to lesions or functional disturbances at the level of the brainstem, which are known to occur in Alzheimer type dementia.⁷ In our view these findings imply that further studies on the visual blink reflex in dementia are warranted.

DLJ TAVY
TCAM VAN WOERKOM

HHE MORRE

JPJ SLAETS

Municipal Hospitals, The Hague,
Departments of Clinical
Neurophysiology and Clinical
Neurology and Psychiatric Centre
Rosenburg, The Hague,
Departments of Neuropsychiatry
and Psychogeriatrics
2545 CH, The Netherlands

References

- ¹ Ron MA, Toone BK, Garalda ME, Lishman WA. Diagnostic accuracy in presenile dementia. *Br J Psychiatry* 1979;134:161-8.
- ² Caine ED. Pseudodementia. *Arch Gen Psychiatry* 1981;38:1359-64.
- ³ Visser SL, Van Tilburg W, Hooijer C, Jonker C, De Rijke W. Visual evoked potentials (VEPs) in senile dementia (Alzheimer type) and in non organic behavioural disorders in the elderly; comparison with EEG parameters. *Electroencephalogr Clin Neurophysiol* 1985;60:115-21.
- ⁴ Malin JP. Der optisch evozierte orbicularis oculi und reflex: methodik, normalwerte, habituation. *Z EEG-EMG* 1982;13:101-7.
- ⁵ Brierly JB, Graham DI, Adam JH, Simpson JA. Neocortical death after cardiac arrest. *Lancet* 1971;2:560-5.
- ⁶ Tavy DLJ, Van Woerkom TCAM, Bots GTAM, Endtz LJ. Persistence of the blink reflex to sudden illumination in a comatose patient. *Arch Neurol* 1984;41:323-4.
- ⁷ Terry RD, Katzman R. Senile dementia of the Alzheimer type. *Ann Neurol* 1983;14:497-506.

Accepted 10 May 1985

Isolated lower motoneuron involvement following radiotherapy

Sir: Progressive transverse myelopathy which follows, after a characteristic latent period, the radiation of tumours in the vicinity of the spinal cord is a well-recognised sequel of radiotherapy.¹ Another myelopathic post-radiation syndrome is characterised by a slowly evolving amyotrophy with paresis and areflexia of legs without sphincter disturbances or sensory impairment, and is presumed to be the result of selective damage to the anterior horn cells in the lumbo-sacral spinal cord.²⁻⁴

Our case, a 64-year-old male, had a right-sided pheochromocytoma, encasing the vena cava and the celiac axis. Surgical removal was only partial, so he received 5800 rads to the affected side (December 1983). Seven months after the treatment, he suffered painful cramps in the legs for two weeks, followed by progressive bilateral leg weakness. There were no sphincter abnormalities or sensory symptoms. The course of illness was one of initial worsening followed by a stabilised state until our observation in May 1985. Neurological examination showed diffuse lower limb wasting, more marked in gluteal muscles, and moderate weakness of the peronei, extensors of the feet and toes and gluteal muscles. Fasciculations were present in the calf and gluteal muscles. Tendon reflexes were absent in the legs and plantar responses were flexor. Sensation was intact. Laboratory evaluations were normal as well as radiographs of the thoracic, lumbar and sacral spine. Magnetic resonance imaging of the thoracolumbar spinal cord was normal, and there was no evidence of intraspinal metastasis. Quantitative EMG⁵ of the left vastus medialis, peroneus longus, gastrocnemius medialis and lateralis and gluteus maximus showed denervation activity with fasciculations and loss of motor units with pronounced signs of reinnervation in all muscles. Motor conduction velocity along the left peroneal and posterior tibial nerves was reduced by 24% and 14%, and distal latencies were prolonged by 51% and 40% respectively. Amplitudes of motor responses were normal. Amplitude and shape of SAPs sensory thresholds and sensory conduction velocity along the left peroneal and sural nerves were normal. Average latencies were prolonged by 22% and 10% along the left peroneal and posterior tibial nerves respectively; The left H reflex had a prolonged latency, a markedly polyphasic shape, and was constantly associated with late waves at 72 ms. SEPs were obtained in the upper and lower limbs on the