Matters arising

- ³ Love JG, Schorn VG. Thoracic disc protrusions. *JAMA* 1965;191:627-31.
- ⁴ Logue V. Thoracic intervertebral disc prolapse with spinal cord compression. *J Neurol Neurosurg Psychiatry* 1952;15:227-41.
- ⁵ Arseni C, Nash F. Thoracic intervertebral disc protrusion. A clinical study. *J Neurosurg* 1960;17:418-30.
- ⁶ Fisher RG. Protrusions of thoracic disc. The factor of herniation through the dura mater. *J Neurosurg* 1965;6:591-3.
- ⁷ Tovi D, Strang RR. Thoracic intervertebral disc protrusions. *Acta Chir Scand* 1960;267:41.
- ⁸ Crafoodi C, Hierton T, Lindblom K, Olsson S-E. Spinal cord compression caused by a protruded thoracic disc. Report of a case treated with antero-lateral fenestration of the disc. *Acta Orthop Scand* 1958;28:103-7.
- ⁹ Patterson RHJ, Arbit E. A surgical approach through the pedicle to protruded thoracic disks. *J Neurosurg* 1978;48:768-72.
- ¹⁰ Perot P, Munro DD. Transthoracic removal of midline thoracic disc protrusions causing spinal cord compression. *J Neurosurg* 1969;31:452-8.
- ¹¹ Ranshoff J, Spencer F, Siew F, Gage LJ. Transthoracic removal of thoracic disc. Report of three cases. *J Neurosurg* 1969;31:459-61.
- ¹² Sebban LN, Jannetta PJ. Thoracic disc herniation, operative approaches and results. *Neurosurgery* 1983;12:303-5.
- ¹³ Simeone FA. The modern treatment of thoracic disc disease. *Orthop Clin North Am* 1971;2:453-62.

The relationship of peripheral trauma and pain to dystonia

Sir: Schott¹ has recently reported four subjects in whom minor peripheral injury was responsible for the development of segmental dystonia. Although the mechanisms underlying trauma-induced dystonia are not known, activation of endogenous endorphins and adrenocorticotropin hormone (ACTH) might be involved. All patients experienced severe pain, sufficient to have activated central endogenous endorphins and ACTH.² The endogenous endorphins and ACTH² have been shown to be involved not only with the regulation of or reaction to pain, but also a wide range of motor and behavioural responses in laboratory animals.³⁻⁶

However, we are not aware of any report that endogenous administration of endorphins or other opioids produced dystonic movements. In fact, in one study⁵ morphine or beta-endorphin injected directly into the brainstem of rats caused catalepsy and rigidity but not dystonic movements. However, ACTH N-terminal fragments, but not

ACTH itself, administered in the same manner produced postural asymmetry and dystonic movements resembling human dystonia.^{5,6} Jacquet and Abrams^{5,6} have suggested that some forms of human dystonia may be related to a genetic abnormality of the ACTH molecule. In the cases reported by Schott¹ it is possible that the patients may have had an underlying mutation in the structure of the ACTH molecule. Thus, it is conceivable that the pain and associated stress of the patients activated the cerebral production and/or release of this abnormal compound leading to the development of the segmental dystonia. Although this mechanism is a conjecture, it is of interest that exogenously administered ACTH was reported to ameliorate symptoms of torsion dystonia in one patient.⁷

Taken together it appears that abnormalities of the molecular structure of ACTH may be implicated, at least in part, in the pathophysiology of human dystonia. We are currently studying this issue.

(Supported in part by NINCDS grant NS15441 and the Muscular Dystrophy Association).

REUVEN SANDYK*

PAUL CONSROE†

Departments of Internal Medicine
(Neurology)* and
Pharmacology/Toxicology*†
University of Arizona,
Tucson, AZ 85724, USA

References

- ¹ Schott GD. The relationship of peripheral trauma and pain to dystonia. *J Neurol Neurosurg Psychiatry* 1985;48:698-701.
- ² Haynes RC, Murad F. Adrenocorticotrophic hormone: adrenocorticosteroids and their synthetic analogs: inhibitors of adrenocortical biosynthesis. In: Gilman AG, Goodman LS, Rall TW, Murad F, eds. *The Pharmacological Basis of Therapeutics*. 7th ed. New York: Macmillan Publishing Co, 1985:1459-89.
- ³ Sandyk R. The endogenous opioid system in neurological disorders of the basal ganglia. *Life Sci* 1985;37:1655-63.
- ⁴ Grillner S, Zangger P. How detailed is the central pattern generation for locomotion. *Brain Res* 1975;88:367-71.
- ⁵ Jacquet YF, Abrams GM. Postural asymmetry and movement disorder after unilateral microinjection of adrenocorticotropin 1-24 in rat brainstem. *Science* 1982;218:175-7.
- ⁶ Jacquet YF. "Dystonia"-like postural asymmetry after microinjection of ACTH N-terminal fragments but not after ACTH₁₋₃₉ in rat brainstem suggests role of neuropeptide mutation in genetic movement disorder. *Brain Res* 1984;294:144-7.
- ⁷ Baumann L, Baumann J. Betrachtungen zu einem Fall von Torsionsdystonie. *Acta Neurol Scand* 1963;39:Suppl 14:221-5.

Transient global amnesia after whiplash trauma

Sir: We are pleased to see the letter from Hofstad and Gjerde on whiplash amnesia.¹ Their note strengthens our opinion that whiplash amnesia is probably a particular form of transient loss of short-term memory.² Whiplash amnesia seems to present characteristic features as the reported cases^{1,3} show. For instance, selective pain in the neck and dizziness are not found in transient global amnesia and definite retrograde amnesia is rarely absent in amnesia by concussion. We thus appreciate the work of Hofstad and Gjerde and we think that the description of clinical features in newly diagnosed cases of whiplash amnesia confirm our opinion.

I BUENAVENTURA

J MATIAS-GUIU

The Neurology Service,
Hospital del Valle de Hebrón,
Universidad Autonoma Barcelona,
Spain

References

- ¹ Hofstad H, Gjerde IO. Transient global amnesia after whiplash trauma. *J Neurol Neurosurg Psychiatry* 1985;48:956-7.
- ² Matias-Guiu J, Buena Ventura I, Cervera C, Codina A. Whiplash amnesia. *Neurology (Cleveland)* 1985;35:1259.
- ³ Fisher CM. Whiplash amnesia. *Neurology (NY)* 1982;32:667-8.

Free light chains in the cerebrospinal fluid

Sir: My letter concerns the article "Free light chains in the cerebrospinal fluid: an indicator of recent immunological stimulation", published in your journal 1985;48:995-8.

I do not agree with the following statements of the authors since they misinterpret the findings of our laboratory:

- (1) "... Bollengier *et al* suggested that the free light chains occurred in CSF..." Nowhere in our papers did we suggest the presence of free light chains; the word "suggest" refers to a mere hypothesis and in fact we clearly demonstrated and quantified those free light chains in radial immunodiffusion by using a specific anti free light chain antiserum (ref. 9 in the article by Vakaet and R. Thompson)
- (2) "... antiserum directed against Bence-Jones protein which had been previously adsorbed with heavy-chains..." We did no such thing for the very reason