and Down's syndrome cases, data from all cases were pooled for analysis. The number of tangles in the aqueous homogenates varied enormously from brain to brain (range 14–325, median 102/mg wet weight) and were not related to age, sex, duration of dementia or post-mortem interval. For each brain the number of tangles after extraction with SDS or NaOH was expressed as a percentage of the number of tangles in the aqueous (control) homogenate. The percentage of tangles insoluble in 1% SDS ($57 \pm 25\%$, mean $\pm$ SD), 5% SDS ($43 \pm 17\%$) and 0.2M NaOH ($37 \pm 22\%$) was not significantly related to age, sex, or post-mortem interval. There was, however, a significant negative correlation ($r = -0.57$, $p < 0.02$) between the percentage of tangles insoluble in 1% SDS and duration of dementia: that is, tangle solubility in 1% SDS increased with increasing duration of dementia (fig). This relationship was also significant ($r = -0.62$, $p < 0.05$) when only the Alzheimer-type dementia cases were considered. Duration of dementia was not significantly correlated with the percentage of tangles insoluble in 5% SDS or 0.2M NaOH. Plaque cores were essentially insoluble in all three agents.

A wide variation in tangle counts between Alzheimer-type dementia brains has been

reported using homogenates² and tissue sections.³ Although younger cases of Alzheimer-type dementia usually have more tangles than older cases,⁴ we observed no correlation between age and tangle counts, possibly because our cases were selected on the basis of severe pathology. The fact that the percentage of tangles soluble in 1% SDS increased with increasing duration of dementia suggests that in any one brain with severe Alzheimer-type dementia, tangles become more soluble and less stable with the passage of time. This enhanced solubility may reflect transformation of intracellular tangles to extracellular "ghost" tangles.⁵

For the supply of post-mortem brain tissue, we thank clinicians and pathologists at Bangour General, Bangour Village and Gogarburn Hospitals, West Lothian; Strathmartine and Ninewells Hospitals, Dundee; the Royal Edinburgh Hospital.

S HUSSEY*

CM YATES*

JE CHRISTIE*

RA ELTON†

A GORDON‡

MRC Brain Metabolism Unit,*

Edinburgh University Department of

Pharmacology,

1 George Square,

References

- Tomlinson BE, Corsellis JAN. Ageing and the dementias. In: Hume Adams J, Corsellis JAN, Duchen LW, eds. *Greenfield's Neuropathology*. London: Edward Arnold, 1985:958–66.
- Iqbal K, Zaidi T, Thompson CH, et al. Alzheimer paired helical filaments: bulk isolation, solubility and protein composition. *Acta Neuropathol (Berl)* 1984;62:167–77.
- Wilcock GK, Esiri MM, Bowen DM, et al. Alzheimer's disease. Correlation of cortical choline acetyltransferase activity with the severity of dementia and histological abnormalities. *J Neurol Sci* 1982;57:407–17.
- Tomlinson BE. Plaques, tangles and Alzheimer's disease. *Psychol Med* 1982;12:449–59.
- Rasool CG, Abraham C, Anderton BH, et al. Alzheimer's disease: Immunoreactivity of neurofibrillary tangles with anti-neurofilament and anti-paired helical filament antibodies. *Brain Res* 1984;310:249–60.

Accepted 13 September 1986

Matters arising

Limb apraxia in patients with damage confined to the left basal ganglia and thalamus

Sir: We read with interest the important paper by De Renzi *et al*¹ on limb apraxia caused by a left basal ganglia and thalamic lesion. In a previous publication² we described in detail four cases of moderate/severe apraxia caused by small left subcortical strokes. In our series involvement of the basal ganglia was not a prominent finding. Instead the lesions appeared on the higher CT cuts, lateral to the body of the lateral ventricles, involving the periventricular white matter. Lesions in this location damage both longitudinal white matter tracts (superior occipito frontal and superior longitudinal fasciculus) and the anterior callosal radiations. In De Renzi *et al*'s cases 2

and 3, CT scans showed infarcts involving the white matter adjacent to the left ventricle. In case 4 the lesion also extended to upper CT slices, while in both haemorrhagic cases 1 and 5, longitudinal white matter tracts might be functionally disturbed by edema and pressure from the haematoma. Damage to the periventricular white matter causes apraxia by disconnecting the parietal from the frontal lobe and preventing visuokinaesthetic motor patterns from the parietal lobe from contributing to the premotor cortex in programming correct movements.

This comment does not imply denying a role to the basal ganglia on the execution of learned movements. In fact they are also important in gesture recognition,³ suggesting that spatio-temporal engrams of common and over-learned gestures may be stored on the basal ganglia.

JOSÉ M FERRO

Neurobehavioral Unit,

Centro de Estudos Egas Moniz,

Department of Neurology,

Hospital Santa Maria,

Lisboa,

Portugal

References

- De Renzi E, Faglioni P, Scarpa M, Crisi G. Limb apraxia in patients with damage confined to the left basal ganglia and thalamus. *J Neurol Neurosurg Psychiatry* 1986;49:1030–8.
- Kertesz A, Ferro JM. Lesion size and location in ideomotor apraxia. *Brain* 1984;107:921–33.
- Ferro JM, Martins IP, Mariano G, Castro-Caldas A. CT scan correlates of gesture recognition. *J Neurol Neurosurg Psychiatry* 1983;46:943–52.

De Renzi *et al* reply:

We agree with Dr Ferro's claim that white matter damage may cause apraxia by interrupting the tracts linking the posterior regions of the brain with the premotor cortex. We doubt, however, that this was the case in our patients. As indicated by Dejerine,¹ (figs 219–23), the arcuate fascicle runs above the level at which our CT scan upper cut was taken and is located between the external and the extreme capsule in the white matter underlying the sylvian operculum, namely, more laterally with respect