- active multiple sclerosis: weekly analysis over a six month period. *Ann Neurol* 1983;13:418–25.
- 16 Bach M-A, Martin C, Cesaro P, Eizenbaum JF, Degos JD. T cell subsets in multiple sclerosis—a longitudinal study of exacerbating remitting cases. *J Neuroimmunol* 1985;7:331–43.
 - 17 Hughes PJ, Kirk PF, Compston DAS. Suppressor T cells in family members of patients with multiple sclerosis. *Brain* 1986;109:969–85.
 - 18 Craig JC, Hawkins SA, Swallow MW, *et al.* Subsets of T lymphocytes in relation to T lymphocyte function in multiple sclerosis. *Clin Exp Immunol* 1985;61:548–55.
 - 19 Oger J, Kastrukoff L, O'Gorman M, Paty DW. Progressive multiple sclerosis: abnormal immune functions in vitro and aberrant correlation with enumeration of lymphocyte subpopulations. *J Neuroimmunol* 1986;12:37–48.
 - 20 Antel JP, Bania MB, Reder A, Cashman N. Activated suppressor cell dysfunction in progressive multiple sclerosis. *J Immunol* 1986;137:137–41.
 - 21 Antel JP, Nicholas MK, Bania MB, Reder AT, Arnason BGW, Joseph L. Comparison of T8+ cell mediated suppressor and cytotoxic function in multiple sclerosis. *J Neuroimmunol* 1986;12:215–24.
 - 22 Chofflon MM, Weiner HL, Hafler DA. Loss of functional suppression is linked to decreases in circulating suppressor inducer (CD4+ 2H4+) T cells in multiple sclerosis. *Neurology* 1987;37(suppl 1):294.
 - 23 Trotter JL, Rodey GE, Gebel HM. Azathioprine decreases suppressor T cells in patients with multiple sclerosis. *N Engl J Med* 1982;306:365–6.
 - 24 Oger JJ-F, Antel JP, Kuo HH, Arnason BGW. Influence of azathioprine (Imuran) on in vitro immune function in multiple sclerosis. *Ann Neurol* 1982;11:177–81.

From the Archives of Surgery¹ Sir Jonathan Hutchinson's (1829–1913) modern description of symmetrical temporal arteritis.

“The subject ... was an old man ... the father of a well-remembered beadle at the London Hospital College 30 years ago ... I was asked to see him because he had red ‘streaks on his head’ which were painful and prevented his wearing his hat. The ‘red streaks’ proved ... to be his temporal arteries which ... were ... inflamed and swollen. Pulsation could be feebly detected in the affected vessel, but it finally ceased; the redness then subsided, and the vessels were left impervious cords. The old gentleman lived, I believe, several years after this without any other manifestation of arterial disease.”

RT ROSS, MD

Reference

- 1 Hutchinson J. Diseases of the Arteries. *Arch Surg* 1890;1:323–33.