The temporal sequence of aura-sensations in patients with complex focal seizures with particular attention to ictal aphasia

K KANEMOTO, D JANZ

From The Department of Neurology, University Hospital Rudolf-Virchow, Free University of Berlin, Berlin, Federal Republic of Germany

SUMMARY The sequences of aura sensations in 143 patients with complex partial seizures, were analysed with special emphasis on aphasic symptoms. Anxiety, epigastric sensation and visual hallucination were experienced early in the course of the aura, while illusion of familiarity and aphasia occurred late in the course of the aura. Three groups of interconnections of aura sensations were found which corresponded possibly to the types of seizure constellations proposed by Wieser. Close interconnections between impairment of verbal comprehension during seizures and paroxysmal thought disorder, as well as between paroxysmal paraphasia and illusion of familiarity were noted. Paroxysmal aphasia in patients with complex partial seizures was characterised as a positive symptom in contrast to stable aphasia.

The phenomenology of epileptic aphasia has been intensively studied by earlier authors.¹⁻¹³ Nevertheless, except for McKeaver *et al*¹⁴ and some sporadic case reports,¹⁵⁻²² aphasic phenomena during seizures have not attracted much attention recently. The statement of Critchley that epileptic aphasia is dispensable as a source of information to elucidate the structure of language disorder has not yet lost its influence.²³

We believe, however, that the temporal sequence of aura sensations within the aura could reveal dynamic linkage between various brain functions. This follows the principle pioneered by Janz²⁴ that this sequence follows definite patterns. As Castell has already pointed out,⁵ the study of ictal aphasia and aura sequences could shed a new light on interconnections between the language system and other brain functions by supplementing the deficiencies of "static" neuropsychology.

Because few authors have paid attention to the chronological sequences of aura sensations within the aura,²⁴⁻²⁷ we have tried to obtain a comprehensive survey of interrelations between aura sensations. We have also studied the specific position of epileptic aphasia from the viewpoint of aura sequences.

Address for reprint requests: Dr Kousuke Kanemoto, Utano National Hospital, Ukyo-ku, Narutaki, Ontoyama-cho 8, Kyoto-fu, Kyoto-shi, Japan

Received 10 May 1987 and in revised form 4 August 1988.
Accepted 8 August 1988

Materials and methods

We analysed auras of 143 patients with complex focal seizures retrospectively from the case notes in our clinic between 1973 and 1987. Patients directly treated by one of the authors (D.J.) were selected for our study because adequate and homogeneous documentation of auras were indispensable. Among these, 24 patients complained of aphasic symptoms during the aura (table 1).

Aphasia is defined in our study as follows: Impairment of comprehension, paraphasia, or word finding difficulty. Speech arrest and vocalisation without other signs of speech

Table 1 Frequency of different aura-sensations in our patients

	N	%
Epigastric	50	35
Familiarity (Déjà vu)	45	31
Aphasia	28	19
Anxiety	24	17
Thought-disorder	23	16
"Es"	19	13
Visual	16	10
Somatosensory	12	8
Gustatory	12	8
Porroopia-Porracousia	8	6
Olfactory	7	5
Auditory	7	5
Dysarthria	4	3
Others	5	4
No aura	31	22
Total N	143	

disturbance are not counted as ictal aphasia, in accordance with earlier studies suggesting their topographical and clinical independence from aphasic symptoms.^{8,9,22,29} Besides ictal aphasia, some other aura sensations should also be explained. "Es"²⁴ is non-specific warning difficult to express otherwise. "Es" in German corresponds approximately to "it" in English. We call the aura sensation "Es", if the experience can not be put into more precise terms. Patients can only say "It is a strange, indescribable feeling." *Epigastric sensation* refers to sensations rising from epigastric area. With *Porropsia* and *Porrakousia* the objects seem to move away from the patient or sounds seem to come from a distant place. In *thought disorder* there is crowding of thought, forced-thinking² and other changes in the quality of thought-process (for example, acceleration of thought). *Illusion of familiarity*: Besides déjà vu and jamais vu, this includes all the experiences of altered familiarity to the environment. Because déjà vu or jamais vu indicates visual familiarity literally, we adopt illusion of familiarity as a wider concept applicable to other modalities of sensations as well as visual.

Aura sequences were reconstructed on the basis of the patient's own report supplemented by the statements of witnesses. The same type of aura sequence in a patient was regarded as a unit. Eighty-one patients had only one type of aura sequence. Twenty-six patients had two types of aura sequences and five had more than two types. Thirty-one patients were recorded altogether ($31 \times 0 + 81 \times 1 + 26 \times 2 + 2 \times 3 + 2 \times 4 + 1 \times 5$).

EEG foci were determined interictally through multiple surface recordings.

Results

(A) Content and frequency of aura sensations

Frequencies of different aura sensations in our study are summarised in table 1. Epigastric aura and illusion of familiarity were the most frequent. Ictal aphasia, anxiety and thought-disorder followed them by a wide margin.

Among auditory hallucinations, we recorded six cases of elementary (machine sounds etc.) and one case of complex type (voice). Half of the visual hallucinations were elementary (coloured spot etc.) and half were complex (scene or picture). With regards to the ictal thought-disorder, five patients reported acceleration of thought ("The thought accelerates itself more and more rapidly, as if it began to gallop"). It was similar to crowding of thought, but the accelerated thought here lacked the strangeness characteristic of crowding of thought. Another six patients complained of a kind of forced thinking ("Thoughts push their way in my head and I cannot get rid of them"). This forced thinking was often reported to have an iterative nature ("It is always the same idea that comes and goes in my head"), but ordinarily they could not formulate what it was. Furthermore, a variety of altered thought-processes was experienced ("Thoughts blended into one another," "I could think only slowly and all ideas went out of me," "The thought was hardened as if it

Table 2 Distribution of laterality of EEG focus and type of aura sensation

	Left	Right	Both	No	Total
Epigastric	22	13	5	10	50
Familiarity (déjà vu)	16	12	6	11	45
Aphasia	8	8	3	9	28
Anxiety	16	1	2	5	24
Thought-disorder	9	5	3	6	23
"Es"	14	0	2	3	19
	5	7	1	3	16
Total	49	34	11	49	143

Only aura sensations occurring in more than 10% of patients are listed.

were a material"). Ictal aphasia consisted of eight cases with impairment of comprehension without paraphasia, five cases of paraphasia with simultaneous impairment of comprehension, nine cases of paraphasia with intact comprehension and two cases with difficulty in finding words. It should be noted also that the paraphasias reported or witnessed in our study contained neologisms in most cases.

(B) Correlation between laterality of EEG foci and type of aura (table 2)

Among our 143 patients, 34% had only left foci; 24%, only right foci. In comparison, left foci predominated in patients with ictal aphasia (67%) and ictal thought disorder (73%). Application of the χ^2 test indicated a significant association between these two aura sensations and a focus in the left cerebral hemisphere ($\chi^2 = 11.5$, 15.43 respectively, $p = 0.05$). In the patients with thought disorder, 14 had temporal EEG foci and two had extratemporal foci. We did not find a predominance of right foci among patients suffering from illusion of familiarity. Application of the χ^2 test fails to suggest any significant associations between aura sensations listed in table 2 and a focus in the right hemisphere.

(C) The sequence of aura sensations

We examined the distribution of auras with only one and more than one aura sensation. Especially "Es"-aura (69%) but also epigastric sensations (55%), olfactory sensations (57%) and somatosensory sensations (58%) tended not to be combined with other aura sensations. On the contrary, anxiety, gustatory sensations, thought disorder, ictal aphasia, dysarthria and visual hallucinations manifested themselves mostly in combination with other aura sensations (more than 80%).

Table 3 shows the time sequence of aura sensations within auras with more than one aura sensation. More than 70% of illusion of familiarity, ictal aphasia, dysarthria and porropsia-porracousia occurred at the end of the aura sequence. "Es"-aura, anxiety and epigastric sensation were almost exclusively found at

have not been pointed out previously. First, the frequency of thought disorder is strikingly high in our study (13%), especially considering that previous authors³⁰⁻³² did not adopt it as an independent item. Weber *et al.*²⁶ who studied aura psychopathologically, did pay attention to thought disorder, but did not note the predominance of left foci in thought disorder. Unlike previous authors, we included patients with extratemporal foci as well as those with temporal foci in our study. This alone, however, can not explain the discrepancy between our study and the previous studies, because the current series contained only a limited number of patients with extratemporal EEG foci in patients with thought disorder.

The predominant side of EEG foci in patients with déjà vu has been long disputed. In contrast to Mullan *et al.*³³ and Gupta *et al.*³⁴ as well as Cole *et al.*³⁵ did not find a predominance of right EEG foci among patients suffering from déjà vu.

Some of the rules about the sequences of aura sensations have been already reported in previous studies.^{24 25 27} Janz²⁴ and Paillas *et al.*²⁵ pointed out that visual hallucination is experienced early in the course of aura sequences. Janz,²⁴ and Hallen²⁷ mentioned the precedence of anxiety and epigastric sensations to the gustatory and olfactory sensations. The finding that illusion of familiarity occupies the endposition of an aura sequence was described by Janz.²⁴ However, previous studies have not examined the positions occupied by aphasia and thought disorder within aura sequences. Nor have they performed a comprehensive survey of aura sequences in patients with complex focal seizure. To offer such a survey, interrelations between main aura sensations including thought disorder and aphasia are schematically redrawn in fig 2. Three separate flows of aura sequences can be seen in the figure. After starting from anxiety, the flow going through visual hallucination does not join other aura sensations before reaching illusion of familiarity. Within the group of aura sensations starting from epigastric aura, the same is true for the flow going through gustatory sensation. The flow going through thought disorder has also an independent course from the other aura sequences on the way from epigastric sensation to aphasia. In view of this flow of aura sequences, three different functional circuits within the limbic system can be suggested: (1) Visual function → memory function, (2) autonomic function → alimentary function → memory function → language function and (3) autonomic function → thought function → memory function → language function. This grouping of aura sensations corresponds, at least to some extent, to the types of seizure constellation proposed by Wieser.³⁶ The first ring to Wieser's type V (posterior neocortical type), the second ring to Wieser's type I and type II (temporobasal type and

Fig 2 Schematic representation of fig 1.

temporal-pole type) and the third ring to Wieser's type III (fronto-cingulate type).

Taylor *et al.*³⁷ classified aura sensations tentatively into three groups, namely "simple primitive", "special senses" and "intellectual". The first group of aura sensations corresponds to anxiety, epigastric sensation and "Es" in our study. Auditory hallucination, visual hallucination and gustatory sensation belong to their "special senses". Thought-disorder, illusion of familiarity and possibly aphasia correspond to "intellectual auras". According to our study, aura-sequences always develop unidirectionally from "simple primitive" to "special senses" and from "special senses" to "intellectual" (fig 2). Taylor speculated that most patients with temporal lobe epilepsy with early onset experienced simple primitive aura and that this could reflect a change in temporal lobe function between childhood and adult life. If that is the case, the aura sequence is supposed to mimic ontogenetical development in view of the rule summarised by us.

In view of the close interrelations between paraphasia and illusion of familiarity and between thought disorder and impairment of verbal comprehension, it could be supposed that at least some parts of ictal paraphasia and ictal impairment of comprehension stem from a disinhibition of memory- and thought-process respectively. This hypothesis is supported also by the fact that some cases with paraphasia, as in the cases of Kisker⁶ and Wilson *et al.*²¹ characteristically developed from a preceding state of readiness, which should be regarded as some kind of illusion of familiarity limited to words. In such cases, the paraphasia and illusion of familiarity interacted with each

other so closely that the manifestation of speech symptoms could be interpreted as an expression of illusion of familiarity. To some extent, this was also true of the interrelation between paroxysmal thought disorder and impairment of verbal comprehension. This is to say, most of our patients not only lost some parts of their ability but also experienced something extra during the ictal aphasia. Therefore, if we call what is lost through illness negative symptoms and what appears extra positive symptoms, ictal aphasia in our patients could be described as a positive symptom.

Contrary to Critchley's opinion, ictal aphasia relating to limbic seizures contrasted well with defect aphasia caused by cerebral infarction and provided information that could not be obtained through studies of stable aphasia.

The first author wishes to thank his deceased wife, Yoshiko, for her help in transcribing raw materials into a definite format and Mrs. Susan Hodge in correcting the English manuscript.

References

- 1 Jackson JH. On slight and severe cerebral paroxysms (epileptic attacks) with auditory warning. *Lancet* 1894;ii:182.
- 2 Penfield W, Jasper H. *Epilepsy and Functional Anatomy of the Human Brain*. Boston: Little, Brown, 1954:366-7.
- 3 Chavany JA, Lobel G, Hagenmuller D. Etude clinique des troubles du langage d'essence épileptique. *Presse Méd* 1956; 64:1449-51.
- 4 Hécaen H, Piercy M. Paroxysmal dysphasia and the problem of cerebral dominance. *J Neurol Neurosurg Psychiatry* 1956; 19:194-201.
- 5 Castells C, Fuster B, Maslenikow V. Crisis disfasicas de origen temporal. *Acta Neurol Latinoam* 1957;3:172-89.
- 6 Kisker KP. Sprachliche Stereotypien bei Temporallappen-Epilepsie (Ein Beitrag zur Konstitution verbaler Sprachteile). *Nervenarzt* 1957;28:366.
- 7 Bingley T. Mental symptoms in temporal lobe epilepsy and temporal lobe gliomas. *Acta Psychiatr Neurol Scand Suppl* 1958;120:96-101.
- 8 Penfield WG, Roberts L. *Speech and Brain Mechanisms*. Princeton, NJ: Princeton University Press, 1959:88-9.
- 9 Hécaen H, Angelergues. Epilepsie et troubles du langage. *Encéphale* 1960;49:138-69.
- 10 Alajouanine T, Sabouraud A. Les perturbations paroxystiques de langage dans l'épilepsie. *Encéphale* 1960;49:95-133.
- 11 Boudouresques J, Roger J, Gastaut H. Crises aphasiques subintrantes chez un épileptique temporal. *Rev Neurol (Paris)* 1962;106:381-93.
- 12 Serafetinides EF, Falconer MA. Speech disturbances in temporal lobe seizures: a study in 100 epileptic patients submitted to anterior temporal lobectomy. *Brain* 1963;86:333-46.
- 13 Acevedo de Mendilaharsu S, Bogacz J, Mendilaharsu C. Étude électroclinique ictale des crises dysphasiques provoquées par activation cardiazolique. *Neuropsychologia* 1964;1:299-312.
- 14 McKeever M, Holmes GL, Russmann BS. Speech abnormalities in seizures: a comparison of absence and partial complex seizures. *Brain Lang* 1983;19:25-32.
- 15 De Pasquet EG, Gaudin ES, Bianchi A. Prolonged and monosymptomatic dysphasic status epilepticus. *Neurology* 1976;26: 244-7.
- 16 Hamilton NG, Matthews T. Aphasia: The sole manifestation of focal status epilepticus. *Neurology* 1979;29:745-8.
- 17 Racy A, Osborn MA, Vern BA, Molinari GF. Epileptic aphasia: first onset of prolonged monosymptomatic status epilepticus in adults. *Arch Neurol* 1980;37:419-22.
- 18 Dinner DS, Lueders H, Ledermann R, Gretter TE. Aphasic status epilepticus: A case report. *Neurology* 1981;31:888-90.
- 19 Gilmore RL, Heilman KM. Speech arrest in partial seizures: Evidence of an associated language disorder. *Neurology* 1981; 31:1016-9.
- 20 Marrosu F, Brundu A, Rachele MG, Marrosu G. Epileptic aphasia as dynamic disturbance: a case report. *Acta Neurol* 1983;5:43-6.
- 21 Wilson A, Petty R, Perry A, Clifford Rose F. Paroxysmal language disturbance in an epileptic treated with clobasam. *Neurology* 1983;33:652-4.
- 22 Rosenbaum DH, Siegel M, Barr WB, Rowan AJ. Epileptic aphasia. *Neurology* 1986;36:822-5.
- 23 Critchley M. Troubles de la parole dans les cas d'épilepsie. *Encéphale* 1960;49:134-7.
- 24 Janz D. *Die Epilepsien*. Thieme, Stuttgart 1969.
- 25 Paillass JE, Vigouroux R, Darcourt G, Naquet R. Considerations sur l'épilepsie occipitale. *Neurochirurgie* 1959;5:3-15.
- 26 Webber WC, Jung CR. Über die epileptische Aura. *Z Gesamte Neurol Psychiatr* 1940;170:211-65.
- 27 Hallen O. Zur Differenzierung der psychomotorischen Anfälle in klinische Formen *Dtsch Z Nervenheilkd* 1962;183:199-217.
- 28 Kawai I, Ohashi H. Ictal speech disturbance and cerebral dominance. *Stud Phonol* 1975;9:40-4.
- 29 Penfield W, Rasmussen T. Vocalisation and arrest of speech. *Arch Neurol Psychiatry* 1949;61:21-7.
- 30 Currie SH, Heathfield KWG, Henson RA, Scott DF. Clinical course and prognosis of temporal lobe epilepsy-survey of 666 patients. *Brain* 1971;94:173-90.
- 31 King DW, Ajmonemarsan CA. Clinical features and ictal patterns in epileptic patients with EEG temporal lobe foci. *Ann Neurol* 1977;2:138-47.
- 32 Meyer-Mickeleit RW. Die Dämmeratacken als charakteristischer Anfallstyp der temporal Epilepsie (psychomotorische Anfälle, Äquivalente, Automatismen). *Nervenarzt* 1953;24:331-46.
- 33 Mullan S, Penfield W. Illusion of comparative interpretation and emotion. *Arch Neurol* 1959;81:269-84.
- 34 Gupta AK, Jeavons PM, Hughes RC, Covanis A. Aura in temporal lobe epilepsy: clinical and electroencephalographic correlation. *J Neurol Neurosurg Psychiatry* 1983;46:1079-83.
- 35 Cole M, Zangwill OL. Déjà vu in temporal lobe epilepsy. *J Neurol Neurosurg Psychiatry* 1963;26:37-8.
- 36 Wieser HG. Electroclinical features of the psychomotor seizure. Stuttgart, Fischer, 1983:193-206.
- 37 Taylor D, Lochery M. Temporal lobe epilepsy: origin and significance of simple and complex auras. *J Neurol Neurosurg Psychiatry* 1987;50:673-81.