Matters arising

University Medical Centre
61105 Ljubljana, Zaloška cesta 7
Yugoslavia

References

- Jesell M, Isch-Treussard, Isch F. Electromyography of striated muscle of anal and urethral sphincters. In: Desmedt JE, ed. *New Developments in Electromyography and Clinical Neurophysiology*. Basel: Karger, 1973; 2:406-20.
- Fowler CJ, Kirby RS, Harrison MJG, Milroy EJJ, Turner-Warwick R. Individual motor unit analysis in the diagnosis of disorders of urethral sphincter innervation. *J Neurol Neurosurg Psychiatry* 1984;47:637-41.
- Stålberg E, Trontelj JV. *Single Fibre Electromyography*. Woking (Surrey): Mirvalle Press, 1979.
- Neill ME, Swash M. Increased motor unit fibre density in the external anal sphincter muscle in ano-rectal incontinence: a single fibre EMG study. *J Neurol Neurosurg Psychiatry* 1980;43:343-7.
- Vodušek DB, Janko M, Lokar J. EMG, single fibre EMG and sacral reflexes in assessment of sacral nervous system lesions. *J Neurol Neurosurg Psychiatry* 1982;45:1064-6.
- Krieger MS, Gordon D, Stanton SL. Single fibre EMG: A sensitive tool for evaluation of the urethral sphincter. Proceedings, International Continence Society 18th Annual Meeting, Oslo, Norway 1st-3rd September 1988. *Urology and Urodynamics* 1988; 7:239-40.
- Stålberg E, Trontelj JV. Clinical neurophysiology: The motor unit in myopathy. In: Rowland EP, ed. *Diseases of Muscle*. Vinken PJ, Bruyn GW, eds. *Handbook of Clinical Neurology*, revised series. Amsterdam: North-Holland Publ Co, in press, 1989.
- Payan J. The blanket principle: a technical note. *Muscle Nerve* 1978;1:432.
- Vodušek DB, Janko M, Lokar J. Direct and reflex responses in perineal muscles on electrical stimulation. *J Neurol Neurosurg Psychiatry* 1983;46:67-71.

Cyclosporin neurotoxicity in cardiac transplant recipient

Sir: We read with interest the report of Lane RJM *et al*¹ explaining the toxicity of cyclosporin on the central nervous system through changes of the blood-brain barrier. However, we feel that their explanation concerning the abnormal MRI seems hazardous. Indeed patient 1 and patient 3 described in the report experienced status epilepticus a few days before the MRI. The occurrence of CT scan hypodensities slightly enhanced by contrast injection following severe epileptic seizures is well described.^{2,4} These hypodensities spontaneously regres-

sive within months and are thought to be due to alterations of the blood brain-barrier produced by local hypoxia, lactic acidosis and loss of vascular autoregulation.

Furthermore, MRI T2-weighted abnormalities were described by Stone *et al*⁵ in one patient and Lesser *et al*⁶ in three patients with intractable focal seizure. These abnormalities were not correlated with the findings on subsequent pathologic examination.

Thus, it seems that the MRI features in Lane's report are probably due to the seizures. The sensitivity of MRI will probably lead to the description of more transient abnormalities secondary to severe epileptic seizures.

In consequence, the discovery of abnormal signal on MRI following severe epileptic seizures cannot be considered as an unequivocal proof of toxic effects of cyclosporin on the blood-brain barrier.

N MAVROUDAKIS
D ZEGERS DE BEYL
Service de Neurologie,
Hôpital Erasme,
Route de Lennik, 808,
1070 Bruxelles, Belgium

References

- Lane RJM, Roche SW, Leung AAW, *et al*. Cyclosporin neurotoxicity in cardiac transplant recipient. *J Neurol Neurosurg Psychiatry*, 1988;51:1434-7.
- Zegers de Beyl D, Hermanus N, Colle H, Goldman S. Focal seizures with reversible hypodensity on the CT-scan. *J Neurol Neurosurg Psychiatry*, 1985;48:187-8.
- Rumack M, Guggenheim MA, Fasules JW, Burdick D. Transient positive postictal computed tomographic scan. *J Pediatr* 1980; 97:263-4.
- Rougiez A, Loiseau P, Rivel J, Cohadon F, Orgogozo JM. Epilepsies partielles de l'adolescence avec anomalies tomodensitométriques, réaction astrocytaire localisée, évolution spontanément régressive. *Rev Neurol (Paris)*, 1984;140:171-8.
- Stone JL, Hughes JR, Barr A, *et al*. Neuroradiological and Electroencephalographic features in a case of temporal lobe status epilepticus. *Neurosurgery* 1986;18:212-6.
- Lesser RP, Modic MT, Weinstein MW, *et al*. 1.5-Tesla magnetic resonance studies in patients with intractable focal seizures. *Neurology* 1985;35(suppl. 1):132 (abstr).
- Sethi PK, Kumar BR, Madan VS, Mohan V. Appearing and disappearing CT scan abnormalities and seizures. *J Neurol Neurosurg Psychiatry* 1985;48:866-9.
- De Groen PC, Aksamit AJ, Rakela J, Forbes GS, Krom RAF. Central nervous system toxicity after liver transplantation. *N Engl J Med* 1987;317:861-6.
- Boon AP, Adams DH, Carey MP, Williams A, McMaster P, Elias E. Cyclosporin-associated cerebral lesions in liver transplantation. *Lancet* 1988;ii:1457.
- Bhatt BD, Meriano FV, Buchwald D. Cyclosporin-associated central nervous system toxicity. *N Engl J Med* 1988;318:788.

Frozen shoulder and other shoulder disturbances in Parkinson's disease

Sir: We read with interest the article by Riley *et al*¹ on the high incidence of shoulder pain in Parkinson's disease. We ourselves have made a similar observation.

As part of a survey to investigate a possible aetiological relationship between Parkinson's disease and essential tremor² we interviewed 100 consecutive Parkinson's disease patients attending the clinic. We were particularly concerned to establish the nature of the first symptom experienced and our interest was aroused when five consecutive patients gave shoulder pain as the initial symptom. On completion of the survey, we found a total of 15 patients (eight female) giving upper limb pain as the first symptom. Pain occurred in the shoulder (12 patients),