SHORT REPORT

Improvement of Parkinsonian features correlate with high plasma levodopa values after broad bean (*Vicia faba*) consumption

J M Rabey, Y Vered, H Shabtai, E Graff, A D Korczyn

Abstract

Five healthy volunteers and six patients with Parkinson's disease (PD) ate 250 g cooked broad beans after 12 hours without treatment. During the next four hours a substantial clinical improvement was noted, three patients showed severe dyskinesias, and plasma levodopa concentrations rose.

(J Neurol Neurosurg Psychiatry 1992;55:725-727)

Levodopa was identified in the seedlings, pods and beans of the broad bean (*Vicia faba*—VF) by Guggenheim in 1913.¹ Although anecdotal cases of PD improvement after ingesting broad beans have been presented,² we are not aware of any studies which have systematically examined this question.

Patients and methods

Six patients with PD participated in the study, after signing written consent (mean age 63.5 years; mean disease duration 13 years; stage III, Hoehn and Yahr scale; mean daily dose of broad beans 541 mg and 54.1 mg carbidopa; 3 also took bromocriptine and 3 took lisuride hydrogen maleate (table 1). Three of the patients also had moderate drug-induced dyskinesia. Five healthy volunteers (mean age 49 years) also participated in the study after giving their written consent. The trial was approved by the hospital Helsinki Committee.

After 12 hours off medication, six PD patients had breakfast at 7 am consisting of tea or coffee with biscuits or a slice of bread. At 8–8.30 am they ate 250 g of cooked VF. Blood samples were obtained before eating VF (time

0) and every 30 minutes thereafter for four hours. Clinical evaluation was scored according to the Webster scale.³ Dyskinesia was evaluated on a separate scale ranging from 0 (no dyskinesia) to 4 (severe dyskinesia). Five days later, the same patients were given orally 125 mg levodopa plus 12.5 mg carbidopa at time 0, after 12 hours off medication. Blood samples were then obtained at time 0 and each 30 minutes thereafter for four hours. The clinical evaluation was performed as described above. Five healthy volunteers also ate 250 g cooked VF at 8–8.30 am. Blood samples were obtained at time 0 and every 30 minutes for four hours.

DOPA MEASUREMENTS

Plasma concentrations of levodopa were determined by an HPLC–ECD method.⁴ Catecholamines were absorbed onto acid washed alumina at pH 8.5 directly from plasma. After being shaken with the alumina, the residue plasma was aspirated. Catecholamines were then eluted from the alumina by a small volume of 0.1 M HC10₄. The HPLC system consisted of an LDC/Milton Roy high pressure pump, a rheodyne injector, and RP chromatographic column (BAS ODS column, 4 × 250 mm, 5 µm) and an electrochemical detector (LC-4, BAS) equipped with a TL-5 glassy carbon electrode. The applied potential at the working electrode was +0.64 V. The mobile phase consisted of monochloroacetate buffer containing 2 mM EDTA and an ion pair agent (SOS, 50 mg/L). The lower limit of detection was 200 pg/ml and the intertest coefficient of variation was 5%.

Statistical analysis for the comparison of clinical performance and levodopa plasma levels in the PD group while eating VF versus levodopa was performed using Wilcoxon and Student's paired *t* tests.

Results

PD patients showed a significant improvement in their motor features after eating VF which was similar to the improvement measured after receiving 125 mg of levodopa plus 12.5 mg of carbidopa (table 2). Moreover, in three patients we observed after VF ingestion severe dyskinesias which were similar to those observed after levodopa administration (table 3). Plasma levodopa levels increased substantially (although somewhat less) after the broad beans compared with levodopa in PD

Departments of Neurology and Clinical Pathology, Tel Aviv Elias Sourasky Medical Center and Sackler Faculty of Medicine, Tel-Aviv University, Israel
J M Rabey
Y Vered
H Shabtai
E Graff
A D Korczyn

Correspondence to:
Dr Rabey, Department of Neurology, Ichilov Hospital, 6 Weizman Street, Tel-Aviv 64239, Israel

Received 19 August 1991
Accepted 23 October 1991.

Table 1 Patients' characteristics

Patient	Sex/age	Disease duration (y)	Stage (Hoehn-Yahr)	Treatment (mean daily dose mg)	Dyskinesia
1	M/60	9	III	500 LD 10 Bro	+++
2	M/65	10	III	750 LD 15 Bro	+++
3	M/70	15	III	500 LD 3 Lis	++
4	F/56	12	III	625 LD 1 Lis	
5	M/66	20	III	500 LD 2 Lis	
6	F/64	12	III	375 LD 20 Bro	

LD—levodopa; Bro—bromocriptine; Lis—lisuride.
Dyskinesia (see method).

Table 2 Cumulative Motor Score (Webster) (n = 6)

Minutes/	0	30	60	90	120	150	180	240
VF	140	114	97	95	96	105	123	130
LD	136	119	99	100	102	115	124	132

VF = *Vicia faba* ingestion.

LD = 125 mg levodopa + 12.5 mg DCI.

Table 3 Cumulative Dyskinesia Score (n = 3)

Minutes/	0	30	60	90	120	150	180	240
VF	0	6	12	8	4	2	0	0
LD	0	5	12	10	6	0	0	0

VF = *Vicia faba* ingestion.

LD = 125 mg levodopa + 12.5 mg DCI.

patients (figure 1). The mean (SD) Cmax was also not significantly different ([Cmax (VF) 0.84 (0.2) $\mu\text{g/ml}$ vs Cmax (levodopa) 1.22 (0.31) μg]). The mean (SD) AUC was similar ([VF 2.95 (0.85) $\mu\text{g/ml/H}^{0-4}$ vs levodopa 3.21

Figure 1 Plasma levodopa levels measured after the ingestion of VF and levodopa in six PD patients.

Figure 2 Plasma levodopa levels measured after the ingestion of VF in six PD patients and five healthy volunteers.

(0.59) $\mu\text{g/ml/H}^{0-4}$). The mean Tmax was also similar.

Five volunteers also showed an increase in plasma levodopa levels after eating broad beans which were similar although considerably lower than the levels measured in PD patients after they had eaten broad beans (fig 2).

Discussion

Three natural potential precursors for dopamine synthesis which can be supplied in the diet, already exist in the brain: phenylalanine, tyrosine and levodopa. Phenylalanine hydroxylase has not been found in the brain⁵ but phenylalanine may be a substrate for tyrosine hydroxylase and can be converted to tyrosine by this enzyme.⁶ High doses of phenylalanine, however, reduce catecholamine synthesis, perhaps due to inhibition of tyrosine hydroxylation.^{7,8} Several investigators found that dl-phenylalanine either had no effect or exacerbated Parkinsonism.^{9,10}

Tyrosine is the normal precursor for catecholamine synthesis. Tyrosine administration does not affect dopamine synthesis in normal animals.⁷ However, if the dopaminergic system is lesioned, the increased firing rate of the remaining neurons will require more tyrosine and in this case, tyrosine administration will enhance dopamine synthesis.^{11,12} Administration of tyrosine to Parkinsonian patients increased the CSF concentration of the dopamine metabolite homovanillic acid (HVA) by 75%.¹³ If, however, tyrosine hydroxylase was severely depleted, as in more advanced Parkinsonism, tyrosine addition would be of little benefit.

The levodopa content of the normal diet is very low or absent.¹⁴ The low concentration of levodopa found in the plasma apparently reflects the activity of peripheral tyrosine hydroxylase in sympathetic neurons rather than dietary intake.¹⁵

In our study, plasma levodopa levels induced in PD patients were higher than the levels measured in healthy volunteers after eating the same amount of broad beans. This difference may be due to the previous treatment of the patients with carbidopa, a potent peripheral decarboxylase inhibitor with a prolonged plasma half-life.¹⁶

- Guggenheim M. Dioxypheylalanin, eine neue Aminosäure aus *Vicia faba*. *Zeitschr Physiol Chem* 1913; 88:276-84.
- Spengos M, Vassilopoulos D. Improvement of Parkinson's disease after *Vicia Faba* consumption. Ninth Int Symposium on Parkinson's disease. Jerusalem, Israel. *Abs* 1988:46.
- Webster DD. Clinical analysis of the disability in Parkinson's disease. *Mod Treat* 1968;5:257-82.
- Shoup RE, Keife J. Plasma catecholamines assayed conveniently and rapidly by LCEC. *Current Separation* 1980;3:1-30.
- Tourian A, Groddard J, Puck TT. Phenylalanine hydroxylase activity in mammalian cells. *J Cell Physiol* 1969; 73:159-70.
- Kaufman S. Properties of the pterin-dependent aromatic amino acid hydroxylase. *Ciba Found Symp* 1974; 22:85-108.
- Gibson CJ, Deikel SM, Young SN, Binik YM. Behavioural and biochemical effects of tryptophan, tyrosine and phenylalanine in mice. *Psychopharmacology* 1982; 76:118-21.
- Nagatsu T, Levitt M, Vdenfriend S. Tyrosine hydroxylase: The initial step in norepinephrine biosynthesis. *J Biol*

- Chem* 1964;239:210–17.
- 9 Cotzias GC, Van Woert MH, Schiffer LM. Aromatic amino acids and modification of parkinsonism. *N Engl J Med* 1967;276:374–9.
 - 10 Granerus AK, Jagenburg R, Roger S, Svanborg A. Phenylalanine absorption and metabolism in parkinsonian patients. *BMJ* 1971;4:262–4.
 - 11 Agid Y, Savoy F, Glavinolli J. Hyperactivity of remaining dopaminergic neurons after partial destruction of the nigrostriatal dopaminergic synthesis in the rat. *Nature (New Biol)* 1973;245:150–1.
 - 12 Melamed E, Hefti F, Wurtman RJ. Tyrosine administration increases striatal dopamine release in rats with partial nigrostriatal lesions. *Proc Natl Acad Sci* 1980;77:4305–9.
 - 13 Growdon JH, Melamed E. Effects of oral l-tyrosine administration on CSF tyrosine and HVA levels in patients with Parkinson's disease. *Neurology* 1980;30:396.
 - 14 Hoeldtke R, Baliga I, Essenberg P, Wurtman RJ. Dihydroxyphenylalanine in rat food containing wheat and oats. *Science* 1972;175:761–2.
 - 15 Eishofer G, Goldstein DS, Ropchak TG, Kopin IJ. Source and physiological significance of plasma 3,4-dihydroxyphenylalanine in the rat. *J Neurochem* 1988;51:1204–13.
 - 16 Vickers S, Stuart EK, Bianchine JR, *et al.* Metabolism of carbidopa, an aromatic amino acid decarboxylase inhibitor in the rat, dog, rhesus monkey and man. *Drug Metab Dispos* 1974;2:9–22.

Neurological stamp

Franciscus de le Boë or Sylvius 1614–72

The chief interest of Franciscus Sylvius was alchemy. He was one of the founders of the iatrochemical school which thought that all phenomena of life and disease were based on chemical action. (He should not be confused with Jacques Dubois, 1478–1555, also called Sylvius, who was an instructor of anatomy in Paris).

Franciscus Sylvius was considered a great teacher of his time and was among the first to introduce ward instruction. His name is linked eponymously to the Sylvian fissure, but his relationship to this structure is obscure. The aqueduct between the third and fourth ventricles has been previously attributed to both Jacobus and Franciscus Sylvius but it had been described long before by others.

Sylvius noted that some tremors were always present but others appeared only with movement. He explored the concept of thermal and tactile senses, had an understanding of ductless glands and was one of the first to propose that the pancreas produced a juice, which together with saliva and a form of fermentation, aided digestion. He was the first to describe tubercles and the process of cavitation. The Netherlands honoured him with this stamp in 1937 (Stanley Gibbons no 470, Scott no B95).

L F HAAS