considerably on the Tensilon test, six had detectable anti-AChR abs, five showed decremental pattern on RNS performed on proximal muscles (deltoid and biceps brachii). In the patient without fluctuations of clinical signs, in whom the Tensilon test was negative, the diagnosis of myasthenia gravis was confirmed by the presence of serum anti-AChR abs. Muscle biopsy performed in this patient as well in another case with no detectable anti-AChR abs, showed normal findings. All patients had a follow up period of at least two years and alternative diagnoses were excluded.

Eight patients had chronic progressive external ophthalmoplegia, 3 males and 5 females, aged 35–71, mean (SD) 56 (14.9) with duration of disease ranging from 1.5–17 years mean (SD) 5.3 (5.1). All patients had clinical weakness restricted to ocular muscles and five complained of mild fluctuations of ptosis. In this group, serum anti-AChR ab titration and repetitive nerve stimulation gave negative results, while the response to the Tensilon test was negative in four cases and equivocal (minimal improvement of ptosis) in the other four. The diagnosis of chronic progressive external ophthalmoplegia was confirmed by the presence of typical ragged red fibres on muscle biopsy in all cases.

Single fibre EMG was performed in the orbicularis oculi muscle during slight voluntary contraction. A single fibre EMG electrode (Medelec SF 25) was inserted in the inferior and lateral portion of the muscle and recordings were made on a Medelec Mistro electromyograph. For each muscle tested, 20 action potential pairs were analysed and the overall mean jitter as mean consecutive difference (MCD), the percentage of potential pairs with prolonged jitter and the percentage of potential pairs with blocking were evaluated. In agreement with other authors,⁷ we considered a jitter study pathological when either a value for mean MCD was above the 95% upper limit for age or a jitter was above the 95% upper limit for potential pairs in more than 10% of pairs. The reference values for these parameters, in different age groups, were derived from literature data.⁷

Single fibre EMG studies were performed, at least 48 hours after discontinuing therapy, on patients with myasthenia.

Results of the single fibre EMG in the orbicularis oculi muscle were pathological in 13 of 14 patients (93%) with purely ocular myasthenia gravis. Eleven patients showed abnormal mean jitter and more than 10% potential pairs had prolonged jitter (in 8 patients blocking was recorded); two patients had mean MCD below the 95% upper limit and respectively 20% and 40% potential pairs with prolonged jitter.

In all patients with chronic progressive external ophthalmoplegia, single fibre EMG studies showed values within the normal limits. The mean jitter did not exceed 25 μ sec. Only one patient had 5% of potential pairs with prolonged jitter and no blocking was recorded.

In patients with ocular myasthenia gravis, the positivity rate with single fibre EMG was higher than with the Tensilon test (86%), RNS (36%) or anti-AChR abs (43%).

Serum anti-AChR abs, which are present in more than 90% of patients with generalised myasthenia gravis, are detectable in a much lower percentage of patients with the

ocular form of the disease. Single fibre EMG has proved the most sensitive technique in detecting the neuromuscular transmission defect in ocular myasthenia gravis, especially when facial muscles are tested. The orbicularis oculi muscle shows abnormal results more often than the frontalis muscle.^{6,8} Unfortunately, single fibre EMG abnormalities can also be seen in primary neuropathic and myopathic disorders as a result of abnormal conduction of the impulse in degenerating or reinnervating nerve terminals and newly formed endplates.^{3,4} In CPEO patients, Krendel and Sanders described increased jitter and blocking in facial (frontalis) and in arm muscles⁶ and a primary defect of neuromuscular transmission has been suggested in this condition.⁵ In our study, single fibre EMG performed in the orbicularis oculi muscle revealed a neuromuscular transmission defect in 93% of patients with purely ocular myasthenia gravis and so demonstrated a diagnostic capacity significantly higher than the Tensilon test, repetitive nerve stimulation or anti-AChR ab titration. Also, in our experience, the sensitivity of single fibre EMG is higher in facial muscles, particularly in the orbicularis oculi, than in the extensor digitorum communis muscle.

M MILONE
ML MONACO
A EVOLI
S SERVIDEI
P TONALI
Institute of Neurology,
Catholic University,
Rome, Italy

Correspondence to: Dr. Milone, Institute of Neurology, Catholic University, Policlinico A. Gemelli, Lgo Gemelli 8, 00168 Rome, Italy

- 1 Evoli A, Tonali P, Bartoccioni E, Lo Monaco M. Ocular myasthenia: diagnostic and therapeutic problems. *Acta Neurol Scand* 1988; 77:31–5.
- 2 Sanders DB, Howard JF, Johns TR. Single-fiber electromyography in myasthenia gravis. *Neurology* 1979;29:68–76.
- 3 Stalberg E, Schwartz MS, Trontelj JV. Single fiber electromyography in various processes affecting the anterior horn cell. *J Neurol Sci* 1975;24:403–15.
- 4 Stalberg E, Trontelj J. Single fiber electromyography. Old Woking, Surrey: Mirvalle Press, 1979.
- 5 Krendel DA, Sanders DB, Massey JM. Single fiber electromyography in chronic external ophthalmoplegia. *Muscle Nerve* 1985;8:624.
- 6 Sanders DB, Howard J. AAEE minimonograph # 25: single fiber electromyography in myasthenia gravis. *Muscle Nerve* 1986;9: 809–19.
- 7 Ad hoc committee of the AAEM special interest group on single fiber EMG, Gilchrist JM, coordinator. *Muscle Nerve* 1992;15: 151–161.
- 8 Trontelj JV, Khuraibet A, Mihelin M. The jitter in stimulated orbicularis oculi muscle: technique and normal values. *J Neurol Neurosurg Psychiatry* 1988;51:814–9.

Soluble interleukin-2 Receptor levels in serum of patients with demyelinating polyneuropathy associated with monoclonal gammopathy

Monoclonal gammopathies, especially monoclonal gammopathy of uncertain significance (MGUS) with IgM M-components, are associated with demyelinating polyneuropathy (PN). In approximately 50% of these patients, the IgM M-component reacts with myelin associated glycoprotein (MAG) or other subcomponents of

myelin and a pathogenetic role of the M-component has been proposed. However, other mechanisms, besides the M-component acting as antibody, may be operating. In vitro studies have shown that monoclonal antibodies against MAG in a patient with monoclonal gammopathy were subject to T-cell regulation,¹ but the role of a T-cell mediated immune response has not been further investigated.

Interleukin-2 (IL-2), which is synthesised by antigen or mitogen activated T lymphocytes, plays an essential role in triggering T-lymphocyte proliferation, T-cell differentiation, and surface expression of IL-2 receptors (IL-2R) on T-cells. It also induces maturation and proliferation of B-lymphocytes. Surface bound IL-2 has a rapid turnover. Therefore the measurement of soluble (s) IL-2R in serum is preferable as a means to evaluate T-cell activation. Elevated serum levels of sIL-2R have been demonstrated in several autoimmune disease states, for example, multiple sclerosis², and the Guillain-Barré syndrome.³

We analysed sIL-2R levels in serum as a measure of T-cell activation in 19 patients (12 men and 7 women, age 43–85 years, mean age 68) with monoclonal gammopathy (15 with MGUS and 4 with Waldenström's macroglobulinaemia; 17 of IgM- and 2 of IgG-isotype) and demyelinating PN. Seventeen of these patients had anti-MAG and/or anti-peripheral nerve myelin (PNM) antibodies. Sera from 19 patients (13 men and 6 women, age 43–80 years, mean age 66) with monoclonal gammopathy (15 with MGUS and 4 with multiple myeloma; 13 of IgG-, 2 of IgM-, 2 of IgA-isotype, and 2 with light chain proteinuria) without PN and without antibodies against MAG or PNM, and sera from 15 healthy individuals (9 men and 6 women, age 47–80 years, mean age 66) without monoclonal gammopathy served as controls. Antibodies against MAG and PNM were analysed by enzyme linked immunosorbent assays (ELISA) essentially as described previously.^{3,4} In 15 patients with monoclonal gammopathy and PN, 2 serum samples or more were obtained during or shortly after tapering off the immunosuppressive treatment. Serum samples were stored at –70°C, and thawed only once. Serum sIL-2R levels were measured using a commercial ELISA kit (Cellfree, T Cell Diagnostics, Inc, Cambridge, MA) according to instructions from the manufacturer. Values exceeding the mean value +2SD (> 919 U/ml) of 50 blood donors were considered pathological (manufacturers information).

Seven of 19 patients with M-component associated PN had increased sIL-2R levels in the first serum sample, compared with 2 of 19 M-component patients without PN ($p = 0.06$ according to Fisher's exact test, one-tail) and 1 of 15 healthy controls ($p < 0.05$ according to Fisher's exact test, one-tail) (fig). At the time of the sample 4 patients with M-component associated PN and 4 M-component patients without PN were receiving immunosuppressive treatment. One of 4 treated patients with M-component associated PN had elevated serum sIL-2R levels and 6 of 15 untreated patients had elevated serum sIL-2R levels.

In 15 patients with M-component associated PN, subsequent serum samples were taken during or shortly after tapering off the immunosuppressive treatment. Seven of

Figure Soluble IL-2R levels in serum at the first sampling occasion of individual patients with M-protein associated with polyneuropathy, M-protein without polyneuropathy and in controls without M-protein. Horizontal bars denote mean values and the horizontal line denotes mean value +2SD of 50 blood donors.

these patients showed decreased (> 10% difference in OD value as compared with the first serum sample), 1 increased and 3 unchanged sIL-2R levels. In the remaining patients no judgement could be made because of continuous treatment in 3 patients and a serum sample taken several years after treatment in one patient. In 3 out of 4 patients with elevated sIL-2R levels at the first sampling occasion, consecutive samples showed remaining high sIL-2R levels.

In a study by Hartung *et al*⁵ comparing serum IL-2 levels in Guillain-Barré syndrome and other neurological diseases, patients with IgM M-component associated demyelinating PN (n = 6) did not have elevated IL-2 levels. However, their patient material was limited, and IL-2 has a rapid turnover making detection of transient elevated IL-2 levels difficult.

In an *in vitro* study¹ it was shown that B-cells from a patient with neuropathy and anti-MAG IgM M-protein were stimulated by pokeweed mitogen activated T-helper cells, and also partially by T-cells in the absence of pokeweed mitogen. T-cells might thus act as regulators of B-cell activation and secretion of immunoglobulins reacting with myelin components. Our findings of activated T-cells in some patients may be due to the action of such regulatory T-cells. We found no statistically significant correlation between serum sIL-2R levels and amount of anti-MAG IgM antibodies. Some of the blood samples, however, were taken at different occasions for analysis of sIL-2R concentration and levels of anti-MAG IgM antibodies.

A substantial number of PN patients, however, had normal sIL-2R levels, arguing for mechanisms other than activated T-cells in the pathogenesis of PN. Also some M-component patients without PN showed

signs of T-cell activation, possibly in response to antigens other than peripheral nerve myelin. The 7 patients with M-component associated PN and elevated sIL-2R levels did not differ from the 12 patients with M-component associated PN with normal sIL-2R levels regarding disease activity or duration of disease.

Factors other than activation of autoreactive (or immunoregulating) T-cells, for example, viral infection, might have been responsible for the elevated levels of sIL-2R in this study. However, no signs or symptoms of infection were noted in our patients. Furthermore, repeated blood samples on different occasions showed persistently elevated sIL-2R levels in three out of four tested patients.

M VRETHEM

Department of Neurology*

T SKOGH

JERNERUDH*

Department of Transfusion Medicine and Clinical Immunology University Hospital, Linköping

B EKSTEDT

Department of Neurology, Örebro Hospital, Örebro

O ANDERSEN

J LYCKE

University Hospital, Göteborg, Sweden

Correspondence to: Dr Vrethem, Department of Neurology, University Hospital, S-581 85 Linköping, Sweden.

We acknowledge the laboratory assistance by Mrs Carina Bergvall. Supported by grants from the University Hospital of Linköping.

- 1 Latov N, Godfrey M, Thomas Y, *et al*. Neuropathy and anti-myelin associated glycoprotein IgM M-proteins: T cell regulation of M-protein secretion *in vitro*. *Ann Neurol* 1985;18:182-8.
- 2 Hartung H-P, Hughes R, Taylor W, Heininger K, Reiners K, Toyka K. T cell activation in Guillain-Barré syndrome and in MS. *Neurology* 1990;40:215-8.
- 3 Cruz M, Jiang Y-P, Ernerudh J, Solders G, Olsson T, Osterman P-O, Link H. Antibodies to myelin-associated glycoprotein are found in cerebrospinal fluid in polyneuropathy associated with monoclonal serum IgM. *Arch Neurol* 1991;48:66-70.
- 4 Vrethem M, Skogh T, Berlin G, Holmgren H, Ernerudh J. Antibodies to peripheral nerve myelin may occur without clinical neuropathy in healthy persons. *J Neuroimmunol* 1991;32:219-22.
- 5 Hartung H-P, Reiners K, Schmidt B, Stoll G, Toyka KV. Serum interleukin-2 concentrations in Guillain-Barré syndrome and chronic idiopathic demyelinating polyradiculoneuropathy: comparison with other neurological diseases of presumed immunopathogenesis. *Ann Neurol* 1991;30:48-53.

HLA class II susceptibility and resistance genes in patients with multiple sclerosis from Northern Spain, by DNA-RFLP genotyping

Table HLA-DR and DQ allele and haplotype frequencies (%) in patients with multiple sclerosis and controls from Asturias (Northern Spain), by DNA-RFLP genotyping

Alleles Haplotypes	All MS patients (n = 96)	Remittent MS (n = 85)	Progressive MS (n = 11)	Controls (n = 123)	p	p*	p†
DR15(DR2)	39	35	39	19	< 0.05	NS	NS
DR16(DR2)	1	1	0	3	NS	NS	NS
DQw5(DQw1)	31	30	0	52	NS	NS	NS
DQw6(DQw1)	49	42	48	38	NS	NS	NS
DR13(DRw6)	19	19	10	41	< 0.05	< 0.05	NS
DR4	28	24	48	26	NS	NS	NS
DQw8(DQw3)	22	19	38	18	NS	NS	NS
DR15/DQw6	36	33	33	18	< 0.05	NS	NS

NS not significant.

p All MS patients vs controls.

p* Remittent MS vs controls.

p† Progressive MS vs controls

The association between susceptibility to multiple sclerosis (MS) and the HLA system has been shown in previous population studies, but the associated HLA antigens have varied in different ethnic groups.¹ Southern Europe is an area of interest for the HLA-MS association, since the associations observed in most Northern European populations are usually absent. Previous studies have reached conflicting conclusions on whether particular HLA types influence the clinical course of the disease.²

HLA typing has traditionally been accomplished serologically or immunologically, but some subtypes have so far evaded serological recognition. An alternative method is typing at the DNA level with restriction fragment length polymorphism (RFLP). This HLA genotyping detects further subtypes of serologically defined -DR and -DQ specificities, which will permit a better understanding of the HLA-MS association.³

We examined MS associated HLA-DR and DQ alleles, characterised by RFLP, at the genomic level in 96 MS patients (63 women and 33 men) from Asturias, a region in Northern Spain with a medium MS prevalence of 24/100000. MS was defined clinically or by laboratory support using the Poser *et al* criteria.⁴ Eleven had primarily chronic progressive MS and 85 had relapsing-remitting MS. The latter group also included patients with a secondary progressive evolution of symptoms. A total of 123 healthy unrelated Spanish individuals were used as controls.

To carry out the HLA typing by DNA-RFLP analysis, the DNA from peripheral blood leucocytes was digested with the restriction endonucleases Taq I and hybridised with probes to DR beta, DQ beta and DQ alpha genes, using standard methodological and analytical procedures. Haplotypes DR-DQ were assigned according to the pattern of bands following the Bidwell method.³ The results of HLA-class II frequencies in MS patients and controls were compared by using the Chi-square test with Yates's correction and p values were multiplied by the number of allotypes tested.

DR15 (p < 0.05) and DR15/DQw6 (p < 0.05) were significantly increased in the whole MS group compared with the controls (table). DRw13 was significantly decreased (p < 0.05). DQw5 and DRw13/DQw5 were also decreased but the differences were not significant when corrected for the number of allotypes tested.

The frequency of MS patients that were positive for DR15 and/or DR16 (splits of DR2) appears increased compared with the