- 31 Durelli L, Cocito D, Riccio A, *et al*. High dose intravenous methylprednisolone in the treatment of multiple sclerosis: clinical-immunologic correlations. *Neurology* 1986;36:238-43.
- 32 Chong G, Mourad G, Levy P, Nedelec G, Guidicelli C, Mion C. Chorio-retinite à cytomegalovirus: complication rare et grave chez le transplanté rénal. Intérêt du diagnostic précoce et du traitement prolongé par le DHPG. *Néphrologie* 1990;11:255-8.
- 33 Tokunaga N, Ichikawa Y, Kuboshiro M, Higashi T, Fujino K, Kaji M. A case of pulmonary tuberculosis associated with adult respiratory distress syndrome during corticosteroid treatment of rheumatoid arthritis. *Kekkaku* 1990;65:539-43.
- 34 Walsh DA, Durance RA. Fatal acute pyelonephritis following pulsed methylprednisolone for rheumatoid arthritis. *Ann Rheum Dis* 1990;49:955-6.
- 35 Chikanza C, Fernandes L. Arrhythmia after pulse methylprednisolone therapy. *Br J Rheumatol* 1991;30:392-3.
- 36 Tvede N, Nielsen LP, Andersen V. Bradycardia after high-dose intravenous methylprednisolone therapy. *Scand J Rheumatol* 1986;15:302-4.
- 37 Jones JG. Avascular necrosis and pulsed methylprednisolone in RA. *Br J Rheumatol* 1988;27:497-8.
- 38 Gardiner PV, Griffiths ID. Sudden death after treatment with pulsed methylprednisolone. *BMJ* 1990;300:125.
- 39 Lyons PR, Newman PK, Saunders M. Methylprednisolone therapy in multiple sclerosis: a profile of adverse effects. *J Neurol Neurosurg Psychiatry* 1988;51:285-7.
- 40 Williams IA, Mitchell AD, Rothman W, Tallett P, Williams K, Pitt P. Survey of the long term incidence of osteonecrosis of the hip and adverse medical events in rheumatoid arthritis after high dose intravenous methylprednisolone. *Ann Rheum Dis* 1988;47:930-3.
- 41 Snead III OC, Benton JW, Myers GJ. ACTH and prednisone in childhood seizure disorders. *Neurology* 1983;33:966-70.
- 42 Dorfman A, Apter NS, Smull K, Bergenstal DM, Richter RB. Status epilepticus coincident with use of pituitary adrenocorticotrophin hormone. *JAMA* 1951;146:25-7.
- 43 Hasaerts D, Dulac O. Traitement par hydrocortisone des épilepsies généralisées secondaires du grand enfant. *Arch Fr Pediatr* 1989;46:635-9.
- 44 Stephens EHM, Noad KV. Status epilepticus occurring during cortisone therapy. *Med J Aust* 1951;38:334-5.
- 45 Conford E, Oldendorf W. Epilepsy and blood brain barrier. In: Delgado-Escueta A, Ward A, Woodbury D, Porter R. eds. *Advances in neurology: basic mechanisms of epilepsies*. New York: Raven Press, 1986:787-812.
- 46 Sokrab T-E O, Kalimo H, Johansson BB. Parenchymal changes related to plasma protein extravasation in experimental seizures. *Epilepsia* 1990;31:1-8.
- 47 Siemes H, Siebert M, Aksu F, Emrich R, Hanefeld F, Scheffner D. CSF protein profile in infantile spasms. Influence of etiology and ACTH or dexamethasone treatment. *Epilepsia* 1984;3:368-76.
- 48 Miller DH, Thompson AJ, Morrissey SP, *et al*. High dose steroids in acute relapses of multiple sclerosis: MRI evidence for possible mechanisms of therapeutic effect. *J Neurol Neurosurg Psychiatry* 1992;55:450-3.

NEUROLOGICAL STAMP

Pierre Joseph Pelletier (1788–1842) and Jean Bienaime Caventou (1795–1887)

The French chemist Pelletier's major work was the investigation of drugs, which he began in 1809. Later he worked with the pharmacist and organic chemist Caventou. Their collaborative studies, which began in 1817, included the isolation of strychnine from nuxvomica (1818), caffeine (1821), and quinine from cinchona bark (1820). It was not until 1936 that Wolff reported on the effectiveness of quinine in relieving myotonia. Later, in 1955, Geschwind and Simpson noted the "quinine-like" action of procainamide on repetitive firing of cardiac muscle and suggested that it might be effective in combating myotonia.

Caventou, an expert in toxicology (Professor of Toxicology at the Ecole de Pharmacie (1835–1860)), also reported on cases of arsenical poisoning. Philatelically both were honoured by France in 1970 to commemorate the 150th anniversary of the discovery of quinine (Stanley Gibbons 1870, Scott 1268). Pelletier and Caventou are regarded as the founders of alkaloid chemistry. Caventou's early successes were not repeated in later life. After Pelletier's death in 1842 Caventou published nothing further.

L F HAAS