- 18 Williams DB, Lyons MK, Yanagihara T, Colgan JP, Banks PM. Cerebral angiotropic large cell lymphoma (neoplastic angioendotheliosis): therapeutic considerations. *J Neurol Sci* 1991;103:16–21.
- 19 Lim HW, Anderson HM. Angioendotheliomatosis associated with histiocytic lymphoma: response to systemic chemotherapy. *J Am Acad Dermatol* 1985;13:903–8.
- 20 Axelsen RA, Laird PP, Horn M. Intravascular large cell lymphoma: diagnosis on renal biopsy. *Pathology* 1991;23:241–3.
- 21 Petroff N, Koger OW, Fleming MG, et al. Malignant angioendotheliomatosis: an angiotropic lymphoma. *J Am Acad Dermatol* 1989;21:727–33.
- 22 Nishikawa K, Sekiyama S, Suzuki T, et al. A case of angiotropic large cell lymphoma manifesting nephrotic syndrome and treated successfully with combination chemotherapy. *Nephron* 1991;58:479–82.
- 23 Yousem SA, Colby TV. Intravascular lymphomatosis presenting in the lung. *Cancer* 1990;65:349–53.
- 24 Stroup RM, Sheibani K, Moncada A, Purdy LJ, Battifora H. Angiotropic (intravascular) large cell lymphoma. A clinicopathologic study of seven cases with unique clinical presentations. *Cancer* 1990;66:1781–8.
- 25 Kao NL, Broy S, Tillawi I. Malignant angioendotheliomatosis mimicking systemic necrotizing vasculitis. *J Rheumatol* 1992;19:1133–5.

NEUROLOGY IN LITERATURE

Visual disorders

With the exception of the physiological observations of Norman Douglas and Edgar Allan Poe (of which Poe more impresses), the extracts I have chosen principally refer to vascular disorders affecting vision. Charlotte Brontë's father presumably suffered an episode of occipital ischaemia, probably related to transient occlusion of the terminal basilar artery. A similar process probably affected Nero. The accounts by Wilkie Collins and Thomas Hardy are intriguing. The subject of Collins' description is in his 20s. Prodromal pain does not exclude a diagnosis of Leber's optic atrophy although it would usually be distributed frontally or in the orbital region. Hardy seems to suggest that Yeobright's ocular pain is photophobic and it is perhaps more likely that he is describing an acute iritis than optic neuritis. If the latter had been the case, Hardy would have antedated the descriptions of Nettleship, Parinaud, Uhthoff, Buzzard, and Marcus Gunn!

Edgar Allan Poe, 1841, Tales of mystery and imagination. The murders in the Rue Morgue

To look at a star by glances—to view it in a side-long way by turning towards it the exterior portions of the retina (more susceptible of feeble impression of light than the interior) is to behold the star distinctly—is to have the best appreciation of its lustre—a lustre which grows dim just in proportion as we turn our vision fully upon it. A greater number of rays actually fall upon the eye in the latter case, but, in the former, there is the more refined capacity for comprehension.

Elizabeth Gaskell, 1857, The life of Charlotte Brontë
My dear father has borne the severe winter very well, a circumstance for which I feel the more thankful as he had many weeks of very precarious health last summer, following an attack from which he suffered in June, and which for a few hours deprived him totally of sight, though neither his mind, speech nor even his powers of motion were in the least affected. I can hardly tell you how thankful I was, when, after that dreary and almost despairing interval of utter darkness, some gleam of daylight became visible to him once more. I had feared that paralysis had seized the optic nerve. A sort of mist remained for a long time; and, indeed, his vision is not yet perfectly clear, but he can read, write, and walk about. . .

Wilkie Collins, 1857, The dead secret
About the last work he did, poor fellow, was the repairing of my watch—here it is; goes as regular as a

steam-engine. I hadn't got it back into my fob very long before I heard that he was getting a bad pain at the back of his head, and that he saw all sorts of moving spots before his eyes. String him up with lots of port wine, and give him three hours a day on the back of a quiet pony—that was my advice. Instead of trying it, they sent for doctors from London, and blistered him behind the ears and between the shoulders, and drenched the lad with mercury, and moped him up in a dark room. No use. The sight got worse and worse, flickered and flickered, and went out at last like the flame of a candle. . . . His father was half out of his mind. Took him to oculists in London, and oculists in Paris. All they did was to call the blindness by a long Latin name, and to say that it was hopeless and useless to try an operation. Some of them said it was the result of the long weakness from which he had twice suffered after illness. Some said it was an apoplectic effusion in the brain. . . . So they brought him back home, blind, blind he is now; and blind he will remain, poor fellow, for the rest of his life.

Thomas Hardy, 1878, The return of the native
One morning, after a severer strain than usual, he awoke with a strange sensation in his eyes. The sun was shining directly upon the window-blind, and at his first glance thither-ward a sharp pain obliged him to close his eyelids quickly. At every new attempt to look about him the same morbid sensibility to light was manifested, and excoriating tears ran down his cheeks. He was obliged to tie a bandage over his brow while dressing; and during the day it could not be abandoned

. . . Towards evening he arrived, and pronounced the disease to be an acute inflammation induced by Clym's night studies, continued in spite of a cold previously caught, which had weakened his eyes for the time.

. . . He was not to be blind: that was enough. To be doomed to behold the world through smoked glass for an indefinite period was bad enough, and fatal to any kind of advance.

Arnold Bennett, 1906, Whom God hath joined
"When I woke up the next morning in the hotel, I couldn't make out what was the matter with me. Something with my left eye—I couldn't see out of it, and I was so feeble, I just managed to get as far as a doctor's. The doctor said I'd had a haemorrhage into my retina while I was asleep."

Continued on page 370

stretched over the transverse process of the atlas during forced head extension.^{6,11}

The described method of transcranial Doppler monitoring of blood flow velocities of posterior cerebral arteries may have advantages compared with angiography, as various combinations of head movements can be performed sequentially and analysed for their influence. The high time resolution of transcranial Doppler is particularly useful to show rapid and brief haemodynamic changes. This approach measures the net haemodynamic effect of possible obstructions at different levels and is therefore likely to correlate with clinical symptoms. The absence of a so called temporal bone "ultrasound window" may preclude investigation of posterior cerebral arteries in a considerable percentage of elderly people. This is not usually a problem when examining adolescents, the population at risk for stretch syncope. The technique reported here of monitoring both posterior cerebral arteries by a temporal approach using a two channel Doppler system as compared with basilar artery and vertebral artery monitoring by a suboccipital transforamenal approach with the freehand technique has the advantage of stable Doppler signals despite head movements.¹² Suboccipital transducer position is hampered by large probe motion artefacts during head movements as well as the risk of signal loss due to changing depth of the vessel of interest and difficulties maintaining vessel insonation.¹³

The patients described here show that it may be difficult to prove vertebral artery compression during angiography because of the inability to reproduce the exact offending

posture. Also, angiography is invasive and not without risk. Transcranial Doppler monitoring on the other hand, although able to prove haemodynamically significant vertebral artery compression, does not identify its level and the underlying cause.¹⁴ Avoidance of the triggering posture in most patients is all that is necessary to eliminate further attacks.

- 1 Manolis AS, Linzer M, Salem D, Estes MNA. Syncope: current diagnostic evaluation and management. *Ann Intern Med* 1990;112:850-63.
- 2 Pelekanos JT, Dooley JM, Camfield PR, Finley J. Stretch syncope in adolescence. *Neurology* 1990;40:705-7.
- 3 Buedingen HJ, Staudacher T. Evaluation of vertebrobasilar disease. In: Newell DW, Aaslid R, eds. *Transcranial Doppler*. New York: Raven Press, 1992:167-95.
- 4 Grad A, Baloh R. Vertigo of vascular origin. *Arch Neurol* 1989;46:281-4.
- 5 Gilles FH, Bina M, Sotrel A. Infantile atlanto-occipital instability: the potential danger of extreme extension. *Am J Dis Child* 1979;133:30-7.
- 6 Toole JF, Tucker SH. Influence of head position upon cerebral circulation. Study of blood flow on cadavers. *Arch Neurol* 1960;2:616-23.
- 7 Faris AA, Poser CM, Wilmore DW, Agnew CH. Radiologic visualization of neck vessels in healthy men. *Neurology* 1963;13:386-96.
- 8 Schmidt H, Sartor K, Heckel RW. Bone malformations of the craniocervical region. In: Vinken PJ, Bruyn GW, eds. *Handbook of clinical neurology*. Vol 32. Amsterdam: North-Holland Publishing Co, 1978:1-98.
- 9 Coria F, Rebollo M, Quintana F, Polo JM, Berciano J. Occipitoatlantal instability and vertebrobasilar ischemia. Case report. *Neurology* 1982;32:303-5.
- 10 Wiesel SW, Rothman RH. Occipitoatlantal hypermobility. *Spine* 1979;4:187-91.
- 11 Saver JL, Easton JD, Hart RG. Dissections and trauma of cervicocerebral arteries. In: Barnett HJM, ed. *Stroke. Pathophysiology, diagnosis and management* 2nd ed. New York: Churchill Livingstone, 1992:671-88.
- 12 Brautaset NJ. Provokable bilateral vertebral artery compression diagnosed with transcranial Doppler. *Stroke* 1992;23:288-91.
- 13 Fujioka KA, Ernsberger AM, Nicholls SC, Spencer MP. Transcranial Doppler assessment of mechanical compression of the vertebral arteries. *Journal of Vascular Technology* 1991;15:254-9.
- 14 George B, Laurian C. Impairment of vertebral artery flow caused by extrinsic lesions. *Neurosurgery* 1989;24:206-14.

NEUROLOGY IN LITERATURE

Visual disorders

Continued from page 366

Norman Douglas, 1917, South wind

"Or perhaps you strained one of your eyes bathing. It has happened to me occasionally. Darkness is the best remedy. It rests the optic nerve."

Marcel Proust, 1920, 1921, 1922, Remembrance of things past. Vol 2

"Look in this cabinet. I have all the hats worn by Madame Elisabeth, by the Princesse De Lamballe, and by the Queen. They don't interest you, it's as though you couldn't see. Perhaps you are suffering from an affection of the optic nerve."

Marcel Proust, 1923, 1925, 1927, Remembrance of things past. Vol 3

"I was very distressed when I was told that he had lost his sight." "Yes, it's true that his eyes were affected by his stroke. For a time he could see nothing at all, just imagine, during the cure, which as a matter of fact did him a great deal of good, he was for several months unable to see more than a man born blind."

Gaius Suetonius (AD69-140), The twelve Caesars - translated by Robert Graves, 1957

In reference to Nero

A warning portent made him cancel the Alexandrian voyage, on the very day when his ship should have sailed: during his farewell round of the temples he had sat down in the shrine of Vesta, but when he rose to leave, the hem of his gown caught fire and then a temporary blindness overcame him.

G D PERKIN
Regional Neurosciences Centre,
Charing Cross Hospital,
London W6 8RF, UK