NEUROLOGICAL INVESTIGATIONS

Multimodal monitoring in neurointensive care

P J Kirkpatrick, M Czosnyka, J D Pickard

The pathophysiological mechanisms occurring in cerebral ischaemia are multiple, complex, and incompletely understood.¹ However, despite the diverse aetiology for brain injury, different processes operate to cause common manifestations such as a raised intracranial pressure (ICP), derangements in cerebral blood flow (CBF), and brain hypoxia.^{1,2} Such changes in the pathophysiological state of the cerebral tissues may be transient and last only a few minutes. Although intermittent monitoring with serial cranial imaging methods (such as enhanced computed and emission tomography) in specialised institutions³⁻⁵ provide good spatial information, they are likely to miss transient events. Also, the necessary intensive support for these precarious patients is difficult to maintain within such imaging facilities. Thus methods for assessing brain function in an uninterrupted fashion have attracted increased clinical attention,⁶⁻¹¹ particularly those that can be adapted for bedside monitoring, which reduces the need for patients' transfer.

A single monitored cerebral event, such as a period of raised ICP, may be a manifestation of various different pathophysiological changes. Cerebral swelling from ischaemia (oligaemia), and increased cerebral blood volume from hyperaemia are examples in which the contrast in pathology is extreme. Blind ICP treatment in both instances using agents such as mannitol may be beneficial in the first case, but potentially aggravate the raised ICP in the second.^{12,13} Thus directing treatment according to one measured variable may be inappropriate. Similarly, whereas controlled hyperventilation has traditionally been used to treat raised ICP by encouraging reactive vasoconstriction, recent evidence suggests that in situations of cerebral oligoemia these manoeuvres can increase cerebral ischaemia and lactic acid production.¹⁴ By monitoring several different variables, each providing relevant information on different aspects of brain physiology, a greater understanding of the individual situation can be gathered. The aim would be a more accurate targeting and policing of treatment. Computer support of multimodality monitoring¹⁵ helps the observer to identify important cerebral events among the background noise and artefacts (often induced within the hostile environment of an intensive

care unit), and helps in the interpretation of complex information.

The purpose of this chapter is to provide an introduction to the novel methods that are available for the continuous assessment of cerebral perfusion, haemodynamics, and oxygenation. Continuous monitoring techniques for different variables concerning the health of the brain are now available, and these include measurements of ICP, cerebral perfusion pressure (CPP), jugular venous oxygen saturation (SjO₂), and cortical electrical activity.^{6-11,16-18} General systems monitors, such as pulse oximetry, end tidal CO₂, and temperature are clearly of importance but will not be discussed here. More recently additional methods have been introduced such as transcranial Doppler,^{6,8} laser Doppler flowmetry,⁹⁻¹⁰ and near infrared spectroscopy.¹¹

Computing support of data analysis

In an established neurointensive care facility enormous quantities of data can be captured from each patient from which information on cerebral autoregulation, oxygenation, metabolite production, and function can be obtained. Recognition of changing cerebrovascular haemodynamics and oxygenation demands not only reliable monitoring techniques, but also complex and time consuming signal analysis. This can only be provided by dedicated computer support.

The first specialised computer based systems for neurointensive care were introduced at the beginning of the 1970s. Initially these systems were oriented to the monitoring of ICP and arterial blood pressure (ABP) allowing calculation of CPP and a basic analysis of the pulsatile ICP waveform.^{19,20} By contrast, contemporary systems are highly complex multichannel digital trend recorders with built in options for complex signal processing. The considerable flexibility of such systems^{15,20,21} permits almost unlimited signal analysis which in itself can generate a state of data chaos. Thus the modern user is faced with the problem of which variables should be considered, and how the data should be interpreted. This information should then be presented in a manner which is comprehensible to medical and nursing staff. The mechanism of presentation is also important.

MRC Cambridge
Centre for Brain
Repair and Academic
Neurosurgical Unit,
Addenbrooke's
Hospital, PO Box 167,
Cambridge CB2 2QQ,
UK
P J Kirkpatrick
M Czosnyka
J D Pickard
Correspondence to:
Mr Kirkpatrick

Although personal computers with designated software are portable, they have yet to gain widespread clinical acceptance as an intensive care tool. They are seen as stand alone instruments requiring specialised skills for their operation, and occupying precious space. By contrast, commercial hardware systems with a customised console are more user friendly, but are far more expensive and less flexible.

The intensive multimodality monitoring system adopted in the Cambridge Neurosurgical Intensive Care Unit is based on software for the standard IBM compatible personal computer, equipped with a digital to analogue converter and RS232 serial interface. It was introduced into clinical practice in Poland, Denmark, and the United Kingdom in the late 1980s and has recently been extended into a system for multimodal neurointensive care monitoring and waveform analysis.¹⁵ Most data have been derived from patients with head injuries, common occupants of the neurointensive care unit. However, the same techniques are being increasingly applied to those with severe stroke, subarachnoid haemorrhage, cerebral infections, and encephalopathies.²²

Intracranial pressure and cerebral perfusion pressure

Pathophysiological mechanisms responsible for stabilisation of ICP within a rigid skull vault are complex.⁶ Changes in the CSF and cerebral blood volume may compensate for longstanding volumetric changes of cerebral tissue. However, in acute injury small changes in cerebral parenchymal volume can cause gross changes in ICP and CPP. Accompanying cerebral haemodynamic deterioration may occur with the possible sequel of secondary ischaemic brain damage. Thus the continuous assessment of ICP is a key component of any cerebral multimodality monitoring system.

Reliable measurement of ICP still depends on invasive systems. Non-invasive methods, such as transfontanometry,²³ tympanic membrane displacement,²⁴ and assessment of transcranial Doppler flow velocity pulse waveform,²⁵ are difficult to calibrate and have not achieved suitable accuracy. The least invasive systems available use epidural probes, but there is still uncertainty regarding the precise relation between ICP and pressure in the extradural space. Intraventricular or subdural fluid filled catheters with external pressure sensitive elements can measure ICP directly but display signal drift, have limited frequency response,²⁶ and present a risk of infection. By contrast, ICP microtransducers measure CSF or intraparenchymal pressures with high accuracy, minimum signal drift, and a good frequency response,^{27–30} and as their support bolt provides an airtight seal with the skull bone they can be safely used for long term monitoring without concern for infection.

Intracranial pressure measurements are used to estimate CPP; providing that ICP estimates intracerebral venous pressure:

$$\text{mean CPP} = \text{mean ABP} - \text{mean ICP}$$

Sufficient CPP is required to maintain a stable CBF with an autoregulatory reserve. A CPP below 60–70 mm Hg may result in a compromise in various haemodynamic modalities.^{3 8 31 32} There is also an increased chance of a poor outcome if CPP falls below these thresholds in patients with head injury.^{33–35} However, policies to therapeutically maintain a high CPP are controversial. Non-reactive vessels may result in hyperaemia, increasing vasogenic oedema, and secondary increase in ICP. It is also probable that there are considerable patient dependent differences in the optimal level of CPP. Thus although many authors evaluate such thresholds in their group analysis^{31 32 35 36} and demonstrate critical values ranging from 55 to 80 mm Hg, a general threshold between adequate and non-adequate CPP for each patient is difficult to define. The threshold of CPP causing haemodynamic deficit should be considered as a time dependent factor,^{37 38} hence the real time assessments of the relation between haemodynamic modalities and CPP is essential.

One approach to identifying a “safe pressure” zone in the individual patient is to use information derived from the ICP and CPP waveforms. For example, the analysis of the relation between the mean ICP and the amplitude of its waveform,^{39 40} the analysis of the shape of pulse wave³⁹ and its relation to respiratory oscillations,⁴¹ the transmission of the ABP wave into the intracranial compartment,^{42 43} and the spectral analysis of the fundamental^{44 45} and subsequent ICP waveform harmonics⁴⁶ have all been considered. The basic phenomenon of an increase in ICP amplitude with rising mean ICP was seen in the early recordings made by Ryder *et al* in 1953.⁴⁷ Using a monoexponential model of cerebrospinal pressure-volume relation Langfitt *et al*⁴⁸ later postulated that if an increase in cerebral blood volume during one heart contraction was constant, it would produce a higher pressure response when the ICP level was raised. However, when the ICP becomes very high with a compensatory maximal vasodilatation a secondary decrease in the ICP pulse amplitude is seen (fig 1). The linear correlation coefficient between mean ICP and ICP pulse amplitude values can be calculated and has been termed RAP (R = symbol of correlation, A = amplitude, P = pressure).^{44 49} This index describes time dependent changes in the relation between mean ICP and the pulse amplitude. The advantage is that the coefficient has a normalised value from –1 to +1 (fig 1) allowing comparison between patients. The relation of RAP and ICP or CPP in pooled analysis of patients with head injury shows that a positive index close to +1 is expected in patients with head injury with moderately raised ICP (>15 mm Hg) and CPP above 50 mm Hg. Decrease in RAP to 0 is found with very high ICP and very low CPP (fig 2), which are predictive of a poor outcome (fig 3).⁵⁰ The RAP index as a time related factor often anticipates brainstem herniation due to

Figure 1 Pulse amplitude (AMP) of ICP waveform increases with mean ICP until a critical threshold is reached above which a decrease occurs (upper graph). The correlation coefficient between AMP and mean ICP (RAP—bottom graph) marks this threshold by switching from positive to negative values. Redrawn from Pickard and Czosnyka.⁶

Figure 2 Relation between the mean value of RAP index and the day average ICP (upper graph) and CPP (bottom graph) in a group of 56 patients after head injury (bars denote 95% confidence intervals for mean value). Redrawn from Czosnyka et al.⁵⁰

Figure 3 A terminal increase of mean ICP and decrease in CPP leading to brain stem herniation at 12:30 (x axis: time in hours). RAP coefficient decreased from positive to negative values 1.5 hours earlier showing generalised deterioration of cerebrovascular reactivity reserve. AMP = pulse amplitude of ICP waveform. Redrawn from Czosnyka et al.¹⁵

an excessive rise in ICP (fig 4). Because the RAP coefficient is calculated using the fundamental harmonic of the ICP pulse wave the practical advantage of this index is that there is no need to use a pressure transducer with a wide bandwidth; a simple subdural catheter connected to an external membrane transducer will suffice.

Transcranial Doppler ultrasonography

Transcranial Doppler ultrasonography⁵¹ allows non-invasive measurement of blood flow velocity in basal cerebral arteries. Most data have been derived from the middle cerebral artery as this vessel is technically the simplest to insonate, and is the most relevant as 80% of supratentorial cerebral blood flow passes through it. Although blood flow velocity cannot express volume flow,⁵² the dynamic changes of CBF are almost always reflected in transcranial Doppler readings. Experience with transcranial Doppler in neurointensive care monitoring is still limited because of problems of long term fixation of the ultrasound probe and of interfacing with a computerised system.¹⁵ However, the high dynamic resolution provided and confirmed correlation with other haemodynamic modalities is encouraging increasing numbers of neurointensivists to adopt the technique.^{31 32 38 53-55} In addition to calculating the time averaged meanflow velocity, recent transcranial Doppler machines provide information on the flow waveform which is affected by pulsations of CPP and by the resistance and compliance (mechanoelectric properties) of the cerebrovascular bed.⁵⁶ Thus transcranial Doppler provides great potential in cerebrovascular investigations^{32 37 38} for assessment of cerebrovascular autoregulatory reserve, reactivity, CPP, cerebral hyperaemia, post-traumatic spasm, and in the estimation of cerebral tamponade.⁵⁷

CEREBRAL AUTOREGULATION

With continuous transcranial Doppler, autoregulation can be assessed by observing the responses to spontaneous changes in ABP, transient changes in CPP induced by a three second carotid compression, or longer periods (20–40 s) of reduced CPP induced by inflating and releasing large blood pressure cuffs applied to the legs.⁵⁸⁻⁶¹ Continuous assessment of transcranial Doppler pulsatility indices which vary according to the state of autoregulation^{31 32 37 38} can be useful for the on line monitoring of critical thresholds in CPP (see later). More recently an evaluation of the gradient of linear regression between the different components of the flow velocity waveform (systolic, mean, and diastolic) and the CPP (fig 5)³⁷ show that these gradients are dependent on mean CPP and correlate with outcome after head injury (fig 6).

CEREBROVASCULAR REACTIVITY

The response of flow velocity to changes in CO₂ concentration characterises vascular reactivity. Decreased reactivity is reported with decreasing CPP⁶² and in patients with poor outcome after head injury.⁶³

Figure 4 Mean values of CPP, ICP, and RAP over a 24 hour period in 42 patients with head injury in different outcome groups. ICP is significantly lower for moderate/good (m/g) patients. RAP is significantly lower for dead or persistent vegetative (PVS/d) patients. RAP = correlation coefficient between pulse amplitude (first harmonic) and mean intracranial pressure; sd = severely disabled patients. Redrawn from Czosnyka et al.³⁷

SPASM AND HYPERAEMIA

Flow velocity in excess of 100 cm/s occurs in 10% to 20% of patients after head injury,^{64 65} but the differentiation between hyperaemia and vasospasm can be difficult and often requires measurement of a further variable such as jugular venous oximetry.⁴⁸ If the ratio of middle cerebral artery flow velocity to the internal carotid artery flow velocity exceeds 3 then vasospasm is likely.⁶⁶ Detailed flow velocity waveform analysis and detection of a dicrotic notch in the pulse pattern⁶⁷ are not very helpful.

NON-INVASIVE ASSESSMENT OF CEREBRAL PERFUSION PRESSURE

The pulsatility index (flow velocity amplitude / flow velocity mean) is a dimensionless index that is independent of sampling variation provided the signal to noise ratio is good and the gain setting of the instrument is constant. Most modern software packages provided with transcranial Doppler automatically calculate the pulsatility index which is averaged from several cardiac cycles. The potential importance of the pulsatility index in brain injury is that increases in the index provide a non-invasive artefact free measure of failing autoregulation.^{8 21 25} Clinical experience has shown a closer correlation between CPP and pulsatility index than between mean flow velocity and CPP which has facilitated the use of this variable for the non-invasive estimation of CPP in patients with head injury (fig 7).

DETECTION OF CEREBRAL TAMPONADE

Transcranial Doppler has been used to assist in the diagnosis of brain death. At very low levels of CPP the critical closing pressure for cerebral arterioles is reached resulting in the collapse of the microcirculation and vascular infarction. Net forward blood flow diminishes

Figure 5 Three specific patterns of relation between systolic (FVs), diastolic (FVd) and time averaged (FV) blood flow velocity and CPP: Left: both systolic and diastolic are pressure-passive (autoregulation exhausted). Middle: diastolic and time average FV pressure-passive but systolic FV not pressure passive (autoregulatory reserve compromised). Right: systolic and diastolic FV are not pressure passive (good autoregulatory reserve). Figure redrawn from poster presented at the 3rd International Neurotrauma Symposium, Toronto, 1995 by Czosnyka et al.

Figure 6 Analysis of the gradient of average, systolic, and diastolic flow velocity time versus CPP in different outcome groups (62 patients with head injury; transcranial Doppler examinations done daily). Black bars show 95 % confidence intervals for mean. m/g = moderate/good outcome; sd = severe disability; PVS/d = persistent vegetative or dead patients. syst FV = systolic flow velocity. Redrawn from Czosnyka et al.³⁷

and the transcranial Doppler pattern shows reversal of flow during diastole.⁵⁷

Laser Doppler flowmetry

Laser Doppler flowmetry is a technique which provides a continuous measure of relative microcirculatory flow.⁶⁸⁻⁷⁰ The final signal generated is a measure of microcirculatory red cell flux (the product of red cell concentration and the red cell velocity). Experimental use of laser Doppler flowmetry in vitro and in vivo has consistently shown a close linear correlation between laser Doppler flowmetry flux and CBF measured with standard methods, and the laser light used does not seem to alter the morphological and physiological characteristics of the vascular bed examined.⁷¹⁻⁷³ The method has shown particular use in the observation of changes in microcirculatory flow induced by physiological and pharmacological stimuli.⁷⁴ However, laser Doppler flowmetry is not quantitative, records from a small tissue volume, and provides no information on the direction of blood flow. Further, experience has shown that the flux signal is very sensitive to the artefacts of local tissue pressure and movement, so that the reliability of the tech-

Figure 7 Relation between real CPP (x axis) and estimated CPP (y axis) in 42 patients with head injury. Graph shows that the method overestimates real CPP for pressure under 60 mm Hg. From Czosnyka et al.²⁵

Figure 8 Laser Doppler flowmetry in patients with head injury: Left: closely correlated changes in middle cerebral artery flow velocity and cortical laser Doppler capillary flow during plateau wave activity. From Kirkpatrick et al.¹⁰ Right: uncoupled "big" and "small tube" flow in head injury. FV = blood flow velocity in the middle cerebral artery. LDF = cortical laser Doppler flux.

Figure 9 Effect of mannitol infusion on laser Doppler flux (upper) and estimated cerebrovascular resistance (eCVR calculated using relative changes in cortical laser Doppler signal—lower graph) after a 200 ml bolus of 20% mannitol compared with a 200 ml bolus of normal saline.

nique is critically dependent on the method of application.⁵⁵ Despite these drawbacks, laser Doppler flowmetry has already shown potential for blood flow measurements in several clinical disciplines including neurosurgery.^{10 75-77}

Using rigid support bolts which are fixed into the skull, reliable long term recordings from ventilated patients with head injury has been achieved.¹⁰ Figure 8 (left) shows an example of a recording captured from one patient with a severe diffuse head injury during spontaneous changes in CPP. Fluctuations in ICP resulted in variations in CPP which were accompanied by changes in the laser Doppler flowmetry signal. Although cerebral events resulting in changes of CPP usually cause very similar trends in middle cerebral artery flow velocity, and laser Doppler flowmetry, uncoupling between flow velocity and laser Doppler flowmetry can occur (fig 8 (right)).

The advantage of laser Doppler flowmetry is that the technique provides a real time measure of relative changes in capillary perfusion which is particularly suitable for assessing the microcirculatory response to a therapeutic challenge. Figure 9 shows the effect of mannitol infusion on cerebral microcirculatory flow in patients with head injury indicating a fall in cerebrovascular resistance with increased red cell flow independent of any significant change in CPP.

Jugular venous oximetry

Cerebral tissues can tolerate limited changes in CPP without compromising neuronal function. A fall in CPP to below certain thresholds results in loss of neuronal electrical activity and, on further reduction, loss of membrane stability resulting in cell death.¹ The critical thresholds at which damage ensues is difficult to ascertain in a patient, and may be different

with differing pathologies. A measure of the metabolic response to cerebral hypoperfusion can be obtained by measuring the oxygen and metabolite concentrations in the venous effluent from the brain.^{15 16 78} This is usually achieved by passing a cannula or optic fibre into the jugular vein and subsequently into the jugular bulb.^{15 16 78} The second allows continuous measurement of Sjo₂. Intermittent measures of arteriovenous oxygen and lactate difference provides additional measures of the cerebral oxygenation and metabolic state.⁷⁸ However, the technique is notoriously difficult and a high percentage of Sjo₂ readings are erroneous due to complications of catheter position, impaction, or thrombus formation.⁷⁸⁻⁸¹ Further, it is still unclear as to which side should be cannulated.⁸² Consequently, other cerebral variables need to be monitored to assist in the interpretation of changes in Sjo₂. Despite these difficulties the method has added support to the notion that thresholds of CPP exist below which the cerebral oxygenation and metabolic state deteriorates, and that such states are associated with a poor outcome.³¹

Near infrared spectroscopy

Near infrared spectroscopy is a non-invasive method which attempts to measure cerebral concentrations of oxyhaemoglobin and deoxyhaemoglobin by observing the absorption of near infrared light. The method has been used most extensively in the neonate where interpterional transmission of light (transmission spectroscopy) can occur.^{83 84} An absolute measure of changing brain haemoglobin saturation and blood volume is possible, and has provided a means of monitoring the cerebrovascular response to certain therapeutic manipulations in critically ill infants.⁸⁵

In adults, scattering of light during passage through a greater thickness of tissue prevents adequate transmission of light to the opposite side of the skull.⁸⁶⁻⁸⁸ Thus scattered light has to be sampled by a receiving probe placed ipsilateral to the source probe (reflectance spectroscopy). This results in limited topographical resolution, as it is not clear to what depth near infrared light penetrates the adult brain. Further, the thicker extracranial tissue will influence the sampled signal to a greater proportion when near infrared spectroscopy is used in adults, and the significance of this remains unresolved.⁸⁹ Hence, although an estimate of the light path length transgressed is possible, the use of near infrared spectroscopy in the adult brain is presently considered non-quantitative. Despite these concerns, the technique has been used to demonstrate predictable physiological changes in cerebral oxyhaemoglobin and deoxyhaemoglobin content during respiration,⁹⁰ in response to various manoeuvres such as a CO₂ stress test,⁹¹ and in response to internal carotid artery cross clamping during carotid endarterectomy.⁹²

Experience using near infrared spectroscopy in the neurointensive care setting is limited due in part to the practical difficulties of main-

Figure 10 Two successive episodes of peripheral desaturation occur in this head injured patient as recorded with peripheral pulse oximetry. The second event arrowed shows the SaO_2 falling to below 80% and is accompanied by a fall in oxyhaemoglobin and reciprocal rise in deoxyhaemoglobin. The SjO_2 monitor failed to register this event. From Kirkpatrick et al.¹¹ ICP = intracranial pressure (mm Hg); CPP = cerebral perfusion pressure (mm Hg); FV = right middle cerebral artery flow velocity (cm/s); LDF = laser Doppler flux from the right frontal region (AU). NIRS = near infrared spectroscopy (recording from the right frontal region); HbO_2 = oxyhaemoglobin ($\mu\text{mol/l}$); Hb = deoxy haemoglobin ($\mu\text{mol/l}$); tHb = total haemoglobin ($\mu\text{mol/l}$); (SjO_2 = right jugular venous oxygen saturation (%)). SaO_2 = peripheral oxygen saturation (%).

taining probe positioning long term. However, our own experience indicates that it can provide warning of a fall in cerebral oxygenation with greater sensitivity than jugular venous oximetry (fig 10).¹¹ In addition, calculations of total haemoglobin allow characterisation of different events causing a fall in CPP (hyper-

aemia *v* primary increases in ICP (fig 11)). The advantage of a non-invasive technique which provides estimates of cerebral oxygenation with high temporal resolution^{11 93} is clear to all those interested in monitoring cerebrovascular status, but the use of near infrared spectroscopy in adults is in a state of evolution requiring considerable efforts with future clinical validation studies.

Intraparenchymal probes

Direct measurement of substances in brain tissue has been slow to evolve in the clinical setting for practical and ethical reasons. However, modern probes can now be placed with minimal risks of added morbidity and experience is accumulating. The measurement of relevant chemicals (such as excitatory amino acids) in traumatised adult brain using microphoresis techniques indicate similar chemical profiles to those seen in experimental animals.⁹⁴ Similarly, oxygen measuring electrodes can be employed for direct measurement of parenchymal oxygen concentration and are beginning to provide novel information regarding the response of the cerebral tissues to specific manoeuvres such as hyperventilation.^{95 96} However, whether such focal measures will provide a sufficiently accurate estimation of the condition of the brain for targeted treatment remains to be seen.

The future for multimodality monitoring

All the aforementioned techniques have largely evolved independently of each other and have all identified their own limitations and artefacts. However, the disadvantages of one modality does not necessarily overlap with

Figure 11 Examples of different types of ICP waveform detected using multimodality monitoring methods. From Kirkpatrick et al.¹¹ Left: NIRS signals register a fall in HbO_2 (arrowed) and rise in Hb indicating cerebral oxygen desaturation confirmed by the fall in SjO_2 . The CPP, FV and LDF also fall indicating primary intracerebral hypertension with a secondary fall in cerebral blood flow. Right: In this event (arrowed) increases in tHb , FV, and LDF all indicate that the rise in ICP was due to cerebral hyperaemia. Abbreviations as in fig 10.

those of another. As a result there has been a growing tendency to adopt a multimodality approach to patients in neurointensive care which allows a more informed interpretation of an individual patient's cerebral state and helps to identify artefacts. Thus an event characterised by changes in several monitored variables (fig 11) provides credence to the finding. As experience gathers we anticipate that certain modalities will eventually evolve as those providing the essential key information. If the errors provided by these selected modalities are acceptable, the system can be trimmed for simplicity and reliability, features which are clearly necessary before these systems gain a wider clinical acceptance.

The advantage of multimodality monitoring is the increased power of interpretation. Most patients with brain injury are treated according to general principles maintaining low ICP and adequate CPP. With increasing experience we are learning to recognise situations in which general principles of treatment may at best be inappropriate, at worst detrimental. We envisage that the future management for brain injury will become more precise and increasingly dependent on monitored information gathered real time.

- 1 Sjeso BK. Pathophysiology and treatment of focal cerebral ischaemia. Parts I and II. *J Neurosurg* 1992;77: 169-84;337-54.
- 2 Macpherson P, Graham DI. Arterial spasm and slowing of the cerebral circulation in the ischaemia of head injury. *J Neurol Neurosurg Psychiatry* 1973;48:560-4.
- 3 Marion DW, Darby J, Yonas H. Acute regional cerebral blood flow changes caused by severe head injuries. *J Neurosurg* 1991;74:407-14.
- 4 Goncalves JM, Vaz R, Cerezo A, et al. HM-PAO SPECT in head trauma. *Acta Neurochir (Wein)* 1994;55: 11-13.
- 5 Obrist WD, Wilkinson WE. Regional cerebral blood flow measurement in humans by xenon-133 clearance. *Cerebrovasc Brain Metab Rev* 1990;2:283-327.
- 6 Pickard JD, Czosnyka M. Management of raised intracranial pressure. *J Neurol Neurosurg Psychiatry* 1993;56: 845-58.
- 7 Cruz J. On-line monitoring of global cerebral hypoxia in acute brain injury. Relationship to intracranial hypertension. *J Neurosurg* 1993;79:228-33.
- 8 Chan KH, Dearden NM, Miller JD, et al. Multimodality monitoring as a guide to treatment of intracranial hypertension after severe brain injury. *Neurosurgery* 1993; 32:547-52.
- 9 Muir JK, Boerschel M, Ellis EF. Continuous monitoring of posttraumatic cerebral blood flow using laser-Doppler flowmetry. *J Neurotrauma* 1992;9:355-62.
- 10 Kirkpatrick PJ, M Czosnyka, P Smielewski, et al. Continuous monitoring of cortical perfusion using Laser Doppler flowmetry in ventilated head injured patients. *J Neurol Neurosurg Psychiatry* 1994;57:1382-88.
- 11 Kirkpatrick PJ, Smielewski P, Czosnyka M, et al. Provisional observations with near infrared spectroscopy in head injured patients. *J Neurosurg* 1996 (in press).
- 12 Marmarou A, Anderson RL, Ward JD, et al. Impact of ICP instability and hypotension in patients with severe head trauma. *J Neurosurg* 1991;75:859-66.
- 13 Miller JD. Head injury. *J Neurol Neurosurg Psychiatry* 1993;56:440-7.
- 14 Gold GE. Does acute hyperventilation provoke cerebral oligoemia in comatose patients after acute head injury? *Acta Neurochir (Wein)* 1989;96:100-6.
- 15 Czosnyka M, Whitehouse H, Smielewski P, et al. Computer supported multimodal monitoring in neuro intensive care. *International Journal of Clinical Monitoring and Computing* 1994;11:223-32.
- 16 Cruz J, Miner ME, Allen SJ, et al. Continuous monitoring of cerebral oxygenation in acute brain injury: assessment of cerebral hemodynamic reserve. *Neurosurgery* 1991;29: 743-49.
- 17 Gopinath PS, Robertson CS, Contant CF, et al. Jugular venous desaturation and outcome after head injury. *J Neurol Neurosurg Psychiatry* 1994;57:717-23.
- 18 Robertson CS, Simpson RK Jr. Neurophysiologic monitoring of patients with head injuries. *Neurosurg Clin N Am* 1991;2:285-99.
- 19 Szwedzkiowski J, Dytko P, Kunicki A, et al. Method of estimating intracranial decompensation in man. *J Neurosurg* 1976;45:155.
- 20 Gaab M, Ottens M, Busche F, et al. Routine computerized neuromonitoring. In: Miller JD, Teasdale GM, et al, eds. *ICP VI*. Berlin: Springer Verlag, 1986:240-7.
- 21 Newell DW, Aaslid R, Stooss R, Reulen HJ. The relationship of blood flow velocity fluctuations to intracranial pressure B waves. *J Neurosurg* 1992;76:415-21.
- 22 Lee WM. Acute liver failure. *Am J Med* 1994;96(1A): 3S-9S.
- 23 Plandsoen WCG, Jong DA de, Maas AIR, et al. Fontanelle pressure monitoring in infants with the Rotterdam teletransducer: A reliable technique. *Med Prog Technol* 1987;13:261-71.
- 24 Reid A, Marchbanks RJ, Martin R, et al. Mean intracranial pressure monitoring by an audiological technique—a pilot study. *J Neurol Neurosurg Psychiatry* 1989;52: 610-2.
- 25 Czosnyka M, Kirkpatrick P, Guazzo E, et al. Can TCD pulsatility indices be used for a non-invasive assessment of cerebral perfusion pressure in head injured patients? In: Nagai H, Kamiya K, Ishii S, eds. *Intracranial pressure IX*. Berlin: Springer Verlag, 1994:146-9.
- 26 Gaab MR, Heissler HE, Ehrhardt K. Physical characteristics of various methods for measuring ICP. In: Hoff JT, Betz AL, eds. *Intracranial pressure VII*. Berlin: Springer Verlag, 1989:16-21.
- 27 Gopinath SP, Robertson CS, Narayan RG, et al. Evaluation of a microsensor intracranial pressure transducer. In: Nagai H, Kamiya K, Ishii S, eds. *Intracranial pressure IX*. Berlin: Springer Verlag, 1994:2-5.
- 28 Luerssen TG, Shields PF, Vos HR, et al. Clinical experience with fiber optic brain parenchymal pressure monitor. In: Hoff JT, Betz AL, eds. *Intracranial pressure VII*. Berlin: Springer Verlag, 1989:35-7.
- 29 Marmarou A, Tsuji O, Dunbar JG. Experimental evaluation of a new solid state ICP monitor. In: Nagai H, Kamiya K, Ishii S, eds. *Intracranial pressure IX*. Berlin: Springer Verlag, 1994:15-19.
- 30 Czosnyka M, Czosnyka Z, Pickard JD. Laboratory testing of three intracranial pressure microtransducers—a technical report. *Neurosurgery* 1996 (in press).
- 31 Chan KH, Miller DJ, Dearden M, et al. The effect of changes in cerebral perfusion pressure upon middle cerebral artery blood flow velocity and jugular bulb venous oxygen saturation after severe brain trauma. *J Neurosurg* 1992;77:55-61.
- 32 Czosnyka M, Guazzo E, Iyer V, et al. Testing of cerebral autoregulation by waveform analysis of blood flow velocity and cerebral perfusion pressure. *Acta Neurochir suppl (Wein)* 1994;60:468-71.
- 33 Contant CF, Robertson CS, Gopinath SP, et al. Determination of clinically important thresholds in continuous monitored patients with head injury. *J Neurotrauma* 1992;10 (suppl 1):S57.
- 34 Jones PA, Andrews PJD, Midgley S, et al. Assessing the burden of secondary insults in head injured patients during intensive care. *J Neurol Neurosurg Psychiatry* 1993; 56:571-2.
- 35 Chambers IR, Mendelow AD. Receiver operator characteristic (ROC) curve analysis applied to CPP, ICP and outcome in severely head injured patients. In: Nagai H, Kamiya K, Ishii S, eds. *Intracranial pressure IX*. Berlin: Springer Verlag, 1994:64-7.
- 36 McGraw P. A cerebral perfusion pressure greater than 80 mmHg is more beneficial. In: Hoff JT, Betz AL, eds. *Intracranial pressure VII*. Berlin: Springer Verlag, 1989:839-41.
- 37 Czosnyka M, Kirkpatrick PJ, Guazzo E, et al. Assessment of the autoregulatory reserve using continuous CPP and TCD blood flow velocity measurement in head injury. In: Nagai H, Kamiya K, Ishii S, eds. *Intracranial pressure IX*. Berlin: Springer Verlag, 1994:593-4.
- 38 Wong FC, Piper IR, Miller JD. Waveform analysis of blood pressure, intracranial pressure and transcranial Doppler signals and their relationship to cerebral perfusion pressure in head injured patients. In: Nagai H, Kamiya K, Ishii S, eds. *Intracranial pressure IX*. Berlin: Springer Verlag, 1994:144-5.
- 39 Avezaat CJJ, van Eijndhoven JHM. Thesis. The Hague: A Jongbloed and Zoon Publishers, 1984.
- 40 Price JD, Czosnyka M, Williamson M. Attempts to continuously monitor autoregulation and compensatory reserve in severe head injuries. In: Avezaat CJJ, van Eijndhoven JHM, Maas AIR, Tans JTJ, eds. *Intracranial pressure VIII*. Berlin: Springer-Verlag, 1993:61-6.
- 41 Dearden NM, Miller JD. Paired comparison of hypnotic and osmotic therapy in the reduction of raised intracranial pressure after severe head injury. In: Hoff JT, Betz AL, eds. *Intracranial pressure VII*. Berlin: Springer-Verlag, 1989:474-81.
- 42 Piper I, Miller JD, Dearden M, et al. System analysis of cerebrovascular pressure transmission: an observational study in head injured patients. *J Neurosurg* 1990;73: 871-80.
- 43 Portnoy HD, Chopp M, Branch C, et al. Cerebrospinal fluid pulse pressure waveform as an indicator of cerebral autoregulation. *J Neurosurg* 1982;56:666-78.
- 44 Czosnyka M, Wollk-Laniewski P, Batorski L. Analysis of intracranial pressure waveform during infusion test. *Acta Neurochir (Wein)* 1989;93:140-5.
- 45 Czosnyka M, Price JD, Williamson M. Monitoring of cerebrospinal dynamics using continuous analysis of intracranial pressure and cerebral perfusion pressure in head injury. *Acta Neurochir (Wein)* 1994;126:113-9.
- 46 Robertson CS, Narayan RK, Contant CF, et al. Clinical experience with a continuous monitor of intracranial

- compliance. *J Neurosurg* 1989;71:673-80.
- 47 Ryder HW, Epsey FP, Kimbell FD. The mechanism of the change in cerebrospinal fluid pressure following an induced change in the volume of the fluid space. *J Lab Clin Med* 1953;41:428-35.
 - 48 Langfitt TW, Weinstein JD, Kassell NF. Cerebral vasomotor paralysis produced by intracranial pressure. *Neurology* 1965;15:622-41.
 - 49 Czosnyka M, Laniewski P, Batorski L, et al. Remarks on amplitude-pressure characteristic phenomenon. In: Hoff JT, Betz AL, eds. *Intracranial pressure VII*. Springer Verlag, 1989:255-9.
 - 50 Czosnyka M, Guazzo E, Kirkpatrick P, et al. Prognostic value of the ICP pulse waveform analysis after severe head injury. In: Nagai H, Kamiya K, Ishii S, eds. *Intracranial pressure IX*. Berlin: Springer Verlag, 1994:200-3.
 - 51 Aaslid R. Cerebral hemodynamics. In: Newell DW, Aaslid R, eds. *Transcranial Doppler*. New York: Raven Press Ltd, 1992:49-58.
 - 52 Kontos HA. Validity of cerebral arterial blood calculations from velocity measurements. *Stroke* 1989;20:1-3.
 - 53 Nelson RJ, Czosnyka M, Pickard JD, et al. Experimental aspects of cerebrospinal haemodynamics: the relationship between blood flow velocity waveform and cerebral autoregulation. *Neurosurgery* 1992;31:705-10.
 - 54 Czosnyka M, Richards H, Kirkpatrick P, Pickard J. Assessment of cerebral autoregulation using ultrasound and laser doppler waveforms—an experimental study in anaesthetized rabbits. *Neurosurgery* 1994;35:287-93.
 - 55 Richards HK, Czosnyka M, Kirkpatrick P, Pickard JD. Estimation of laser doppler flux biological zero using basilar artery flow velocity in the rabbit. *Am J Physiol* 1995;268:H213-17.
 - 56 Czosnyka M, Richards H, Pickard JD, Iyer V. Frequency-dependent properties of cerebral blood transport—An experimental study in rabbits. *Ultrasound Med Biol* 1994;20:391-9.
 - 57 Hassler W, Steinmetz H, Gawlowski J. Transcranial Doppler ultrasonography in raised intracranial pressure and intracranial circulatory arrest. *J Neurosurg* 1988;68:745-51.
 - 58 Aaslid R, Lindegaard KF, Sorteberg W, Helge N. Cerebral autoregulation dynamics in human. *Stroke* 1989;20:45-52.
 - 59 Giller CA. A bedside test for cerebral autoregulation using transcranial Doppler ultrasound. *Acta Neurochir (Wein)* 1991;108:7-14.
 - 60 Czosnyka M, Pickard J, Whitehouse HS. The hyperemic response to a transient reduction in cerebral perfusion pressure—a modelling study. *Acta Neurochir (Wien)* 1992;15:90-7.
 - 61 P Smielewski, M Czosnyka, V Iyer, et al. Computerised transient hyperaemic response test—a method for the assessment of cerebral autoregulation. *Ultrasound in Medical Biology* 1995;21:599-611.
 - 62 Newell DW, Aaslid R, Stooss R, Reulen HJ. Evaluation of closed head injury patients using transcranial Doppler monitoring. In: Avezaat CJJ, van Eijndhoven JHM, Maas AIR, Tans JTJ, eds. *Intracranial pressure VIII*. Berlin: Springer-Verlag, 1993:309-12.
 - 63 Grosset DG, Strebel S, Straiton J, et al. Impaired carbon dioxide reactivity predicts poor outcome in severe head injury: a transcranial Doppler study. In: Avezaat CJJ, van Eijndhoven JHM, Maas AIR, Tans JTJ, eds. *Intracranial pressure VIII*. Berlin: Springer-Verlag, 1993:322-6.
 - 64 Chan KH, Dearden M, Miller JD. The significance of post-traumatic increase in cerebral blood flow velocity: a transcranial Doppler ultrasound study. *Neurosurgery* 1992;30:697-700.
 - 65 Compton JS, Teddy PJ. Cerebral arterial vasospasm following severe head injury—a transcranial Doppler study. *Br J Neurosurg* 1987;1:435-9.
 - 66 Weber M, Grolimund P, Seiler RW. Evaluation of post-traumatic cerebral blood flow velocity by transcranial Doppler ultrasonography. *Neurosurgery* 1990;27:106-12.
 - 67 Chan KH, Dearden M, Miller JD, et al. Transcranial Doppler waveform differences in hyperemic and nonhyperemic patients after severe head injury. *Surg Neurol* 1992;32:433-6.
 - 68 Haberl RL, Heizer ML, Ellis EF. Laser-Doppler assessment of brain microcirculation: effect of systemic alterations. *Am J Physiol* 1989;256:H1247-54.
 - 69 Haberl RL, Heizer ML, Ellis EF. Laser-Doppler assessment of brain microcirculation: effect of local alterations. *Am J Physiol* 1989;256:H1255-60.
 - 70 Shepherd AP, Riedel GL, Kiel JW, et al. Evaluation of an infrared laser Doppler blood flowmeter. *Am J Physiol* 1987;252:G832-9.
 - 71 Skarphedinsson JO, Harding H, Thoren P. Repeated measurements of cerebral blood flow in rats. Comparisons between the hydrogen clearance method and laser Doppler flowmetry. *Acta Physiol Scand* 1988;134:133-42.
 - 72 Eyre JA, Essex JTH, Flecknell PA, et al. A comparison of measurements of cerebral blood flow in the rabbit using laser Doppler spectroscopy and radionuclide labelled microspheres. *Clin Phys Physiol Meas* 1988;9:65-74.
 - 73 Florence G, Seylaz J. Rapid autoregulation of cerebral blood flow: a laser-doppler flowmetry study. *J Cereb Blood Flow Metab* 1992;12:674-80.
 - 74 Skarphedinsson JO, Delle M, Hoffman P, Thoren P. The effects of naloxone on cerebral blood flow and cerebral function during relative cerebral ischemia. *J Cereb Blood Flow Metab* 1989;9:515-22.
 - 75 Fasano VA, Urciuoli R, Bolognese P, Mostert M. Intraoperative use of laser Doppler in the study of cerebral microvascular circulation. *Acta Neurochir (Wein)* 1988;95:40-8.
 - 76 Meyerson BA, Gunasekera L, Linderth B, Gazeliu B. Bedside monitoring of regional cortical blood flow in comatose patients using laser Doppler flowmetry. *Neurosurgery* 1991;29:50-5.
 - 77 Rosenblum BR, Bonner RF, Oyfield EH. Intraoperative measurement of cortical blood flow adjacent to cerebral AVM using laser Doppler velocimetry. *J Neurosurg* 1987;66:396-9.
 - 78 Andrews PJD, Dearden NM, Miller JD. Jugular bulb cannulation: description of a cannulation technique and validation of a new continuous monitor. *Br J Anaesthesiol* 1991;67:553-8.
 - 79 Cruz J. Contamination of jugular bulb venous oxygen measurements. *J Neurosurg* 1992;77:975-6.
 - 80 Sheinberg M, Kanter MJ, Robertson CS, et al. Continuous monitoring of jugular venous oxygen saturation in head-injured patients. *J Neurosurg* 1992;76:212-7.
 - 81 Dearden NM, Midgley S. Technical considerations in continuous jugular venous oxygen saturation measurement. *Acta Neurochir suppl (Wein)* 1993;59:91-7.
 - 82 Stocchetti N, Paparella A, Bridelli F, et al. Cerebral venous oxygen saturation studied with bilateral samples in the internal jugular veins. *Neurosurgery* 1994;34:38-44.
 - 83 Cope M, Delpy DT. A system for long term measurement of cerebral blood and tissue oxygenation in newborn infants by near infrared transillumination. *Med Biol Eng Comput* 1988;26:289-94.
 - 84 Wyatt JS, Cope M, Delpy DT, et al. Quantification of cerebral oxygenation and haemodynamics in sick newborn infants by near infrared spectrophotometry. *Lancet* 1986;ii:1063-6.
 - 85 Edwards AD, McCormick DC, Roth SC, et al. Cerebral hemodynamic effects of treatment with modified natural surfactant investigated by near infrared spectroscopy. *Pediatr Res* 1992;32:532-6.
 - 86 Delpy DT, Cope M, van der Zee P, et al. Estimation of optical pathlength through tissues by direct time of flight measurement. *Phys Med Biol* 1988;33:1433-42.
 - 87 Elwell CE, Cope M, Edwards AD, et al. Measurement of cerebral blood flow in adult humans using near infrared spectroscopy—methodology and possible errors. *Adv Exp Med Biol* 1992;317:235-4.
 - 88 Hampson NB, Camporesi EM, Stolp BW, et al. Cerebral oxygen availability by NIRS spectroscopy during transient hypoxia in humans. *J Appl Physiol* 1990;69:907-13.
 - 89 Germon TJ, Kane NM, Manara AR, et al. Near infrared spectroscopy in adults: effects of extracranial ischaemia and intracranial hypoxia on estimation of cerebral oxygenation. *J Anaesthesiol* 1994;73:503-6.
 - 90 Elwell CE, Owen-Reece H, Cope M, et al. Measurement of changes in cerebral haemodynamics during inspiration and expiration using near infrared spectroscopy. *Adv Exp Med Biol* 1993;345:619-26.
 - 91 Smielewski P, Minhas P, El Zayat S, et al. Cerebral CO₂ reactivity testing using TCD ultrasound and near-infrared spectroscopy—study in normal volunteers. *Cerebrovascular Diseases* 1994;4(suppl 3):23.
 - 92 Kirkpatrick PJ, Smielewski P, Whitfield P, et al. An observational study of near infrared spectroscopy during carotid endarterectomy. *J Neurosurg* 1995;82:756-63.
 - 93 Kato T, Kamei A, Takashima S, Ozaki T. Human visual cortical function during photic stimulation monitoring by means of near infrared spectroscopy. *J Cereb Blood Flow Metab* 1993;13:516-20.
 - 94 Bullock R, Zauner A, Tsuji O, et al. Excitatory amino acid release after severe human head trauma: effect of intracranial pressure and cerebral perfusion pressure changes. In: Nagai H, Kamiya K, Ishii S, eds. *Intracranial pressure IX*. Berlin: Springer-Verlag, 1994:264-7.
 - 95 Mass AIR, Fleckenstein W, De Jong DA, et al. Monitoring cerebral oxygenation: experimental studies and preliminary clinical results of continuous monitoring of cerebrospinal fluid and brain tissue oxygen tension. *Acta Neurochir Suppl (Wein)* 1993;59:50-7.
 - 96 Von Santbrink H, Maas AIR, Avezaat CJJ. Continuous monitoring of brain tissue pO₂ after severe head injury. In: Nagai H, Kamiya K, Ishii S, eds. *Intracranial pressure IX*. Berlin: Springer-Verlag, 1994:582-4.