SHORT REPORT

Treatment of acute pandysautonomia with intravenous immunoglobulin

Robert A Mericle, William J Triggs

Abstract

Acute pandysautonomia has been suggested to be an uncommon variant of Guillain-Barré syndrome. Acute pandysautonomia does not seem to have been treated with intravenous immunoglobulin or other therapies proved efficacious in Guillain-Barré syndrome. A patient is reported with severe acute pandysautonomia who responded dramatically to intravenous immunoglobulin. The findings are consistent with a dysimmune pathogenesis for this syndrome and suggest a possible treatment for future cases.

(*J Neurol Neurosurg Psychiatry* 1997;62:529-531)

Keywords: acute pandysautonomia; intravenous immunoglobulin; autonomic nervous system disease

Acute pandysautonomia was first described by Young *et al* in 1969.¹ Since this time, 27 cases have been reported.¹⁻²⁴ The disorder presents with clinical features reminiscent of the autonomic failure of severe Guillain-Barré syndrome, including severe orthostasis, impairment of gastrointestinal motility and bladder function, impotence, impairment of pupillary reactivity and accommodation, and dryness of the eyes, nasopharynx, and skin.²⁵⁻²⁷ Recovery occurs slowly and is often incomplete. Commonly, patients are left with disabling residual symptoms.^{8 13 15 17 19 24}

Similarity between acute pandysautonomia and the autonomic failure of severe Guillain-Barré syndrome suggests that acute pandysautonomia may be an uncommon variant of Guillain-Barré syndrome.^{1 3 4 6 8 12 14 15 17 19 20 25} This hypothesised relation is strengthened by the existence of cases of acute pandysautonomia with varying degrees of sensory loss and dysaesthesiae.^{5 8 15 16} Raised CSF protein in many cases of acute pandysautonomia is also consistent with this hypothesis.²⁷

Little is known regarding the treatment of acute pandysautonomia. Corticosteroids have been given with disappointing mixed results.^{1 7 9 15 19 20 23 27} More recently, one patient with acute pandysautonomia was treated symptomatically with 400 mg L-threo-3,4-dihydroxyphenylserine (L-DOPS) daily.²¹

This patient required daily symptomatic treatment for more than two years. To our knowledge, however, treatment of acute pandysautonomia with therapies proved efficacious in Guillain-Barré syndrome²⁸⁻³¹ has not been reported. We hypothesised that acute pandysautonomia would respond to treatment with intravenous immunoglobulin (IVIg).

Case report

PRESENTATION

A previously healthy 26 year old woman was admitted to hospital at the University of Florida Health Science Center in November of 1994 with severe acute pandysautonomia. Four weeks earlier, she had developed a generalised rash while receiving trimethoprim/sulfamethoxazole for treatment of a urinary tract infection. She presented with a clinical syndrome characterised by orthostasis, syncope, abdominal pain, anorexia, nausea, vomiting, photophobia, dryness of her mouth, eyes, and skin, diarrhoea, and incontinence of stool. She also complained of burning dysaesthesiae in her feet. Evaluation at an outside hospital disclosed severe orthostatic hypotension without secondary tachycardia, due to absent responses to baroreceptors. One blood pressure recording was 130/70 mm Hg while supine and it dropped to 50/0 mm Hg while standing. Normal laboratory studies included renal and liver functions, complete blood count, erythrocyte sedimentation rate, antinuclear antibodies, serological testing for syphilis, and porphyrins. There were no cells in the CSF; CSF protein was 78 mg/dl and glucose 75 mg/dl. Abdominal CT, a rectal biopsy, nerve conduction studies, and EMG were all normal. An upper gastrointestinal series disclosed pooling of barium in the stomach and much increased transit time. The patient was then transferred to our institution. She had no history of travel, botulism, exposure to toxins, medications, or infections other than that previously mentioned. Family history was negative, and there was no evidence of Adie's, Riley-Day, or Shy-Drager syndromes.

EXAMINATION

Examination showed a photophobic young woman lying supine. On sitting upright, her blood pressure fell from 110/70 to 40/0 mm

Department of Neurology, University of Florida Health Science Center, Gainesville, Florida, USA

R A Mericle
W J Triggs

Correspondence to:
Dr William J Triggs,
Department of Neurology,
Box 100236, JHMHC,
University of Florida,
Gainesville, FL 32610-0236,
USA.

Received 4 November 1996
and in revised form
13 December 1996
Accepted 23 January 1997

Hg and she lost consciousness transiently. Although she lost peripheral pulses, her heart rate remained stable during hypotension (105 bpm supine, 110 bpm standing). Her mouth, pharynx, and skin were very dry. Higher cortical functions were intact. Her pupils were 5 mm bilaterally and unreactive to light or accommodation. Motor examination was normal. Sensory examination was notable for severe hyperaesthesia in the patient's feet. Deep tendon reflexes were normal.

A "cold pressor test" was performed by submerging the patient's hands into a large bucket of ice water for 60 seconds. This test was abnormal, with an absent rise in blood pressure. Nerve conduction studies and concentric needle EMG were normal. The sympathetic skin response³² was absent. A nuclear medicine gastric emptying study showed very reduced gastric emptying with no appreciable response to metoclopramide. ECG telemetry for eight days showed no sign of dysautonomic arrhythmia.

TREATMENT

Intravenous immunoglobulin (IVIg) was given on days 1, 2, 3, 5, and 8 at a dose of 0.4 g/kg/day. Clinical improvement was obvious by day 3 in both subjective and objective measurements. Pupillary responses returned, now constricting from 5 mm to 3 mm bilaterally. This was accompanied by resolution of her photophobia. Her orthostasis improved and she regained heart rate responses to baroreceptors; she was now capable of developing tachycardia in the face of hypotension (supine blood pressure 125/82 mm Hg, heart rate 88 bpm; standing blood pressure 82/42 mm Hg, heart rate 126 bpm). She reported resolution of postural lightheadedness. Her mouth and pharynx became moist and sweat appeared on her skin. The incontinence

resolved. On days 4, 6, and 7, in between IVIg treatments, we noted transient worsening of dysaesthesiae. The dysaesthesiae had disappeared by the completion of IVIg treatment. Nausea and anorexia resolved over the next few weeks. The patient was discharged after 16 days in hospital. On follow up over one year later, examination was notable only for milder, asymptomatic orthostatic hypotension (supine blood pressure 117/78 mm Hg, heart rate 85 bpm; standing blood pressure 104/62 mm Hg, heart rate 98 bpm). She was living independently with no complaints and no symptoms.

Discussion

This patient showed dramatic improvement of acute pandysautonomia temporally associated with administration of IVIg. It is conceivable that symptomatic improvement in our patient was merely coincident with the IVIg. Indeed, spontaneous resolution of acute pandysautonomia is not infrequent.^{1-7 9-12 14-16 18 20-23} However, our analysis of previously reported cases (table) showed that recovery from this condition is usually protracted and incomplete. Partial recovery in previously reported cases of acute pandysautonomia occurred after an average of 30 months and residual disability was not uncommon. By contrast, our patient improved rapidly after receiving IVIg and recovered much more quickly than previously reported cases. Given the natural history of acute pandysautonomia, we suggest that improvement in our patient was not spontaneous, but was likely related to administration of IVIg.

To our knowledge, this case represents the first time that acute pandysautonomia has been treated with therapy proved efficacious in Guillain-Barré syndrome.²⁸⁻³¹ Plasma exchange is associated with various medical complications, including hypotension and cardiac arrhythmia.³³⁻³⁴ Intuitively, plasma exchange imposes an even greater risk of these complications in patients with severe dysautonomia. By contrast, administration of IVIg in this setting probably poses considerably less risk. The extremely low incidence of acute pandysautonomia makes a clinical trial of treatment of this disorder unlikely. However, our findings suggest that acute pandysautonomia is in fact a variant of Guillain-Barré syndrome. We suggest that early treatment of IVIg deserves consideration in cases of this syndrome.

We have no commercial or proprietary interest in the items discussed in this manuscript

Previously reported cases of acute pandysautonomice

Case No	Time until recovery (months)	Authors
1	18	Young <i>et al</i> 1969 ¹
2	28	Andersen <i>et al</i> 1972 ²
3	192	Apenzeller and Kornfeld 1973 ³
4	6	Bost <i>et al</i> 1983 ⁴
5	18	Colan <i>et al</i> 1980 ⁵
6	9	Cortelli <i>et al</i> 1990 ⁶
7	60	Ducla-Soares <i>et al</i> 1993 ⁷
8	No recovery/death at 36 months	Fagius <i>et al</i> 1983 ⁸
9	12	Feldman <i>et al</i> 1991 ⁹
10	24	Goulon <i>et al</i> 1971 ¹⁰
11	9	Guidi <i>et al</i> 1981 ¹¹
12	10	Hart and Kanter 1990 ¹²
13	No mention of recovery	Hirayama 1989 ¹³
14	4	Hopkins <i>et al</i> 1974 ¹⁴
15	48	Hopkins <i>et al</i> 1974 ¹⁴
16	60	Low <i>et al</i> 1983 ¹⁵
17	No recovery	Low <i>et al</i> 1983 ¹⁵
18	31	Okada <i>et al</i> 1975 ¹⁶
19	11	Okada <i>et al</i> 1975 ¹⁶
20	No mention of recovery	Sannomiya <i>et al</i> 1989 ¹⁷
21	12	Sekine <i>et al</i> 1991 ¹⁸
22	No recovery / death	Stoll <i>et al</i> 1991 ¹⁹
23	24	Thomashefsky <i>et al</i> 1972 ²⁰
24	24 (but temporary improvement with daily L-DOPS)	Ushiyama <i>et al</i> 1996 ²¹
25	24	Wichser <i>et al</i> 1972 ²²
26	6	Yahr and Frontera 1975 ²³
27	No recovery at 72 months	Yee <i>et al</i> 1976 ²⁴
28	15 days	Mericle and Triggs 1997 our patient

Mean (SD) recovery time was 30 (39) months.

- Young RR, Asbury AK, Adams RD, Corbett JL. Pure pandysautonomia with recovery. *Transactions of the American Neurological Association* 1969;94:355-7.
- Andersen O, Lindberg J, Modigh, Rese-Nielsen E. Subacute dysautonomia with incomplete recovery. *Acta Neurol Scand* 1972;48:510-9.
- Appenzeller O, Kornfeld M. Acute pandysautonomia. *Arch Neurol* 1973;29:334-9.
- Bost M, Rossignol AM, Tachker D, Batellier H, Jeannoel P. A case of acute reversible dysautonomia in an adolescent. *Pediatrics* 1983;38:29-36.
- Colan RV, Snead OC, Oh SJ, Kashlan MB. Acute autonomic and sensory neuropathy. *Ann Neurol* 1980;8:441-4.
- Cortelli P, Contin M, Lugaesi A, Baruzzi A, Montaga P.

- Severe dysautonomic onset of Guillain-Barré syndrome with good recovery. A clinical and autonomic follow-up study. *Ital J Neurol Sci* 1990;11:159-62.
- 7 Ducla-Soares JL, Guerreiro AS, Pova P, Alvares E, Guerreiro L, Carrilho F, et al. Pure autonomic failure. *Acta Med Port* 1993;6:533-7.
 - 8 Fagius J, Westerberg CE, Olsson Y. Acute pandysautonomia and severe sensory deficit with poor recovery. A clinical, neurophysiological, and pathological case study. *J Neurol Neurosurg Psychiatry* 1983;46:725-33.
 - 9 Feldman EL, Bromberg MB, Blaivas M, Junck L. Acute pandysautonomic neuropathy. *Neurology* 1991;41:746-8.
 - 10 Goulon M, Nouailhat F, Grosbuis S, Gajdos P. Hypotension orthostatique à pouls invariable: étude hémodynamique d'une neuropathie amyloïde et d'une forme idiopathique transitoire. *Rev Neurol (Paris)* 1971;125:257-72.
 - 11 Guidi L, Zeppilli P, Sassara M, and Ghirlanda G. Cardiovascular autonomic impairment in one case of acute pandysautonomia with recovery. *G Ital Cardiol* 1981;11:1151-9.
 - 12 Hart RG, Kanter MC. Acute autonomic neuropathy. Two cases and a clinical review. *Arch Intern Med* 1990;150:2373-6.
 - 13 Hirayama K. Acute curable pandysautonomia and chronic progressive pandysautonomia. *Bulletins of the Osaka Medical College* 1989;35:113-7.
 - 14 Hopkins A, Neville B, Bannister R. Autonomic neuropathy of acute onset. *Lancet* 1974;i:769-71.
 - 15 Low PA, Dyck PJ, Lambert EH, Briminjion WS, Trautmann JC, Malagelada JR, et al. Acute panautonomic neuropathy. *Ann Neurol* 1983;13:412-17.
 - 16 Okada F, Yamashita I, Suwa N. Two cases of acute pandysautonomia. *Arch Neurol* 1975;32:146-51.
 - 17 Sannomiya K, Kudo Y, Mori T, Nakazato O, Okajima T. A case of acute idiopathic pandysautonomia. *Rinsho Shinkeigaku* 1989;29:895-900.
 - 18 Sekine T, Tateyama M, Saito H. A case of subacute idiopathic pure pan-dysautonomia-recovery with prednisolone therapy. *Rinsho Shinkeigaku* 1991;31:445-50.
 - 19 Stoll G, Thomas C, Reinert K, Schober R, Hartung HP. Encephalo-myelo-radiculo-ganglionitis presenting as pandysautonomia. *Neurology* 1991;41:723-6.
 - 20 Thomashefsky AF, Horwith SF, Feingold MH. Acute autonomic neuropathy. *Neurology* 1972;22:251-5.
 - 21 Ushiyama M, Ikeda SI, Suzuki T, Yazawa M, Yanagisawa N, Tsujino SI. Acute pandysautonomia: mass spectrometric and histopathological studies of the sympathetic nervous system during long term L-threo-3,4-dihydroxyphenylserine treatment. *J Neurol Neurosurg Psychiatry* 1996;61:99-102.
 - 22 Wichser J, Vijayan N, Dreyfus PM. Dysautonomia: its significance in neurologic disease. *Calif Med* 1972;117:28-37.
 - 23 Yahr MD, Frontera AT. Acute autonomic neuropathy: its occurrence in mononucleosis. *Arch Neurol* 1975;32:132-3.
 - 24 Yee RD, Trese M, Zee DS, Kollarits CR, Cogan DG. Ocular manifestations of acute pandysautonomia. *Am J Ophthalmol* 1976;81:740-4.
 - 25 Lichtenfeld P. Autonomic dysfunction in the Guillain-Barré syndrome. *Am J Med* 1971;50:772-80.
 - 26 Young RR, Asbury AK, Corbett JL, Adams RD. Pure pandysautonomia with recovery. *Brain* 1975;98:613-36.
 - 27 Ropper AH, Wijdicks EFM, Truax BT. *Guillain-Barré syndrome*. Philadelphia: FA Davis, 1991.
 - 28 Osterman PO, Fagius J, Lundemo G, et al. Beneficial effects of plasma exchange in acute inflammatory polyradiculoneuropathy. *Lancet* 1984;ii:1296-9.
 - 29 Guillain-Barré Syndrome Study Group. Plasmapheresis and acute Guillain-Barré syndrome. *Neurology* 1985;35:1096-104.
 - 30 French Cooperative Group on Plasma Exchange in Guillain-Barré syndrome. Efficiency of plasma exchange in Guillain-Barré syndrome: role of replacement fluids. *Ann Neurol* 1987;22:753-61.
 - 31 Van der Meché FG, Schmitz PI. A randomized trial comparing intravenous immune globulin and plasma exchange in Guillain-Barré syndrome. Dutch Guillain-Barré Study Group. *N Engl J Med* 1992;326:1123-9.
 - 32 Shahani BT, Day TJ, Cros D, Khalil N, Kneebone CS. RR interval variation and the sympathetic skin response in the assessment of autonomic function in peripheral neuropathy. *Arch Neurol* 1990;47:659-64.
 - 33 Bouget J, Chevret S, Chastang C, Raphael JC. Plasma exchange morbidity in Guillain-Barré syndrome: results from the French prospective, double-blind, randomized, multicenter study. The French Cooperative Group. *Crit Care Med* 1993;21:651-8.
 - 34 Thornton CA, Griggs RC. Plasma exchange and intravenous immunoglobulin treatment of neuromuscular disease. *Ann Neurol* 1994;35:260-8.