Predictive value of clinical indices in detecting aspiration in patients with neurological disorders

Fabiola Mari, Monica Matei, Maria Gabriella Ceravolo, Anna Pisani, Alfeo Montesi, Leandro Provinciali

Abstract

Objectives—(1) To evaluate the predictive value of a detailed clinical screening of aspiration in patients with neurological diseases, both with and without symptoms of dysphagia taking videofluoroscopy as the gold standard; (2) to assess the existence of risk factors for silent aspiration, measuring the cost-benefit ratio of radiological examination.

Methods—93 consecutive patients meeting the diagnostic criteria for a neurological disease with a risk of swallowing dysfunctions (cerebrovascular accidents, brain injury, Parkinson's disease, multiple sclerosis, amyotrophic lateral sclerosis, myotonic dystrophy, and abiotrophic diseases) underwent a detailed clinical assessment using a 25 item form to check for symptoms of dysphagia and impairment of the oropharyngeal swallowing mechanism. The 3 oz water swallow test was also performed to assess the aspiration risk. Sensitivity, specificity, positive predictive, and negative predictive values (NPV) of dysphagia, history of cough on swallowing, and 3 oz test positivity, versus videofluoroscopy documented aspiration, taken as the gold standard, were measured in all the patients and in subgroups with different neurological disorders.

Results—Non-specific complaints of dysphagia showed a very poor predictive value, whereas the symptom "cough on swallowing" proved to be the most reliable in predicting the risk of aspiration, with 74% sensitivity and specificity, 71% positive predictive, and 77% negative predictive value. The standardised 3-oz test had a higher predictive potential than the clinical signs, but had low sensitivity. The association of cough on swallowing with the 3 oz test gave a positive predictive of 84%, and an negative predictive value of 78%. In cases where the clinical tests failed to detect any impairment, videofluoroscopy documented only a low risk (20%) for mild aspiration.

Conclusions—The association of two clinical items (such as history of cough on swallowing and 3 oz test positivity) provides a useful screening tool, the cost-benefit ratio of which seems very competitive in comparison with videofluoroscopy in aspiration risk evaluation.

Keywords: dysphagia; aspiration risk; bedside clinical assessment

Food aspiration is a frequent consequence of dysphagia, causing a strong risk of pneumonia¹ and diet alterations.² In some neurological diseases, dysphagia can persist for a long period, without being complained of by patients.³ Therefore, a screening of swallowing alterations in such conditions is suggested, if lung infections are to be prevented.

A prospective, longitudinal cohort study, carried out on patients during the subacute phase after stroke, documented that even silent aspiration is associated with a 5.57-fold increase in risk of pneumonia.⁴ Videofluoroscopy showed the occurrence of a swallowing impairment in many different neurological disorders, such as cerebrovascular accidents,⁵ brain injury,⁶ Parkinson's disease,^{7,8} multiple sclerosis,⁹ amyotrophic lateral sclerosis,^{10,11} myotonic dystrophy,¹² and dementia.¹³

Although videofluoroscopy is, so far, the most accurate instrumental investigation to check the passage of food below the vocal folds,¹⁴ it is an invasive test that requires a specialised diagnostic laboratory with trained staff and a specific set of expensive instruments not available in all centres.¹⁵

Several clinical approaches have been proposed to make the diagnostic process easier and to measure the severity of dysphagia.¹⁶⁻¹⁸ In particular, clinical scales were associated with either non-specific¹⁹ or specific pathology tests²⁰ to predict aspiration.

The aim of this study was to evaluate the predictive value of a detailed clinical screening of aspiration in patients with neurological diseases, both with and without symptoms of dysphagia taking videofluoroscopy as the gold standard.²¹

Furthermore the cost-benefit ratio of the radiological examination was evaluated on the basis of such results and risk factors for silent aspiration were investigated.

Methods

PATIENTS

Ninety three consecutive patients (61 inpatients and 32 outpatients; 59 men, 34 women; mean age 59.8 (SD 16), range 18-80) admitted to the rehabilitation clinic over an 18 month period, were screened to evaluate the presence of aspiration. Inclusion criteria were as follows:

(1) presence of a neurological disorder with a documented risk of swallowing dysfunctions⁵⁻¹³;

Nervous System
Disease Institute,
University of Ancona,
Italy

F Mari
M Matei
M G Ceravolo
L Provinciali

Department of
Radiology Umberto I
Hospital of Ancona,
Italy

A Pisani
A Montesi

Correspondence to:
Professor Leandro
Provinciali, Clinica di
Neuroriabilitazione,
Ospedale Torrette, Ancona,
Italy I-60020.

Received 18 January 1997
and in revised form 14 April
1997

Accepted 15 April 1997

(J Neurol Neurosurg Psychiatry 1997;63:456-460)

(2) informed consent to the investigation; (3) ability to cooperate. Unconscious and demented subjects, as well as bedridden patients, were excluded.

The case mix was represented by 28 acute or subacute stroke (cerebrovascular accident) (five total anterior circulation infarct, 10 partial anterior circulation infarct, four posterior circulation infarct, nine lacunar infarct, as classified according to Bamford *et al*²²), seven recent traumatic brain injury, seven multiple sclerosis (expanded disability status scale ranging between 2.0 and 6.0, median: 3.5), 27 Parkinson's disease (Hoehn and Yahr stage ranging from 2.0 to 4.0, median: 3.0), six amyotrophic lateral sclerosis, five myotonic dystrophy, 13 abiotrophic diseases (four Alzheimer's disease, two multisystemic atrophy, two progressive supranuclear palsy, two olivopontocerebellar atrophy, three Friedreich diseases).

Patients with cerebrovascular accident and traumatic brain injury were examined at least one month after the clinical state had stabilised.

Twenty patients (21%) were free from symptoms of swallowing impairment. Among the others, 17 (18%) complained of dysphagia for liquids, 30 (32%) of dysphagia for solids, and 26 of both.

PROCEDURE

The investigation protocol was approved by the local ethics committee.

Patients underwent a detailed clinical examination based on collection of symptoms and evaluation of the oropharyngeal swallowing mechanism.

A 25 item form, drawn up on the basis of both personal experience and previous reports,^{16 23-25} checked for:

(1) past or present dysphagia for either liquids or solids; (2) "knot in throat" sensation; (3) odynophagia; (4) nasal regurgitation; (5) daily episodes of cough on swallowing during the past week; (6) dysphonia; (7) dysarthria; (8) need for either feeding orthosis or nasogastric tube; (9) history of coma or pneumonia; (10) tracheostomy. The checklist also included 14 indices investigating the function of lips, tongue, mandible, and pharynx both at rest and during swallowing.

In addition, the 3 oz water swallow test^{20 26} was carried out by giving each patient 9 ml water and asking them to drink from a cup without interruption. Coughing during or for up to one minute after completion or the presence of a postswallow wet or hoarse voice quality were considered positive signs of the presence of an abnormal situation.

Videofluoroscopy was used for the evaluation and identification of swallowing disorders. This examination was performed within 72 hours of the clinical examination by an observer blind to the clinical findings.^{13 27} A professional video recorder providing 50 i/s frame by frame analyses, slow motion pictures (from 1/30 to five times less normal speed), and forward and backward mode, was used. Patients sitting upright were given 5 or 10 or 15 ml of fluid and solid barium boluses.

Table 1 Variables considered by videofluoroscopy

Swallowing phase	Radiological issues
Oral containment	Anterior leakage Posterior leakage
Horizontal progression	Deficit of horizontal transport Prolonged transit time Deficit of oral clearance
Pharyngeal containment	Deficit of closure of nasal cavity Delay of swallowing reflex Deficit of laryngeal closure Deficit of epiglottis movements
Vertical progression	Vallecular stasis Deficit of pharyngeal peristalsis Deficit of lingual piston Deficit of UES opening Penetration Aspiration severity Mild <10% Moderate <30% Severe ≥30%

The radiological examination was to assess the four phases of bolus deglutition: (1) oral containment; (2) horizontal progression (the transport from the oral cavity to the pharynx); (3) pharyngeal containment (containment, morphological and volumetric adaptation of the pharynx, closure of the nasal, oral, and laryngeal cavities); (4) vertical progression (transport of bolus from the pharynx to the esophagus) (table 1). Aspiration was accurately assessed checking for the entry of bolus below the vocal folds. With respect to the volume of the aspirated bolus (<10%, 10%-30%, or >30% of the total) aspiration was classified as mild, moderate, or severe respectively.

DATA ANALYSIS

Sensitivity, specificity, positive predictive, and negative predictive values of: (1) the patient's complaint of dysphagia(2); a history of cough on swallowing, and (3) positivity of the 3 oz test versus radiologically documented aspiration taken as the gold standard, were measured in all the patients and in subgroups with different neurological disorders. The positive predictive value associated with each clinical item included in the 25 item checklist was also assessed.

Factor analysis was employed to measure the covariance of indices used to build the checklist. To compute the costs of both clinical and radiological approaches the time spent on performing the investigations and the materials employed were considered.

Results

Table 2 shows the prevalence of symptoms and signs checked for during the clinical examination and of 3 oz test positive findings.

Videofluoroscopy documented the occurrence of aspiration in 43 (46%) patients; it was mild in 25, moderate in seven, and severe in 11. Among the remaining patients, 45 showed an impairment in at least one swallowing mechanism.

Table 2 shows the positive predictive values associated with each clinical item with respect to radiologically documented aspiration.

Table 2 Prevalence and positive predictive value (PPV) of positive findings on the 25 item clinical assessment and of 3 oz test, used to check for aspiration risk in neurological patients

Index	Prevalence (%)	PPV (%)
History collection:		
Dysphagia	78	52
Dysarthria	44	58
Dysphonia	57	60
Knot/lump in throat	32	47
Nasal regurgitation	26	75
Cough during swallowing	48	71
Odynophagia	6	33
Coma	10	67
Nasogastric tube	12	54
Tracheostomy	4	75
Pneumonia	18	82
Clinical evaluation:		
Oral phase		
Corner deviation	30	68
Anterior leakage	29	71
Protrusion deficit	42	54
Drooling	28	66
Tongue motility impairment	42	64
Bolus formation deficit	35	55
Oral clearance impairment	41	63
Mandible movement deficit	19	61
Masseter muscle strength deficit	31	66
Oral sensitivity deficit	17	57
Pharyngeal phase		
Swallow reflex impairment	60	54
Soft palate motility deficit	38	63
Hyoid-laryngeal motility	40	60
Pharyngeal reflex impairment	26	57
3 oz test positivity	34	76

Symptoms of dysphagia showed a very poor predictive value. Silent aspiration occurred in four of 20 (20%) patients who did not complain of any swallowing difficulty and in 49% of dysphagic patients. Among the indices in which the positive predictive value was higher than 70%, history of pneumonia, tracheostomy, evidence of anterior leakage, and nasal regurgitation showed a low occurrence rate (less than 30%).

The symptom cough on swallowing proved to be the most reliable in predicting the risk of aspiration, showing high sensitivity, specificity, positive predictive value, and negative predictive value (table 3). The standardised 3 oz test had a higher predictive potential than the clinical signs, but had a low sensitivity. In particular, subjects with silent aspiration gave negative findings in the 3 oz test. For the whole population, the association of cough on swallowing with the 3 oz test gave a positive predictive value of 84%, and a negative predictive value of 78%.

The analysis of subgroups showed how the correlation between symptoms of dysphagia, 3 oz test positivity, and aspiration differed according to the underlying neurological disease (figure), the positive predictive value of the 3 oz test ranging from 50% in patients with

traumatic brain injury to 78% in patients with a cerebrovascular accident. Silent aspiration occurred more often in patients with Parkinson's disease than in other cases.

The analysis of patients with negativity of clinical criteria produced the following "identikit":

(1) complaint of dysphagia in 30% of patients; (2) higher prevalence of Parkinson's disease (48%) than other neurological diseases; (3) low occurrence of pneumonia, tracheostomy, coma, or feeding through nasogastric tube in the medical history (about 2%); (4) impairment in one or more mechanisms of either oral or pharyngeal swallowing phases in 8% to 25% of patients, without any definite pattern; (5) low frequency of positive findings on videofluoroscopy; aspiration occurred in 21% patients, mostly (18%) of a mild type and always associated with evidence of alteration in gag reflex, penetration, and vallecular stasis.

Factor analysis applied to the 25 item check list allowed the detection of seven domains, each depending on one or more clinical indices: (1) dysarthria, dysphonia, and knot or lump in the throat (2); history of coma, nasotracheal tube application, tracheostomy (3); odynophagia (4); lip corner deviation (5); dyskinesia; (6) impairment of mandible movement (7); the association of complaints of dysphagia, nasal regurgitation, and cough on swallowing, history of pneumonia, and evidence of impairment in the oral and pharyngeal phases as detected by the checklist (anterior leakage, protrusion deficit, drooling, tongue motility deficit, bolus formation and clearance deficit, masseter strength deficit, oral sensitivity deficit, swallow and pharyngeal reflex impairment, and soft palate and hyoid-larynx motility deficit).

For the cost:benefit ratio evaluation, the bedside clinical assessment did not employ any expensive equipment and took 20 minutes on average to complete compared with the 55 minutes required by videofluoroscopy including a global radiation exposure time of up to 5.8 minutes (average 3.6 (SD 0.87) min). Both investigations were well tolerated by patients.

Discussion

The role of clinical tests in evaluating swallowing dysfunction and checking aspiration has often been discussed. Logeman²⁸ took the amount of oral intake and the time of feeding as relevant clinical indicators of dysphagia. DePippo *et al*¹⁹ formulated the Burke dysphagia screening test, associating the 3 oz test with other clinical signs, and administered it to 139 inpatients with stroke, concluding that the test is worthwhile in identifying patients at risk of pneumonia or airway obstruction and death. By contrast, Garon *et al*²⁶ found the opposite in 100 patients affected by different diseases and argued that the 3 oz test, which uses the cough reflex as the sole indicator of aspiration, is not a replacement for the precision and accuracy of videofluoroscopic evaluation.

In agreement with Fleming *et al*¹⁶ and Garon *et al*,²⁶ the checklist employed in this study included both history and clinical indices,

Table 3 Reliability of history collection and 3 oz test in checking for aspiration risk in neurological patients (oropharyngeal dysphagia was defined as a subjective difficulty in swallowing liquids or solids, currently or in the past)

	Oropharyngeal dysphagia	Cough	3 oz test	3 oz test + cough
Sensitivity (%)	88	75	52	50
Specificity (%)	30	74	86	91
PPV (%)	52	71	76	84
NPV (%)	75	77	68	76

PPV=positive predictive value; NPV=negative predictive value.

Prevalence of aspiration documented by videofluoroscopy, dysphagia, and 3 oz water swallow test (WST) test positivity in different neurological groups.

selected on the basis of knowledge of the anatomical structures and functional mechanisms involved in swallowing. Factor analysis allowed us to assess how some indices seem poorly related to impairment in swallowing mechanisms. In particular, items such as dysphonia, dysarthria, oral dyskinesiae, mandible movement deficit, and lip corner deviation, may indicate the presence of brainstem lesions, often responsible for, but not necessarily associated with swallowing impairment.²⁹ Both sensation of a knot or lump in the throat and odynophagia may occur in neurotic syndromes and be detected in neurological patients with depressive reactions, hence being dissociated from the detection of an aspiration risk. History of coma, tracheostomy, and use of a nasogastric tube were highly correlated with each other, leading to the identification of a subgroup of patients who had severe brain lesions. Although such a condition is associated with a high risk of aspiration,⁶ the inclusion of the aforementioned indices in a screening assessment is not advisable, owing to their poor sensitivity.

The choice of videofluoroscopic examination as a gold standard for the detection of aspiration was made in accordance with previous investigations^{19–21, 26}; the pattern of swallowing mechanisms was assessed using a procedure already described by other authors.^{27–30}

Patients included in this research were selected on the basis of the risk of aspiration associated with the neurological disease. The occurrence of aspiration in our patients is similar to that described in the literature for patients with cerebrovascular accident,²⁹ traumatic brain injury,^{6, 31–32} and multiple sclerosis.^{9, 33} No definite data are available on the incidence of swallowing alterations in Parkinson's disease (rates ranging between 0% to 100% are reported),^{7, 10, 34} amyotrophic lateral sclerosis, myotonic dystrophy, and abiotrophic diseases. The few cases examined in this study does not allow us to draw any conclusions about the disease specific incidence rates or the higher prevalence of silent aspiration in Parkinson's disease than in other neurological groups.

The evaluation of the predictive value of the clinical examination showed that the complaint of dysphagia by the patients should be considered a non-specific feature, whatever the underlying neurological disease. Despite a prevalence of symptoms ranging from 56% to 100%, true aspiration has been documented in less than one half of dysphagic patients. Although apparently low, such a percentage is still higher than those reported by Johnson *et al*³⁵ and Perlman *et al*³⁶ who found a prevalence of aspiration of 30% and 39% respectively, in different categories of dysphagic patients undergoing videofluoroscopy. The dissociation between severity of symptoms and swallowing alterations may partially reflect the influence of either neurological deficits (for example, post-stroke facial paresis may cause oral stage dysphagia without affecting the pharyngeal stage of swallowing) or psychic problems (anxious patients may complain of swallowing alterations in the absence of any detectable deficit).

Among all the clinical signs considered, only a history of cough on swallowing proved to be really accurate in predicting aspiration, both in the total population and in subgroups with different diseases. Other conditions such as recurrent pneumonia, or previous tracheostomy, although being highly correlated with the risk of aspiration, have no clinical usefulness, owing to their low occurrence rate, as they only affect patients in severe conditions.

The 3 oz test was confirmed as a valuable tool in the evaluation of dysphagic patients, with a positive predictive value of 76% versus risk of aspiration. These findings almost overlap those described by DePippo *et al*²⁰ in patients with stroke, by Garon *et al* in multiple neurological pathology groups,²⁶ and by Splaingard *et al* in patients with stroke or brain injury.²¹

The association of history collection with objective assessment (cough plus 3 oz test) gives an increase in both positive and negative predictive values. A similar conclusion was provided by DePippo *et al*¹⁹ who analysed the predictive significance of items included in the

Burke dysphagia screening test and found that, in patients with stroke, "coughing associated with feeding or during a 3 oz water swallow test could identify 92% of patients developing pneumonia or recurrent upper airway obstruction.

However, the diagnostic accuracy of the clinical evaluation depends on the preservation of the cough reflex and pharynx sensitivity; all pathological conditions which cause either an impairment in pharynx innervation or silent aspiration are unlikely to be detected by the bedside assessment and need radiological screening.

In our experience, the failure of the clinical evaluation prevented prediction of the aspiration risk in a very low percentage of patients, in whom, moreover, aspiration was always mild (<10% bolus) and never associated with a history of lung infections. According to a recent prospective study, only aspiration of more than 10% on barium test swallows during videofluoroscopy is associated with an increased risk of pneumonia.⁴

In conclusion, the association of the 3 oz test with the clinical index cough on swallowing provides a useful screening tool, the cost:benefit ratio of which seems very competitive with videofluoroscopy in evaluation of risk of aspiration.³⁷

Although videofluoroscopy keeps its diagnostic importance in evaluating disability after different failures of the swallowing mechanism, to plan an appropriate treatment,³⁸⁻³⁹ the use of a detailed bedside assessment, based on a multiple item checklist, is advisable for tailoring the rehabilitation approach to each patient's needs¹⁶ in patients in whom radiological examination is not available.

- 1 Horner J. Aspiration following stroke: clinical correlates and outcome. *Neurology* 1988;38:1359-62.
- 2 Martin AW. Dietary management of swallowing disorders. *Dysphagia* 1993;6:129-35.
- 3 Johnston BT, Li Q, Castell JA, Castell DO. Swallowing and esophageal function in Parkinson's disease. *Am J Gastroenterol* 1995;90:1741-6.
- 4 Holas MA, DePippo KL, Reding MJ. Aspiration and relative risk of medical complication following stroke. *Arch Neurol* 1994;51:1051-3.
- 5 Barer DH. The natural history and functional consequences of dysphagia after hemispheric stroke. *J Neurol Neurosurg Psychiatry* 1989;52:236-41.
- 6 Yorkston KM, Honsinger MJ, Mitsuda PM, Hammen V. The relationship between speech and swallowing disorders in head-injured patients. *Journal of Head Trauma Rehabilitation* 1989;4:1-16.
- 7 Bushmann M, Domeyer SM, Lecker L, Perlmutter JS. Swallowing abnormalities and their response to treatment in Parkinson's disease. *Neurology* 1989;39:1309-14.
- 8 Bird MR, Woodward MC, Gibson EM, Philand DJ, Fonda D. Asymptomatic swallowing disorders in elderly patients with Parkinson's disease: a description of findings on clinical examination and videofluoroscopy in sixteen patients. *Age Aging* 1994;23:251-4.
- 9 Hughes JC, Enderby PM, Langton Hewer R. Dysphagia and multiple sclerosis: a study and discussion of its nature and impact. *Clin Rehabil* 1994;8:18-26.
- 10 Robbins J. Swallowing in ALS and motor neuron disease. *Neurol Clin* 1987;5:213-29.
- 11 Welnets K. Maintaining adequate nutrition and hydration in the dysphagia ALS patients. *Canadian Nurse* 1983;79:30-4.
- 12 Willing TN, Paulus J, Lecan Saint Guily J, Beon C. Swallowing problems in neuromuscular disorders. *Arch Phys Med Rehabil* 1994;75:1175-85.
- 13 Feinberg MJ, Ekberg O, Segall L, Tully J. Deglutition in elderly patients with dementia: findings of videofluorographic evaluation and impact on staging and management. *Radiology* 1992;183:811-4.
- 14 Palmer JB, Kuhlmeier KV, Tippet DC, Lynch C. A protocol for the videofluorographic swallowing study. *Dysphagia* 1993;8:209-14.
- 15 Montesi A, Pesaresi A, Antico E, Piloni V. Dynamic radiological study by videorecording of normal oropharyngeal deglutition. *Radiol Med* 1988;75:166-72.
- 16 Fleming SM, Weaver AW. Index of dysphagia: a tool for identifying deglutition problems. *Dysphagia* 1987;1:206-8.
- 17 Sonies BC, Weiffenbach J, Atkinson JC. Clinical examination of motor and sensory functions of the adult oral cavity. *Dysphagia* 1987;1:178-86.
- 18 Martens L, Cameron T, Simonsen M. Effects of a multidisciplinary management program on neurologically impaired patients with dysphagia. *Dysphagia* 1990;5:147-51.
- 19 DePippo KL, Holas MA, Reding MJ. The Burke dysphagia screening test: validation of its use in patients with stroke. *Arch Phys Med Rehabil* 1994;75:1284-6.
- 20 DePippo K, Holas M, Reding M. Validation of 3-Oz water swallow test for aspiration following stroke. *Arch Neurol* 1992;42:1259-61.
- 21 Splaingard ML, Hutchins B, Sultan L, Chaudhuri G. Aspiration in rehabilitation patients: videofluoroscopy v bedside clinical assessment. *Arch Phys Med Rehabil* 1988;69:637-40.
- 22 Bamford J, Sandercock P, Dennis M, Warlow C. Classification and natural history of clinically identifiable subtypes of cerebral infarction. *Lancet* 1991;337:1521-6.
- 23 Cherney LR. Dysphagia in adults with neurologic disorders: an overview. In: Cherney LR, ed. *Clinical management of dysphagia in adults and children*. Gaithersburg: Aspen, 1994: 1-28.
- 24 Bass NH. Clinical signs, symptoms and treatment of dysphagia in the neurologically disabled. *Journal of Neurologic Rehabilitation* 1990;4:227-35.
- 25 Layne KA, Losinski DS, Zenner PM, Ament JA. Using the Fleming index of dysphagia to establish prevalence. *Dysphagia* 1989;4:39-42.
- 26 Garon BR, Engle M, Ormiston C. Reliability of the 3-oz water swallow test utilizing cough reflex as sole indicator of aspiration. *Journal of Neurologic Rehabilitation* 1995;9:139-43.
- 27 Ekberg O. The role of radiology in the evaluation and treatment of neurologically impaired patients with dysphagia. *Journal of Neurologic Rehabilitation* 1990;4:65-73.
- 28 Logeman JA. *Evaluation and treatment of swallowing disorders*. San Diego: College Hill Press, 1983:6.
- 29 Alberts MJ, Horner J, Gray L, Brazer SR. Aspiration after stroke: lesion analysis by brain MRI. *Dysphagia* 1992;7:170-3.
- 30 Linden P, Kuhlmeier KV, Patterson C. The probability of correctly predicting subglottic penetration from clinical observations. *Dysphagia* 1993;8:170-79.
- 31 Logeman J. Evaluation and treatment planning for the head injured patient with oral intake disorders. *Journal of Head Trauma Rehabilitation* 1989;4:24-33.
- 32 Avery Smith W, Dellarosa DM. Approaches to treating dysphagia in patients with brain injury. *Am J Occup Ther* 1994;48:235-9.
- 33 Herrera W, Zeligman BE, Gruber J, Jones MC, Pautler R, Wriston R, et al. Dysphagia in multiple sclerosis: clinical and videofluoroscopic correlation. *Journal of Neurologic Rehabilitation* 1990;4:1-8.
- 34 Leopold NA, Kagel MC. Prepharyngeal dysphagia in Parkinson's disease. *Dysphagia* 1996;11:14-22.
- 35 Johnson ER, McKenzie SW, Rosenquist CJ, Lieberman JS, Sievers AE. Dysphagia following stroke: quantitative evaluation of pharyngeal transit times. *Arch Phys Med Rehabil* 1992;73:419-23.
- 36 Perlman AL, Booth BM, Grayhack JP. Videofluoroscopic predictors of aspiration in patients with oropharyngeal dysphagia. *Dysphagia* 1994;9:90-5.
- 37 Garon BR, Engle M, Orniston C. Silent aspiration: results of 1000 videofluoroscopic swallow evaluation. *Journal of Neurologic Rehabilitation* 1996;10:121-6.
- 38 Linden P. Videofluoroscopy in the rehabilitation of swallowing dysfunction. *Dysphagia* 1989;3:189-91.
- 39 Logemann JA. Non-invasive approaches to deglutitive aspiration. *Dysphagia* 1993;8:331-3.