SHORT REPORT

Reversible corpus callosum lesion in legionnaires' disease

J C Morgan, R Cavaliere, V C Juel

J Neurol Neurosurg Psychiatry 2004;75:651–654. doi: 10.1136/jnnp.2003.015669

Legionnaires' disease is often associated with neurological findings. Despite such findings, computed tomography and neuropathological investigations are typically normal. This report describes a reversible lesion of the corpus callosum identified on magnetic resonance imaging (MRI) in a patient with legionnaires' disease. MRI may show previously undocumented neuropathology in acute legionnaires' disease. *Legionella pneumophila* infection should be included in the differential diagnosis of conditions associated with reversible lesions of the corpus callosum.

Forty to fifty per cent of patients with legionnaires' disease develop neurological signs and symptoms.^{1,2} Despite frequent neurological findings in patients with this condition, neuroimaging^{1,3} and neuropathological³ studies are typically normal. We report a patient who developed neurological signs and symptoms in the setting of acute legionnaires' disease. Brain magnetic resonance imaging (MRI) showed a transient lesion in the splenium of the corpus callosum (SCC) that resolved in parallel with clinical improvement.

CASE REPORT

A middle aged man suffered three days of fever, malaise, and generalised weakness. He was subsequently admitted to the hospital after he was found unable to stand without assistance. His past medical history was unremarkable and he was not on any drug treatment. He drank two to three cans of beer a week and used no illicit drugs. On initial examination he was febrile, with moderate respiratory distress and left lower lobe rales. He was awake, oriented, and dysarthric and he had word finding difficulties. Extraocular movements were normal. He had bilateral intention tremor and dysmetria on finger–nose testing.

Laboratory abnormalities included: sodium 133 mmol/l (normal, 135–145), creatinine 141.4 µmol/l (61.9–114.9), glucose 16.6 mmol/l (3.9–5.8), white blood cell count 12 300/µl (4000–11 000/µl), platelet count 87 000/µl (150 000–450 000/µl), alanine aminotransferase 126 U/l (7–52), aspartate aminotransferase 170 U/l (13–39), total bilirubin 70.1 µmol/l (5.1–17.1), conjugated bilirubin 30.8 µmol/l (0.0–5.1), prothrombin time 16.8 s (12.5–14.6), international normalised ratio 1.3, and partial thromboplastin time 41.5 s (26.2–37.5). His blood ammonia level was normal at 27 µmol/l (9–33), and his blood ethanol level was <10 mg/dl (repeated). Vitamin B-12, folate, thyroid stimulating hormone, hepatitis A, B, C testing, HIV 1 and 2, rapid plasma reagin, antinuclear antibody, cytomegalovirus antibody, and rickettsial antibody screens were either normal or negative. A chest x ray showed left lower lobe consolidation, and an abdominal ultrasound was unremarkable. Computed tomography (CT) of the head without contrast was normal. Cerebrospinal fluid (CSF) analysis revealed normal cell counts (WBC 1/µl and RBC 1/µl), raised glucose at

6.7 µmol/l (2.2–3.9), and protein of 17 mg/dl (12–40). CSF Gram stain and cultures were negative.

He was treated with intravenous thiamine and began treatment with intravenous ceftriaxone, azithromycin, and doxycycline for community acquired pneumonia or possible rickettsial infection. A brain MRI done two days after admission showed hypointensity and slight oedema in the SCC on T1 weighted images (fig 1A), which did not enhance with contrast (fig 1D). There was markedly increased signal in this region on diffusion weighted imaging (DWI) (fig 1B). The lesion was slightly hyperintense on T2 weighted images (fig 1C). Intracranial magnetic resonance angiography showed no abnormalities (data not shown). A transthoracic echocardiogram revealed normal left ventricular function with no evidence of masses or thrombi.

After admission the patient became more somnolent and inattentive. In this setting, detailed features of a callosal syndrome⁴ could not be demonstrated conclusively. He was intermittently oriented with prolonged response latency, and he had difficulty in following complex commands. He could not perform sequential tasks and had limb apraxias. Although naming and repetition were normal, his speech remained dysarthric. Glabellar, snout, and palmo-mental reflexes were present. Finger–nose testing was accurate bilaterally, but his gait was ataxic. Electroencephalography showed generalised rhythmic slowing without epileptiform discharges, consistent with mild encephalopathy. A sputum sample was positive by direct fluorescent antibody (DFA) for *Legionella pneumophila*, and *Legionella pneumophila* serogroup 1 antigens were present in the urine. In the light of these findings, specific testing to exclude coinfection with *Mycoplasma* was not undertaken. The antibiotic regimen was changed to intravenous ciprofloxacin with subsequent gradual improvement in mentation. A repeat brain MRI with contrast done 13 days after the initial study showed complete resolution of the SCC lesion (fig 1, panels E–H).

He was discharged one month later with normal blood glucose, liver function, and renal function. Despite mild residual dysarthria, he could follow complex commands with less left–right confusion. Although his gait ataxia persisted, he was able to move around independently with a walking frame.

DISCUSSION

Legionnaires' disease is commonly associated with confusion, dysarthria, and cerebellar signs of gait and limb ataxia.^{1,2,5} Our patient had cerebellar dysfunction and frontal release signs, indicating neurological dysfunction beyond areas of the brain which typically contribute fibres to the SCC: the temporo-parieto-occipital junction, the superior parietal lobule, and the occipital cortex.⁶ Despite frequent cerebellar signs in legionnaires' patients, CT and brain scintigraphy have not revealed cerebellar lesions in acute disease.

Occasionally, neuroimaging studies have shown abnormalities associated with various clinical syndromes in patients with neurological deficits and *Legionella* infections (table 1).^{1,2,7–13} Identified lesions vary from abscess⁷ to

Figure 1 Brain magnetic resonance imaging (MRI) showing a callosal lesion on admission (A–D) and repeat brain MRI 13 days later (E–H). (A) Mid-sagittal T1 weighted image showing a hypointense lesion in the splenium of the corpus callosum. (B) Axial diffusion weighted imaging showing symmetrical hyperintensity in the same region. (C) Axial T2 weighted hyperintensity in the same region. (D) Postcontrast axial T1 weighted image showing hypointensity of the lesion without enhancement. (E–H) Corresponding images of the same patient 13 days later, with resolution of the callosal lesion.

Table 1 Brain imaging abnormalities in patients with neurological dysfunction and *Legionella* infections

Reference	Age/sex	Clinical symptoms, signs, outcome	Imaging results
Weir (1982) ¹	53/F	Headache, confusion, nuchal rigidity, dysarthria, extensor left toe, ataxia of lower extremities; 3 years later she had ataxia in her lower extremities more than in her upper extremities	Head CT, normal in acute setting; head CT 3 years later, cerebellar atrophy
Andersen (1987) ⁷	33/M	Headache, seizure, confusion, expressive aphasia, right sided hyperreflexia, dysmetria in right upper extremity; <i>Legionella jordanis</i> serum antibodies present at high titres; "normal" 7 months later except for "slightly impaired memory"	Head CT, left temporoparietal abscess, autologous leucocyte scan with enhancement in same region
Potasman (1990) ⁸	26/F	Fevers, headache, generalised seizure followed by confusion, left hemiparesis, additional generalised seizures; <i>Legionella bozemanii</i> antibodies in CSF; discharged 1 month later with normal neurological examination	Head CT, effacement of sulci
Johnson (1984) ²	Various, 7 patients with imaging (2 abnormal)	Patient with cerebral oedema on head CT had confusion leading to coma with otherwise normal neurological examination; patient survived. Patient with multifocal lesions on brain scan presented in coma and then developed left facial weakness, left hemiparesis, hyperreflexia, and an extensor left toe by day 12 of hospital stay; patient died and had necrotising haemorrhagic leucoencephalitis with bacilli seen on Dieterle stain at necropsy	Head CT, normal in 4 patients; brain scintigraphy, normal in 1 patient; head CT, cerebral oedema in one patient; brain scintigraphy, multifocal lesions in one patient
Karim (2002) ¹⁰	21/M	Unresponsive to verbal commands, with nuchal rigidity on admission, seizures followed, negative herpes simplex virus PCR; <i>Legionella pneumophila</i> by DFA after bronchoalveolar lavage; "the patient's neurologic condition gradually improved"	Head CT, normal initially, slight leptomeningeal enhancement on repeated study; brain MRI, bilateral mesiotemporal FLAIR hyperintensities
Sommer (2000) ¹²	58/M	Three weeks after a probable <i>Legionella pneumophila</i> illness (non-productive cough, fever, diarrhoea, abdominal pain, headache) the patient developed headache, nausea, dizziness, left-beating nystagmus and bilateral horizontal diplopia, lost consciousness; complete recovery 4 weeks later with steroid treatment	Brain MRI, confluent hyperintensity of the bilateral periventricular and subcortical white matter and left cerebellar peduncle (ADEM)
Spieker (1998) ¹¹	35/F	Six weeks after <i>Legionella cincinnatiensis</i> CNS infection (confirmed by PCR detection) the patient developed headache, agitation, hallucinations, paranoia, generalised seizures, mutism, somnolence, right facial dyskinesias, rigidity and dystonia; amnesia for 2 months, other abnormalities on the neurological examination normalised	Brain MRI, bilateral symmetrical hyperintensity of the basal ganglia and left subcortical white matter (ADEM); improved on repeat imaging
Platzek (1990) ⁹	44/M	<i>Legionella bozemanii</i> pneumonia followed by tetraparesis and severe midbrain syndrome; almost complete neurological recovery after several weeks	Head CT, normal; brain MRI, symmetrical, bilateral foci of demyelination in the brain stem
Easterbrook (1992) ¹³	19/M	Patient presented in coma with fever, CSF leucocytosis, <i>Legionella pneumophila</i> and <i>Mycoplasma pneumoniae</i> coinfection documented serologically; developed quadriplegia, anarthria, and dysphagia with little improvement	Head CT, normal on admission; brain MRI, 6 weeks later, multifocal cerebral white matter disease

ADEM, acute disseminated encephalomyelitis; CSF, cerebrospinal fluid; CT, computed tomography; FLAIR, fluid attenuated inversion recovery; MRI, magnetic resonance imaging; PCR, polymerase chain reaction.

cerebral oedema²⁻⁸ on CT, while brain scintigraphy revealed multifocal lesions in one patient.² More recently, brain MRI studies in two patients have shown bilateral symmetrical foci of brain stem demyelination in *Legionella bozemanii* pneumonia⁹ and bilateral mesial temporal FLAIR MRI hyperintensities following seizures in acute legionnaires' disease.¹⁰ Two patients developed acute disseminated encephalomyelitis (ADEM) several weeks after infection with *Legionella* species.¹¹⁻¹² Postinfectious encephalomyelitis has also occurred in a 19 year old man who had documented coinfection with *Legionella pneumophila* and *Mycoplasma pneumoniae*.¹³

The frequent finding of cerebellar signs in legionnaires' patients without neuroimaging evidence of cerebellar lesions may have several potential explanations. CT and brain scintigraphy may not be sensitive enough for routine detection of cerebellar lesions in legionnaires' patients. Additionally, patients developing an acute or postinfectious cerebellar ataxia often have normal cerebellar MRI findings, while functional imaging with single photon emission tomography (SPECT) may show marked diffuse cerebellar perfusion abnormalities.¹⁴

Our patient's SCC lesion may have generated more widespread cortical or cerebellar dysfunction through diaschisis or other mechanisms. Widespread bilateral alterations in cerebral blood flow and metabolism on positron emission tomography (PET) can occur in patients with Marchiafava Bignami disease who have a clinical presentation and neuroimaging findings similar to our patient.¹⁵ Similarly, in a series of 13 cases of intractable epilepsy undergoing total callosotomy, five developed persistent dysarthria and gait difficulties, while those receiving anterior two thirds or subtotal callosotomy did not develop these problems.¹⁶ This implies that posterior corpus callosum damage may be an important cause of dysarthria and gait ataxia.

In addition to Marchiafava Bignami disease, several other conditions with reversible SCC lesions are associated with clinical presentations similar to our patient. High altitude cerebral oedema (HACE), for example, is often associated with confusion, ataxia, and reversible SCC lesions.¹⁷ Reversible SCC lesions also occur in haemolytic-uraemic syndrome,¹⁸ rotavirus infection,¹⁹ and acute cerebellitis.²⁰

Hackett *et al* hypothesised¹⁷ that SCC lesions in HACE are related to vasogenic oedema because they respond to steroids and resolve slowly without significant residual clinical deficit. In haemolytic-uraemic syndrome, *E coli* O-157 verotoxin binding to cerebral vascular endothelium with subsequent microvascular angiopathy and perivascular oedema is proposed as a mechanism for a reversible SCC lesion.¹⁸ A reversible SCC lesion in a child with rotavirus infection was hypothesised to represent cytotoxic oedema, given that the lesion was hyperintense on DWI, though the pathogenic mechanism is unknown.¹⁹

Brain MRI in our patient showed DWI hyperintensity in the SCC that would customarily indicate cytotoxic oedema. Our patient did not receive corticosteroids, given the usual full neurological recovery in legionnaires' patients and the unproven benefit of corticosteroids in treating cytotoxic oedema. The resolution of his SCC lesion by MRI and the parallel clinical improvement is similar to the resolution of DWI changes in patients who develop acute metabolic decompensation and encephalopathy in maple syrup urine disease.²¹ When symptomatic, patients with this condition develop reversible regions of significantly restricted proton diffusion in large areas of white matter without evidence of volume loss or tissue damage. This may represent intramyelinic sheath oedema that reverses after metabolic correction.²¹ Perhaps resolution of intramyelinic sheath oedema explains the reversibility of our patient's SCC lesion and the absence of signs of infarction or necrosis.

The pathogenic mechanism of *Legionella* induced neurological dysfunction is unknown. In very rare instances, the Gram negative bacterium has been demonstrated directly in postmortem brain tissue²; however, this is usually not the case.³ An endotoxin effect accounting for the neurological dysfunction has been hypothesised,^{2-3,22} similar to the mechanism proposed for haemolytic-uraemic syndrome.¹⁸ As most patients with *Legionella* infections and CNS dysfunction have normal CSF leucocyte counts, with few exceptions,^{13,23} an endotoxic effect of *Legionella* spp would be a plausible mechanism for CNS disease. This endotoxin hypothesis is supported by evidence that *Legionella pneumophila* produces a cytotoxin that kills Chinese hamster ovary cells.²⁴ *Legionella* spp can also generate pores in eukaryotic cell membranes, leading to cell death by osmotic lysis.²⁵ Such features of *Legionella pneumophila* may have been responsible for our patient's reversible SCC lesion, though the local vulnerability of the SCC remains unexplained.

Application of newer imaging methods such as MRI may show that brain lesions in legionnaires' disease are more common than previously appreciated. Functional imaging with SPECT and PET may also provide evidence of CNS dysfunction in legionnaires' patients who show neurological signs without structural lesions on MRI. Resolution of oedematous brain lesions may account for the paucity of findings with less sensitive neuroimaging studies (CT and brain scintigraphy), the rarity of neuropathological findings, and the customarily complete neurological recovery in legionnaires' patients. Legionnaires' disease—like haemolytic-uraemic syndrome, rotavirus encephalopathy, and acute cerebellitis—should be included in the differential diagnosis of infectious conditions associated with reversible lesions of the corpus callosum.

ACKNOWLEDGEMENTS

We would like to thank Drs T P Bleck and G F Wooten for helpful criticism of our manuscript.

Authors' affiliations

J C Morgan, Medical College of Georgia, Department of Neurology, Augusta, Georgia, USA

R Cavaliere, V C Juel, University of Virginia, Department of Neurology, Charlottesville, Virginia, USA

Competing interests: none declared

Correspondence to: Dr Vern C Juel, PO Box 800394, Charlottesville, VA 22908-0394, USA; vj2n@virginia.edu

Received 28 March 2003

In revised form 19 August 2003

Accepted 24 August 2003

REFERENCES

- Weir AI, Bone I, Kennedy DH. Neurological involvement in legionellosis. *J Neurol Neurosurg Psychiatry* 1982;**45**:603-8.
- Johnson JD, Raff MJ, Van Arsdaal JA. Neurologic manifestations of Legionnaires' disease. *Medicine* 1984;**63**:303-10.
- Pendlebury WW, Perl DP, Winn WC, *et al*. Neuropathologic evaluation of 40 confirmed cases of Legionella pneumonia. *Neurology* 1983;**33**:1340-4.
- Habib M. Corpus callosum disconnection syndromes and functional organization of the corpus callosum in adults. *Neurochirurgia* 1998;**44**(suppl 1):102-9.
- Shetty KR, Cilyo CL, Starr BD, *et al*. Legionnaires' disease with profound cerebellar involvement. *Arch Neurol* 1980;**37**:379-80.
- de Lacoste MC, Kirkpatrick JB, Ross ED. Topography of the human corpus callosum. *J Neuropathol Exp Neurol* 1985;**44**:578-91.
- Andersen BB, Sogaard I. Legionnaires' disease and brain abscess. *Neurology* 1987;**37**:333-4.
- Potzman I, Liberson A, Krimerman S. Legionella infection mimicking herpes encephalitis. *Crit Care Med* 1990;**18**:453-4.
- Platzack C, Foerster EC, Schneider MU, *et al*. Encephalitis in Legionella bozemanii pneumonia. *Dtsch Med Wochenschr* 1990;**115**:1956-9.
- Karim A, Ahmed S, Rossoff LJ. Legionnaire's disease associated with acute encephalitis and arrhythmia. *Crit Care Med* 2002;**30**:1028-9.

- 11 **Spieker S**, Petersen D, Rolfs A, *et al.* Acute disseminated encephalomyelitis following Pontiac fever. *Eur Neurol* 1998;**40**:169–72.
- 12 **Sommer JB**, Erbguth FJ, Neundorfer B. Acute disseminated encephalomyelitis following *Legionella pneumophila* infection. *Eur Neurol* 2000;**44**:182–4.
- 13 **Easterbrook PJ**, Smyth EG. Post-infectious encephalomyelitis associated with *Mycoplasma pneumoniae* and *Legionella pneumophila* infection. *Postgrad Med J* 1992;**68**:124–8.
- 14 **Daaboul Y**, Vern BA, Blend MJ. Brain SPECT imaging and treatment with IVIg in acute post-infectious cerebellar ataxia: case report. *Neurol Res* 1998;**20**:85–8.
- 15 **Sano M**, Ishii K, Suzuki T, *et al.* Reductions in the bilateral parietal and occipital cerebral blood flow and metabolism in a patient with Marchiafava-Bignami disease. *J Neurol* 1999;**246**:607–8.
- 16 **Cendes F**, Ragazzo PC, da Costa V, *et al.* Corpus callosotomy in treatment of medically resistant epilepsy: preliminary results in a pediatric population. *Epilepsia* 1993;**34**:910–17.
- 17 **Hackett PH**, Yarnell PR, Hill R, *et al.* High-altitude cerebral edema evaluated with magnetic resonance imaging. *JAMA* 1998;**280**:1920–5.
- 18 **Ogura H**, Takaoka M, Kishi M, *et al.* Reversible MR findings of hemolytic uremic syndrome with mild encephalopathy. *Am J Neuroradiol* 1998;**19**:1144–5.
- 19 **Kobata R**, Tsukahara H, Nakai A, *et al.* Transient MR signal changes in the splenium of the corpus callosum in rotavirus encephalopathy: value of diffusion-weighted imaging. *J Comput Assist Tomogr* 2002;**26**:825–8.
- 20 **Kato Z**, Kozawa R, Hashimoto K, *et al.* Transient lesion in the splenium of the corpus callosum in acute cerebellitis. *J Child Neurol* 2003;**18**:291–2.
- 21 **Jan W**, Zimmerman RA, Wang ZJ, *et al.* MR diffusion imaging and MR spectroscopy of maple syrup urine disease during acute metabolic decompensation. *Neuroradiology* 2003;**45**:393–9.
- 22 **Wong KH**, Moss CW, Hochstein DH, *et al.* "Endotoxicity" of the Legionnaires' disease bacterium. *Ann Intern Med* 1979;**90**:624–7.
- 23 **Harris L**. Legionnaires' disease associated with acute encephalomyelitis. *Arch Neurol* 1981;**38**:462–3.
- 24 **Friedman RL**, Iglewski BH, Miller RD. Identification of a cytotoxin produced by *Legionella pneumophila*. *Infect Immun* 1980;**29**:271–4.
- 25 **Kirby JE**, Vogel JP, Andrews HL, *et al.* Evidence for pore-forming ability by *Legionella pneumophila*. *Mol Microbiol* 1998;**27**:323–36.