OPEN ACCESS

REVIEW

Short-term outcome of endoscopic versus microscopic pituitary adenoma surgery: a systematic review and meta-analysis

Mario Ammirati,¹ Lai Wei,² Ivan Ciric³

¹Department of Neurosurgery, Ohio State University, Columbus, Ohio, USA

²Center for Biostatistics, Ohio State University, Columbus, Ohio, USA

³Department of Neurosurgery, North Shore University Health System, Evanston, IL, USA

Correspondence to

Professor Mario Ammirati, Department of Neurosurgery, Ohio State University, 410 West 10th Avenue, N1025 Doan Hall, Columbus, OH 43210, USA; mario.ammirati@osumc.edu

Received 17 May 2012

Revised 18 September 2012

Accepted 13 November 2012

Published Online First

15 December 2012

Open Access
Scan to access more
free content

► <http://dx.doi.org/10.1136/jnnp-2012-304541>

► <http://dx.doi.org/10.1136/jnnp-2012-304583>

► <http://dx.doi.org/10.1136/jnnp-2013-305130>

To cite: Ammirati M, Wei L, Ciric I. *J Neurol Neurosurg Psychiatry* 2013;**84**:843–849.

ABSTRACT

Endoscopic transsphenoidal pituitary surgery has become increasingly more popular for the removal of pituitary adenomas. It is also widely recognised that transsphenoidal microscopic removal of pituitary adenomas is a well-established procedure with good outcomes. Our objective was to meta-analyse the short-term results of endoscopic and microscopic pituitary adenoma surgery. We undertook a systematic review of the English literature on results of transsphenoidal surgery, both microscopic and endoscopic from 1990 to 2011. Series with less than 10 patients were excluded. Pooled data were analysed using meta-analysis techniques to obtain estimate of death, complication rates and extent of tumour removal. Complications evaluated included cerebrospinal fluid leak, meningitis, vascular complications, visual complications, diabetes insipidus, hypopituitarism and cranial nerve injury. Data were also analysed for tumour size and sex. 38 studies met the inclusion criteria yielding 24 endoscopic and 22 microscopic datasets (eight studies included both endoscopic and microscopic series). Meta-analysis of the available literature showed that the endoscopic transsphenoidal technique was associated with a higher incidence of vascular complications ($p < 0.0001$). No difference was found between the two techniques in all other variables examined. Meta-analysis of the available literature reveals that endoscopic removal of pituitary adenoma, in the short term, does not seem to confer any advantages over the microscopic technique and the incidence of reported vascular complications was higher with endoscopic than with microscopic removal of pituitary adenomas. While we recognise the limitations of meta-analysis, our study suggests that a multicentre, randomised, comparative effectiveness study of the microscopic and endoscopic transsphenoidal techniques may be a reasonable approach towards establishing a true valuation of these techniques.

INTRODUCTION

The most widely used approach to remove pituitary adenomas since the 1960s has been the transsphenoidal route, executed using the operating microscope. This approach is generally regarded as being associated with good outcomes.¹ However, starting approximately 15 years ago and more so in the past 5 years, there has been a trend towards using endoscopic transsphenoidal techniques for the removal of pituitary adenomas.^{2–5} Endoscopic techniques have been recommended reportedly because of their lesser invasiveness, fewer complications and

overall better results compared to microscopic microsurgical techniques.⁵ In the absence of significant randomised clinical trials on the subject, a systematic review of published results of microsurgical and endoscopic techniques in the treatment of pituitary adenomas may be appropriate in comparing the two techniques.

METHODS**Search strategy**

The MEDLINE (1990 to present) and the EMBASE (1990 to present) databases were the information source used. The following keywords were searched individually or in association: pituitary, adenoma, sella, transsphenoidal, endoscopy, microscopic, Cushing's disease, prolactinoma, acromegaly, endonasal, minimally invasive surgery, complication, skull base and surgery. The included limits were: English for the language category and humans for the study category. The date of the last search was August 2011.

Study selection criteria

Series with 10 or more patients with pituitary adenoma, operated on using either microscopic or endoscopic techniques were included. Manuscripts in which the approach was guided by the endoscope and tumour removal was carried out using the microscope were included in the microscope group. As the intent of this report was to analyse the results of papers in which transsphenoidal surgery (either endoscopic or microscopic) was deemed to be the best modality to treat pituitary adenoma, we excluded reports in which a decision had been made preoperatively to use a combined transsphenoidal and intracranial approach.⁶ Series including mixed pathology and in which it was not possible to extrapolate the adenoma results were also excluded.^{7–11}

Mined data

Mined data include year of publication, study design, number and sex of subjects, size of the adenoma (micro vs macro), death, cerebrospinal fluid leak, meningitis, vascular complications, visual complications, diabetes insipidus transient and permanent, hypopituitarism, cranial nerve injury and complete versus partial resection. Vascular complications included carotid or other vessels injury, intracerebral haematoma, or any symptomatic intratumoral or intrasellar haemorrhage. Venous bleeding from the cavernous sinus was tabulated as a vascular complication when it prevented the completion of the surgical

procedure. Epistaxis was not included among the vascular complications. Cerebrospinal fluid leak included all postoperative leaks and all patients who developed an intra-operative leak requiring lumbar drainage. Decreased postoperative anterior pituitary function did not include Cushing's patients receiving postoperative cortisol or patients in whom a hypophysectomy was carried out.

Statistical analysis

There were 38 studies that fulfilled the inclusion criteria.^{1 3 12–47} All of them except one that was prospective and randomised²⁷ were retrospective studies. Only eight out of these 38 studies reported data on complications from both the endoscopic and microscopic surgery groups.^{15 17 18 21 27 32 36 45} Only one of these eight studies was prospective and randomised, with only 10 patients in each arm,²⁷ the others being retrospective studies. In six out of these seven studies the endoscopic and microscopic patients were accrued along longitudinally different time frames.^{17 18 21 32 36 45} Furthermore, in two of these seven studies tumour resection was either not evaluated at all,¹⁵ or was judged by the surgeon at the time of the operation.⁴⁵ In addition, not all of these studies reported data on other relevant metrics (ie, tumour size, vascular and or visual complications). Due to the overall limitations of these studies we felt that we could have got more useful information by analysing all the data reported in the literature that fulfilled our inclusion criteria. Given the nature of the studies available in the literature it was not possible to compare the results (eg, complication rates) of endoscopic surgery to microscopic surgery using the traditional meta-analysis method with relative risk or OR. Instead, weighted summary rates and 95% CI of death and major complications/events (cerebrospinal fluid leak, meningitis, vascular complications, visual loss, diabetes insipidus temporary or permanent, hypopituitarism, complete resection, and nerve injury) were provided using the appropriate meta-analysis models (ie, random effects or fixed effects) for endoscopic and microscopic surgery, respectively. Then we compared the outcomes between endoscopic and microscopic surgery using the estimated proportions and SE (ie, 95% CI). Freeman–Tukey double arcsine transformation is used for proportions.⁴⁸ Tests for heterogeneity were performed for each meta-analysis. Based on results of the heterogeneity test, random-effects (heterogeneity present) or fixed-effects (heterogeneity absent) models were chosen to estimate the results.⁴⁹ Using this methodology, each study was weighted more fairly than giving more weight to those studies with larger sample sizes, especially when the interstudy variation was larger than the intrastudy variation. Tabaei *et al*⁵ used this methodological approach in 2009 when reviewing endoscopic pituitary surgery. The meta-analysis was performed using 'meta' package from the R statistical software.⁵⁰

RESULTS

Sixteen manuscripts dealt with transsphenoidal endoscopic removal of pituitary tumours,^{3 13 14 16 20 22 23 25 29 34 35 38 40 43 44 47} and 14 manuscripts reported on transsphenoidal removal of pituitary tumours using the operating microscope.^{1 12 19 24 26 28 30 31 33 37 39 41 42 46} Eight manuscripts included both techniques.^{15 17 18 21 27 32 36 45} Study type and demographics of all the studies are presented in table 1.

Estimated proportions with 95% CI of macro tumour size and women were obtained using meta-analysis techniques.

We failed to detect any statistically significant difference in the two groups regarding the proportion of macro tumour size (endoscopic: 79.75% with 95% CI of 73.20% to 85.58%; microscopic: 66.90% with 95% CI of 47.54% to 83.67%) and

women (endoscopic: 54.49% with 95% CI of 51.44% to 57.52%; microscopic: 54.51% with 95% CI of 47.58% to 61.35%).

We also used meta-analysis techniques to estimate the death and complications rates. Based on the test of heterogeneity results, random-effects pooled or fixed-effects pooled rates were estimated for death, cerebrospinal fluid leak, meningitis, vascular complications, visual complication, diabetes insipidus temporary, diabetes insipidus permanent, hypopituitarism, complete resection and nerve injury with endoscopic and microscopic surgery, respectively (table 2). Fixed-effects pooled estimates of the death rate with endoscopic and microscopic surgery were computed because there was no evidence of heterogeneity ($p=0.84$ and 1). The pooled death rates were 0.49% (95% CI 0.23% to 0.84%) and 0.23% (95% CI 0.10% to 0.42%) for endoscopic and microscopic surgery, respectively. The fixed-effects pooled meningitis rate ($p=0.86$, test for heterogeneity) for endoscopic surgery and the random-effects pooled meningitis rate ($p<0.0001$, test for heterogeneity) for microscopic surgery were 1.11% (95% CI 0.64% to 1.71%) and 2.08% (95% CI 0.83% to 3.86%). The fixed-effects pooled estimates of vascular complications were calculated for endoscopic and microscopic surgery ($p=0.87$ and 0.39 , test for heterogeneity), which were 1.58% (95% CI 1.07% to 2.19%, figure 1) and 0.50% (95% CI 0.28% to 0.78%, figure 2). The fixed-effects or random-effects pooled estimates of visual complication rate ($p=0.50$ and 0.04 , heterogeneity) and nerve injury rate ($p=0.48$ and 0.01 , heterogeneity) were calculated for endoscopic and microscopic surgery, respectively, and are listed in table 2.

The random-effects pooled estimates with 95% CI of cerebrospinal fluid leak rate (both $p<0.0001$, test for heterogeneity), temporary diabetes insipidus rate (both $p<0.0001$, heterogeneity), permanent diabetes insipidus rate ($p=0.02$ and <0.0001 , heterogeneity) and hypopituitarism rate (both $p<0.0001$, heterogeneity) for endoscopic and microscopic surgery were also calculated and listed in table 2. We failed to demonstrate any statistically significant difference in the pooled rate of death, cerebrospinal fluid leak, meningitis, visual complication, diabetes insipidus temporary, diabetes insipidus permanent, hypopituitarism, complete resection and nerve injury between endoscopic and microscopic surgery using the estimated proportions and SE. On the other hand, the vascular complication rate in endoscopic surgery is higher than in microscopic surgery (figures 1 and 2, $p<0.0001$) from those studies reporting the vascular complication information. Furthermore, the macro tumours or the extent of complete tumour resection, two variables that could have affected the vascular complication rate, were found not to be statistically significant in the endoscopic or microscopic series reporting vascular complications ($p=0.24$ and 0.10 for macro tumour size and complete resection, respectively).

DISCUSSION

Transsphenoidal microscopic pituitary surgery

Transsphenoidal microscopic pituitary surgery is a time confirmed effective way of treating pituitary adenomas, as shown by the low incidence of complications in any clinical/practice setting according to the Ciric-sponsored survey of the late 1990s,¹ although a statistically significant difference was noted in terms of better results in the hands of more experienced surgeons. Microscopic pituitary surgery relies on three-dimensional visualisation and thus on the ability to operate in three-dimensional space. Important conceptual nuances that need to be implemented in its execution to lead to optimal results are the recognition that the pituitary gland is a midline, extra-

Table 1 Study types and demographics

Authors	Year	Duration of study	Operation type	No of patients	Mean age, years (range)	Women (%)	Macroadenomas (%)
Jho and Carrau ³	1997	1993–6	Endoscopic	44	38 (14–88)	19 (43)	31 (70)
Cappabianca <i>et al</i> ¹⁴	2002	1997–2001	Endoscopic	146	46.1 (16–74)	76 (52)	125 (86)
White <i>et al</i> ⁴⁵	2004	2000–2	Endoscopic	50	41.1 (NA)	26 (52)	NA
Senior <i>et al</i> ⁴⁰	2008	NA	Endoscopic	148	46 (NA)	81 (55)	114 (81)†
Netea-Meier <i>et al</i> ³⁵	2006	1998–2004	Endoscopic	35	41 (14–68)	25 (71)	6 (17)
Rudnik <i>et al</i> ³⁸	2005	2001–2	Endoscopic	63	48.3 (11–77)	34 (54)	52 (83)
Kenan <i>et al</i> ²⁹	2006	1997–2005	Endoscopic	78	44.7 (11–67)	NA	67 (86)
Jain <i>et al</i> ²⁷	2007	NA	Endoscopic	10	40.1 (NA)	6 (60)	9 (90)
Uren <i>et al</i> ⁴³	2007	2001–5	Endoscopic	32	53 (25–69)	12 (38)	23 (72)
Frank <i>et al</i> ²²	2006	1998–2004	Endoscopic	381	48 (7–89)	227 (60)	284 (75)
Dehdashti <i>et al</i> ²⁰	2008	2004–7	Endoscopic	200	49.9 (20–78)	109 (55)	158 (79)
Charalampaki <i>et al</i> ¹⁶	2007	2004–5	Endoscopic	50	56 (28–84)	30 (60)	41 (82)
Yano <i>et al</i> ⁴⁴	2009	2001–8	Endoscopic	194	53 (15–85)*	120 (62)	NA
Zhang <i>et al</i> ⁴⁷	2008	1998–2005	Endoscopic	78	45.1 (15–76)	42 (54)	67 (86)
Campbell <i>et al</i> ¹³	2010	2005–9	Endoscopic	26	45.7 (20–69)	12 (46)	22
Gondim <i>et al</i> ²³	2010	1998–2009	Endoscopic	301	42 (12–79)	167 (55)	248
Hofstetter <i>et al</i> ²⁵	2010	2004–10	Endoscopic	24	50.7 (22–75)	11 (46)	19
Nakao <i>et al</i> ²⁴	2011	2000–8	Endoscopic	43	55 (31–75)	20 (46)	43
Casler <i>et al</i> ¹⁵	2005	1996–2003	Endoscopic	15	41.6	9 (60)	NA
Duz <i>et al</i> ²¹	2008	1996–2007	Endoscopic	28	NA	NA	NA
Choe <i>et al</i> ¹⁷	2008	2004–7	Endoscopic	12	47±12	7 (58)	9
O'Malley <i>et al</i> ²⁶	2008	2003–8	Endoscopic	25	47.9 (18–73)	10 (40)	22
D'Haens <i>et al</i> ¹⁸	2009	2001–7	Endoscopic	60	37 (10–70)	41 (68)	31
Messerer <i>et al</i> ²²	2011	2006–9	Endoscopic	82	57 (20–82)	35 (43)	82
Jain <i>et al</i> ²⁷	2007	NA	Microscopic	10	31.6 (18–58)	5 (50)	8 (80)
Mortini <i>et al</i> ³³	2005	1990–2002	Microscopic	1140	43 (8–82)	681 (60)	788 (69)
Höybye <i>et al</i> ²⁶	2004	1990–9	Microscopic	34	40 (13–74)	26 (76)	2 (6)
White <i>et al</i> ⁴⁵	2004	1996–9	Microscopic	50	43.5 (NA)	17 (34)	NA
Ciric <i>et al</i> ¹	1997	NA	Microscopic	638	NA	NA	NA
De <i>et al</i> ¹⁹	2003	1980–2001	Microscopic	90	61 (29–86)	46 (51)	61 (68)
Rees <i>et al</i> ²⁷	2002	1980–2000	Microscopic	54	41.3 (14–73)	42 (78)	10 (19)
Shimon <i>et al</i> ⁴²	2002	1990–2000	Microscopic	74	39 (8–72)	64 (86)	3 (4)
Han <i>et al</i> ²⁴	2008	1996–2006	Microscopic	592	42.3 (NA)	274 (46)	592 (100)‡
Kaltsas <i>et al</i> ²⁸	2001	1993–8	Microscopic	67	46 (12–80)*	31 (46)	50 (75)
Kreutzer <i>et al</i> ³¹	2001	1992–8	Microscopic	57	43.9 (16–71)	37 (65)	38 (67)
Semple and Laws ³⁹	1999	1992–7	Microscopic	105	38.5 (6–78)	83 (79)	13 (12)
Zhang <i>et al</i> ⁴⁶	1999	1982–97	Microscopic	208	47.5 (16–71)	110 (53)	208 (100)
Koren <i>et al</i> ³⁰	1999	1993–5	Microscopic	37	NA	NA	23 (62)
Badie <i>et al</i> ¹²	2000	1996–9	Microscopic	34	45.5 (NA)	17 (50)	21 (62)
Sheehan <i>et al</i> ⁴¹	1999	1995–7	Microscopic	70	58.5 (25–85)	21 (30)	70 (100)
Casler <i>et al</i> ¹⁵	2005	1996–2003	Microscopic	15	50.6	10 (67)	NA
Choe <i>et al</i> ¹⁷	2008	1997–2004	Microscopic	11	48±10	9 (82)	8
O'Malley <i>et al</i> ²⁶	2008	2003–8	Microscopic	25	50.8 (23–78)	10 (40)	23
Duz <i>et al</i> ²¹	2008	1996–2007	Microscopic	65	NA	NA	NA
D'Haens <i>et al</i> ¹⁸	2009	2001–7	Microscopic	60	35 (10–68)	16 (27)	27
Messerer <i>et al</i> ²²	2011	2005	Microscopic	82	56.5 (27–84)	35 (43)	82

*Median age.

†Adenoma size was only available in 140 of 148 patients.

‡Including 438 macro and 154 giant.

arachnoidal structure, and that the normal anterior pituitary gland is distended by the macroadenoma into a thin layer of tissue surrounding the pathology, whose preservation maximises postoperative pituitary function.⁵¹ Therefore, microscopic pituitary surgery needs to be extra-arachnoidal, to respect the principle of midline and needs to try to identify and preserve normal pituitary tissue.

Endoscopic pituitary surgery

Endoscopic pituitary surgery has been advocated for the surgical treatment of pituitary adenomas since the late 1990s on the basis of its panoramic improved visualisation and thus the ability to visualise structures easily such as the optico-carotid recesses and to access suprasellar areas visually that are not in the direct line of vision during microscopic pituitary

Table 2 Estimated proportion with 95% CI of complications/events for endoscopy and microscopy

Complication/events	No of studies for endoscopy	Proportion (95% CI) for endoscopy	No of studies for microscopy	Proportion (95% CI) for microscopy
Death	19	0.49% (0.23% to 0.84%)†	18	0.23% (0.10% to 0.42%)†
Cerebrospinal fluid leak	24	7.00% (4.84% to 9.52%)	19	6.34% (3.86% to 9.37%)
Meningitis	13	1.11% (0.64% to 1.71%)†	14	2.08% (0.83% to 3.86%)
Vascular complications*	17	1.58% (1.07% to 2.19%)†	12	0.50% (0.28% to 0.78%)†
Visual loss	13	0.72% (0.37% to 1.19%)†	14	0.60% (0.23% to 1.14%)
Diabetes insipidus temporary	18	9.10% (6.57% to 11.99%)	14	10.23% (6.50% to 14.69%)
Diabetes insipidus permanent	21	2.31% (1.41% to 3.41%)	15	4.25% (1.96% to 7.36%)
Hypopituitarism	17	8.51% (5.16% to 12.59%)	12	11.64% (5.14% to 20.32%)
Complete resection	22	68.77% (64.37% to 73.00%)	18	64.44% (57.62% to 70.98%)
Nerve injury	8	0.28% (0.05% to 0.71%)†	7	0.53% (0.08% to 1.34%)

*Vascular complication rate is higher using the endoscope than using the microscope ($p < 0.0001$).

†A fixed-effects pooled estimate is used for the proportions. Otherwise, a random-effects pooled estimate is used.

surgery.^{5 20 22} This improved visualisation was felt to be theoretically associated with a greater extent of resection and improved safety, less invasiveness and potential better results—compared to microscopic pituitary surgery.⁵ However, endoscopic pituitary surgery is bidimensional and thus depth perception is missing, although manoeuvres have been recommended to minimise this drawback.²¹ The three surgical principles as described under microscopic surgery should also be applicable to endoscopic surgery.

Results of our study

Our study did not demonstrate any short-term advantages of endoscopic pituitary adenoma surgery. The review of the literature suggests an increased incidence of vascular complications associated with it. Recognising the limitations of a systematic review in coming to firm conclusions, it is conceivable that the greater incidence of vascular complications associated with the endoscopic transsphenoidal surgical technique is the consequence of either a more robust sphenoid exposure of the carotid optic recess with a greater potential for carotid artery injury, or the lack of three-dimensional depth perception during suprasellar manoeuvres. Indeed, the majority of vascular complications associated with the endoscopic technique involved

suprasellar vascular injuries including intracerebral haemorrhages, midbrain stroke, mesencephalic haemorrhage, posterior cerebral artery laceration and internal carotid artery injury.^{14 16 22 40}

Other possible advantages of endoscopic pituitary surgery

Other theoretical advantages of endoscopic pituitary surgery such as its limited invasiveness or minimally invasive feature, leading to an overall better quality of life, have not been addressed by our study.^{16 29 38 40 47} Indeed, the only study that looked at outcome and quality of life after endoscopic surgery for pituitary adenomas showed only 'a trend to improved scores in the endoscopic group compared with previous studies in patients treated by conventional approaches'.⁵² A recent study has reported a 29.6% incidence of extracranial complications following endoscopic transsphenoidal sellar surgery, with the majority of them being prolonged nasal crusting (10.8%) and synechiae formation (8.8%).⁵³ Again, without making a definitive statement, it appears plausible that these nasal complications are the consequence of liberal and repetitive passage of instrumentation through the nasal cavity. That is even more likely to occur whenever nasal septal flaps are developed. Postoperative nasal debridement is usually required.⁸

Figure 1 Proportion of vascular complications with endoscopy. Test for heterogeneity: $Q=10.01$ on 16 degrees of freedom ($p=0.87$).

Figure 2 Proportion of vascular complications with microscopy. Test for heterogeneity: $Q=11.67$ on 11 degrees of freedom ($p=0.39$).

Other systematic reviews

There are three systematic reviews comparing endoscopic to microscopic pituitary adenoma removal.^{54–56} One such review⁵⁵ is difficult to evaluate due to the fact that it includes, among other things, mixed patient/pathology cohorts in which endonasal microscopic surgery ‘...in some cases with endoscopic assistance’ was compared to a sublabial microscopic technique.⁵⁷ The other two reviews, one analysing 11⁵⁴ and the other 10 studies,⁵⁶ with the same 10 studies being evaluated by both, reached somewhat different conclusions. Goudakos *et al*⁵⁴ indicated that the occurrence of diabetes insipidus, rhinological complications and length of hospital stay was significantly less in the endoscopic than in the microscopic group, while rates of complete tumour resection, cerebrospinal fluid leak, operative time and blood loss were not different in the two groups. The other review⁵⁶ shows that the purely endoscopic approach was linked to less blood loss, shorter hospital stay, shorter operative times and fewer nasal complications, but a higher incidence of cerebrospinal fluid leak than the microscopic approach, while other outcomes and complication rates were the same in the two groups.

Our review analysed a total of 5643 patients, of which were 2125 operated using the endoscope and 3518 using the microscope; data on vascular complications were available in 1887 patients operated with the endoscope and 3023 patients operated using the microscope. This is in contrast to 812 patients (373 endoscopic and 430 microscopic),⁵⁴ 549 patients (236 endoscopic and 313 microscopic)⁵⁵ and 687 patients (314 endoscopic and 373 microscopic)⁵⁶ collated by the three other systematic reviews.

Perspective

The need for careful evaluation of surgical innovations led to the convening of the Balliol colloquium and the Balliol collaboration at Balliol College in Oxford, UK, between 2007 and 2009. Participants included surgeons, methodologists, statisticians and the evidence-based medicine community. The results of such an effort were published in 2009.⁵⁸ Suffice it to say that notwithstanding transformative innovations such as tracheotomy for tracheal obstruction, leading to ‘an advance that is clear and substantial and that cannot be explained by either chance or bias’ the majority of surgical innovations are incremental and are ‘prone to overoptimistic assessments by their developers and, therefore, need controlled randomised studies, when

possible’.⁵⁸ The colloquium proposed a progressive algorithm, ‘the IDEAL recommendations’, to evaluate surgical innovations.⁵⁸ Stage 1, the idea phase, should be prompted by the need for a new solution to a clinical problem. It is essentially the proof of a concept and is implemented in only a few patients with careful outcome—especially negative outcome—reporting.

Stage 2a, the development phase, involves the planned use of the procedure in a small group of selected patients to accrue outcome on the safety of the procedure and its technical and procedural success or lack thereof. Protocols should be registered beyond the surgeons’ institution to secure equipoise and should undergo some form of ethical approval and informed research consent. It is our belief that endoscopic transsphenoidal surgery has fulfilled splendidly the requirements under these two progressive algorithm stages.

Stage 2b, the exploration phase, involves the evaluation of the outcome of a rather well-defined procedure in a larger group of patients. Rigorous prospective trials looking at short-term outcomes (technical clinical and patient centered) need to be employed in this stage. Clearly, stringent quality controls need to be in place.

Stage 3, the assessment phase, involves well-defined indications, a stable procedure executed by many surgeons and the comparison of this procedure with the gold standard procedure. This is usually done using randomised clinical trials or alternative well-structured trials, such as parallel group non-randomised studies, controlled interrupted-time series studies, tracker trials or similar designs. Again, institutional review board approval and informed research consent are mandatory.

Stage 4, the long-term study phase, is a surveillance study wherein the now established procedure is evaluated for rare and long-term outcomes. No ethical approval is necessary.

Endoscopic pituitary surgery and the IDEAL framework

It appears that 12 years after the publication of one of the first peer-reviewed papers on the use of endoscopic pituitary surgery²⁶ the advantages of the procedure are still not fully consolidated in accordance with the above algorithm for the evaluation of new surgical techniques. Clearly, the introduction of new surgical technology/procedures, such as endoscopic transsphenoidal surgery, is both a laudatory step towards innovation as well as a complex process requiring rigorous scrutiny with evidence-based methods such as discussed above or a similar acceptable variant thereof.^{58–60}

CONCLUSIONS

Meta-analysis of available literature on endoscopic transsphenoidal removal of pituitary adenomas seems to suggest that the endoscopic technique is associated with a higher incidence of vascular complications compared to microscopic transsphenoidal removal of pituitary adenomas. Moreover, in the short term, the same analysis of the literature reveals that there is no clear, unequivocal advantage to endoscopic transsphenoidal removal compared to microscopic transsphenoidal removal of pituitary adenomas.

Contributors MA, LW and IC all contributed to the conception and design of the article, to the data acquisition, analysis and interpretation. They were all involved in critically drafting/revising the article for important intellectual content. Finally, they all gave final approval of the version to be published.

Funding None.

Competing interests None.

Provenance and peer review Not commissioned; externally peer reviewed.

Open Access This is an Open Access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 3.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/3.0/>

REFERENCES

- Ciric I, Ragin A, Baumgartner C, et al. Complications of transsphenoidal surgery: results of a national survey, review of the literature, and personal experience. *Neurosurgery* 1997;40:225–37.
- Cavallo LM, Dal Fabbro M, Jalalod'din H, et al. Endoscopic endonasal transsphenoidal surgery. Before scrubbing in: tips and tricks. *Surg Neurol* 2007;67:342–7.
- Jho HD, Carrau RL. Endoscopic endonasal transsphenoidal surgery: experience with 50 patients. *J Neurosurg* 1997;87:44–51.
- Powell M, Gnanalingham KK. Endoscopic trans-sphenoidal pituitary surgery: is it here to stay? *Br J Neurosurg* 2007;21:315–17.
- Tabaee A, Anand VK, Barron Y, et al. Endoscopic pituitary surgery: a systematic review and meta-analysis. *J Neurosurg* 2009;111:545–54.
- Kabil MS, Eby JB, Shahinian HK. Fully endoscopic endonasal vs. transseptal transsphenoidal pituitary surgery. *Minim Invasive Neurosurg* 2005;48:348–53.
- Graham SM, Iseli TA, Karnell LH, et al. Endoscopic approach for pituitary surgery improves rhinologic outcome. *Ann Otol Rhinol Laryngol* 2009;118:630–5.
- Higgins TS, Courtemanche C, Karakla D, et al. Analysis of transnasal endoscopic versus transseptal microscopic approach for excision of pituitary tumors. *Am J Rhinol* 2008;22:649–52.
- Jho HD. Endoscopic transsphenoidal surgery. *J Neurooncol* 2001;54:187–95.
- Neal JG, Patel SJ, Kulbersh JS, et al. Comparison of techniques for transsphenoidal pituitary surgery. *Am J Rhinol* 2007;21:203–6.
- Shah S, Har-El G. Diabetes insipidus after pituitary surgery: incidence after traditional versus endoscopic transsphenoidal approaches. *Am J Rhinol* 2001;15:377–9.
- Badie B, Nguyen P, Preston JK. Endoscopic-guided direct endonasal approach for pituitary surgery. *Surg Neurol* 2000;53:168–73.
- Campbell PG, Kenning E, Andrews DW, et al. Outcome after a purely endoscopic transsphenoidal resection of growth hormone-secreting pituitary adenomas. *Neurosurg Focus* 2010;29:E5.
- Cappabianca P, Cavallo LM, Colao A, et al. Surgical complications associated with the endoscopic endonasal transsphenoidal approach for pituitary adenomas. *J Neurosurg* 2002;97:293–8.
- Casler JD, Doolittle AM, Mair EA. Endoscopic surgery of the anterior skull base. *Laryngoscope* 2005;115:16–24.
- Charalampaki P, Reisch R, Ayad A, et al. Endoscopic endonasal pituitary surgery: surgical and outcome analysis of 50 cases. *J Clin Neurosci* 2007;14:410–15.
- Choe JH, Lee KS, Jeun SS, et al. Endocrine outcome of endoscopic endonasal transsphenoidal surgery in functioning pituitary adenomas. *J Korean Neurosurg Soc* 2008;44:151–5.
- D'Haens J, Van Rompaey K, Stadnik T, et al. Fully endoscopic transsphenoidal surgery for functioning pituitary adenomas. A retrospective comparison with traditional transsphenoidal microsurgery in the same institution. *Surg Neurol* 2009;72:336–40.
- De P, Rees DA, Davies N, et al. Transsphenoidal surgery for acromegaly in Wales: results based on stringent criteria of remission. *J Clin Endocrinol Metab* 2003;88:3567–72.
- Dehdashti AR, Ganna A, Karabatsou K, et al. Pure endoscopic endonasal approach for pituitary adenomas: early surgical results in 200 patients and comparison with previous microsurgical series. *Neurosurgery* 2008;62:1006–17.
- Duz B, Harman F, Secer HI, et al. Transsphenoidal approaches to the pituitary: a progression in experience in a single center. *Acta Neurochir* 2008;150:1133–39.
- Frank G, Pasquini E, Farneti G, et al. The endoscopic versus the traditional approach in pituitary surgery. *Neuroendocrinology* 2006;83:240–8.
- Gondim JA, Almeida JPC, Albuquerque LAF, et al. Endoscopic endonasal approach for pituitary adenoma: surgical complications in 301 patients. *Pituitary* 2011;14:174–83.
- Han ZL, He DS, Mao ZG, et al. Cerebrospinal fluid rhinorrhea following trans-sphenoidal pituitary macroadenoma surgery: experience from 592 patients. *Clin Neurol Neurosurg* 2008;110:570–9.
- Hofstetter CP, Manna RH, Mubita L, et al. Endoscopic endonasal transsphenoidal surgery for growth hormone-secreting pituitary adenomas. *Neurosurg Focus* 2010;29:E6.
- Hoybe C, Grenback E, Thoren M, et al. Transsphenoidal surgery in Cushing disease: 10 years of experience in 34 consecutive cases. *J Neurosurg* 2004;100:634–8.
- Jain AK, Gupta AK, Pathak A, et al. Excision of pituitary adenomas: randomized comparison of surgical modalities. *Br J Neurosurg* 2007;21:328–31.
- Kaltsas GA, Isidori AM, Florakis D, et al. Predictors of the outcome of surgical treatment in acromegaly and the value of the mean growth hormone day curve in assessing postoperative disease activity. *J Clin Endocrinol Metab* 2001;86:1645–52.
- Kenan K, Ihsan A, Dilek O, et al. The learning curve in endoscopic pituitary surgery and our experience. *Neurosurg Rev* 2006;29:298–305.
- Koren I, Hadar T, Rappaport ZH, et al. Endoscopic transnasal transsphenoidal microsurgery versus the sublabial approach for the treatment of pituitary tumors: endonasal complications. *Laryngoscope* 1999;109:1838–40.
- Kreutzer J, Vance ML, Lopes MB, et al. Surgical management of GH-secreting pituitary adenomas: an outcome study using modern remission criteria. *J Clin Endocrinol Metab* 2001;86:4072–77.
- Messerer M, De Battista JC, Raverot G, et al. Evidence of improved surgical outcome following endoscopy for nonfunctioning pituitary adenoma removal. Personal experience and review of the literature. *Neurosurg Focus* 2011;30:E11.
- Mortini P, Losa M, Barzaghi R, et al. Results of transsphenoidal surgery in a large series of patients with pituitary adenoma. *Neurosurgery* 2005;56:1222–33.
- Nakao N, Itakura T. Surgical outcome of the endoscopic endonasal approach for non-functioning giant pituitary adenoma. *J Clin Neurosci* 2011;18:71–5.
- Netea-Meier RT, van Lindert EJ, den Heijer M, et al. Transsphenoidal pituitary surgery via the endoscopic technique: results in 35 consecutive patients with Cushing's disease. *Eur J Endocrinol* 2006;154:675–84.
- O'Malley B, Grady SM, Gabel BJ, et al. Comparison of endoscopic and microscopic removal of pituitary adenomas: single-surgeon experience and the learning curve. *Neurosurg Focus* 2008;25:E10.
- Rees DA, Hanna FW, Davies JS, et al. Long-term follow-up results of transsphenoidal surgery for Cushing's disease in a single centre using strict criteria for remission. *Clin Endocrinol (Oxf)* 2002;56:541–51.
- Rudnik A, Zawadzki TR, Wojtacha M, et al. Endoscopic transnasal transsphenoidal treatment of pathology of the sellar region. *Minim Invasive Neurosurg* 2005;48:101–7.
- Semple PL, Laws ER Jr. Complications in a contemporary series of patients who underwent transsphenoidal surgery for Cushing's disease. *J Neurosurg* 1999;91:175–9.
- Senior BA, Ebert CS, Bednarski KK, et al. Minimally invasive pituitary surgery. *Laryngoscope* 2008;118:1842–55.
- Sheehan MT, Atkinson JL, Kasperbauer JL, et al. Preliminary comparison of the endoscopic transnasal vs the sublabial transseptal approach for clinically nonfunctioning pituitary macroadenomas. *Mayo Clin Proc* 1999;74:661–70.
- Shimon I, Ram Z, Cohen ZR, et al. Transsphenoidal surgery for Cushing's disease: endocrinological follow-up monitoring of 82 patients. *Neurosurgery* 2002;51:57–62.
- Uren B, Vrodo N, Wormald PJ. Fully endoscopic transsphenoidal resection of pituitary tumors: technique and results. *Am J Rhinol* 2007;21:510–14.
- Yano S, Kawano T, Kudo M, et al. Endoscopic endonasal transsphenoidal approach through the bilateral nostrils for pituitary adenomas. *Neurol Med Chir* 2009;49:1–7.
- White DR, Sonnenburg RE, Ewend MG, et al. Safety of minimally invasive pituitary surgery (MIPS) compared with a traditional approach. *Laryngoscope* 2004;114:1945–48.
- Zhang X, Fei Z, Zhang J, et al. Management of nonfunctioning pituitary adenomas with suprasellar extensions by transsphenoidal microsurgery. *Surg Neurol* 1999;52:380–5.
- Zhang X, Fei Z, Zhang W, et al. Endoscopic endonasal transsphenoidal surgery for invasive pituitary adenoma. *J Clin Neurosci* 2008;15:241–5.
- Miller JJ. The inverse of the Freeman-Tukey double arcsine transformation. *Am Statistician* 1978;32:138.
- Whitehead A. *Meta-Analysis of controlled clinical trials*. Chichester, UK: John Wiley & Sons Ltd, 2002:151–74.

- 50 Schwarzer G. Meta-analysis with R (reference manual of package 'meta' in R). The R foundation for statistical computing (Publisher), Vienna, Austria:2010:1–65.
- 51 Ciric I, Rosenblatt S, Baumgarter C, *et al.* Transsphenoidal microsurgery. *Neurosurgery* 2002;51:161–9.
- 52 Karabatsou K, O'Kelly C, Ganna A, *et al.* Outcomes and quality of life assessment in patients undergoing endoscopic surgery for pituitary adenomas. *Br J Neurosurg* 2008;22:630–5.
- 53 Kilty SJ, McLaughlin N, Bojanowski MW, *et al.* Extracranial complications of endoscopic transsphenoidal sellar surgery. *J Otolaryngol Head and Neck Surg* 2010;39:309–14.
- 54 Goudakos JK, Markou KD, Georgalas C. Endoscopic versus microscopic trans-sphenoidal pituitary surgery: a systematic review and meta-analysis. *Clin Otolaryngol* 2011;36:212–20.
- 55 Rotenberg B, Tam S, Ryu WHA, *et al.* Microscopic versus endoscopic pituitary surgery: a systematic review. *Laryngoscope* 2010;120:1292–97.
- 56 Strychowsky J, Nayan S, Reddy K, *et al.* Purely endoscopic transsphenoidal surgery versus traditional microsurgery for resection of pituitary adenomas: systematic review. *J Otolaryngol Head Neck Surg* 2011;40:175–85.
- 57 Dusk JR, Esposito F, Mattozo CA, *et al.* Endonasal transsphenoidal surgery: the patient's perspective – survey results from 259 patients. *Surg Neurol* 2006;65:332–42.
- 58 McCulloch P, Altman DG, Campbell WB, *et al.* No surgical innovation without evaluation: the IDEAL recommendations. *Lancet* 2009;374:1105–12.
- 59 Ashrafian H, Rao C, Darzi A, *et al.* Benchmarking in surgical research. *Lancet* 2009;374:1045–7.
- 60 Horton R. Surgical research or comic opera: questions, but few answers. *Lancet* 1996;347:984–5.