Methodology 40 eligible ALS patients with respiratory failure were randomised to the breath-stacking BS technique (n=21) or to mechanical insufflator/exsufflator MI-E (n=19).

Findings There were 13 episodes of chest infection in the BS group and 19 episodes in MI-E group ($p=0.92$), requiring 90 and 95 days of antibiotics respectively ($p=0.34$). The mean duration of symptoms per chest infection was 6.9 days in the BS group and 3.9 days in MI-E group ($p=0.16$). There were six episodes of hospitalisation in each group ($p=0.64$). The chance of hospitalisation, in the event of a chest infection was 0.46 in the BS group and 0.31 in MI-E group ($p=0.47$). Median survival in the BS group was 535 days and 266 days in MI-E group ($p=0.34$). The QoL was maintained above 75% of baseline for a median of 329 days in the BS group and 205 days in MI-E group ($p=0.41$).

Interpretation These results suggest non-inferiority of the BS to MI-E. We recommend that the BS technique is prescribed as a low cost, first-line intervention.

182

ROLE OF COUGH AUGMENTATION IN AMYOTROPHIC LATERAL SCLEROSIS

Muhammad Khizar Rafiq, Michael Bradburn, Alison Proctor, Catherine Billings, Stephen Bianchi, Christopher McDermott, Pamela Shaw. *University of Sheffield*

10.1136/jnnp-2014-309236.182

Background There is no clear consensus about how best to enhance cough in patients with ALS.