Book reviews

ACROMÉGALIE-GIGANTISME. TRAITEMENT CHIRURGICAL PAR EXÉRÈSE TRANSSPHÉNOÏDALE DE L'ADÉNOME HYPOPHYSAIRE. *Neurochirurgie*, 19, Suppl 2. By J. Hardy, F. Robert, M. Somma, and J. L. Vezina, with J. M. Blais, P. Dumas, A. Lanthier, R. Maltais. (Pp. 184; illustrated; price not stated.) Masson: Paris. 1973.

This important monograph describes in detail 40 acromegalic patients, including three giants, treated by transsphenoidal surgery, in the department of neurosurgery at Notre Dame Hospital, Montreal. The radiology, clinical presentation, and pre- and postoperative endocrine studies are described in detail. In 75% of the cases there was no spread of the tumour beyond the pituitary fossa, and in 25% the sella was not enlarged but showed localized bulging of the anterior inferior wall on one side. Growth hormone levels were measured in most patients and differing patterns of response to hyperglycaemia are described. Surgery was performed by the rhinoseptal transsphenoidal route, using the operating microscope and radiographic control. The aim of the treatment, when the tumour was confined to the pituitary fossa, was to remove only the adenoma and preserve the normal pituitary gland intact. This was achieved in 25 patients and in 11 of these there was recovery of previously depressed pituitary function.

Clearly, the concept of a growth hormone-secreting microadenoma which causes only minimal change in the radiograph of the pituitary fossa, and which can be selectively removed with preservation of the functioning normal pituitary gland, is an important one. However, these are slow growing tumours and careful follow-up will be needed to show whether this conservative surgery is really curative. It seems possible that some patients, treated by removal of only the adenoma with preservation of the remaining normal gland, will develop further symptoms, either from regrowth of the original tumour, or from the development of another tumour in the remaining anterior pituitary. However, it appears acceptable to treat the smaller tumours in this way, since more radical surgery can be undertaken later if necessary.

The work includes a section on the pathology, describing the gross, microscopic, and electron microscopy appearances, and the main part of the work is preceded by an interesting historical review.

The work is attractively presented and lavishly illustrated, and will be of considerable interest, particularly to endocrinologists.

This is an important therapeutic advance which deserves to be widely known.

R. D. ILLINGWORTH

MULTIPLE SCLEROSIS: CLUES TO ITS CAUSE By Uri Leibowitz and Milton Alter. (Pp. 346; illustrated; \$28.10.) North-Holland: Amsterdam. 1973.

The authors have made many important contributions to the literature on multiple sclerosis which are incorporated in this book. The first part concerns the clinical studies in Israel and a comparison of the clinical pattern of the disease in various populations: European, Afro-Asian, Oriental, and Negro. The second part is chiefly concerned with epidemiological studies in Israel and the world. Their studies of immigrant and native born populations of Israel and Hawaii are of great importance and imply that the age of 15 years marks a critical period for acquiring multiple sclerosis. However, they admit that there are pitfalls in the migration data, chiefly the small number of patients involved. Their studies of the data on ethnic groups living together in similar environments, as well as more formal evaluation of the role of genetics in the aetiology of multiple sclerosis, suggest that genetic factors are less important than environmental factors. They consider, after much speculation about possible aetiological factors suggested by epidemiological research, that the agent which causes multiple sclerosis is most likely to be connected in some way with drinking water.

The book is full of interesting epidemiological information and is a worthy memorial to Dr Leibowitz who, unfortunately, died before his book was published.

J. H. D. MILLAR

MEDICAL NEUROBIOLOGY By William D. Willis, and Robert G. Grossman. (Pp. 457; illustrated; £9.) Kimpton: London. 1973.

This book is an account of the structure and function of the nervous system and is intended as an introduction to clinical studies. As such, it is appropriate for medical students undergoing integrated courses in anatomy and physiology but it should also be helpful to graduate students as an introduction to the neuro-

logical sciences. Early sections cover the structure and physiology of neurones, sensory receptors and the organization of peripheral nerves. Later sections deal with the structure and function of the brain and spinal cord. A final chapter deals with the clinical examination of the nervous system and its applied anatomy, and the appendices include a detailed photographic atlas of the brain and spinal cord.

The presentation is thorough and concise, and care has been taken to provide a balanced account of neuroanatomy and physiology which is reasonably up to date and clinically relevant. References are given at the end of each chapter but it is a pity that these are not cited in the text as some subjects are dealt with in considerable depth, and it is difficult to identify key references for further study. The book as a whole can be thoroughly recommended and many students will be grateful to the authors who have assembled a vast amount of information in attractive form. The illustrations are clear and the price does not seem excessive for what is offered.

J. A. R. LENMAN

EVOKED POTENTIALS IN PSYCHOLOGY, SENSORY PHYSIOLOGY AND CLINICAL MEDICINE By D. Regan. (Pp. 328; illustrated; £6.) Chapman and Hall: London. 1972.

It is 27 years since George Dawson described in this Journal a method for recording the potential change evoked in the human cerebral cortex by flashes of light. With the development of cheap methods of electronic averaging the method has been widely applied to psychovisual studies. Despite early promise the results have been disappointing to the clinician except for the measurement of conduction delays in the peripheral optic pathways. Responses to more sophisticated visual stimulation such as pattern, movement, or colour are more promising. Regan has shown that evoked potentials, and in particular the steady state potentials, are capable of precise psychophysiological correlations. Although his analytical techniques are advanced, his experimental methods are relatively simple as he has the rare gift of stripping difficult problems to their essentials. This review has been delayed as I felt it necessary to study the book carefully. It has been worth the effort and the book is highly recommended.

J. A. SIMPSON

REFLEX TESTING METHODS FOR EVALUATING C.N.S. DEVELOPMENT By M. R. Fiorentino. 2nd edn. (Pp. 57; \$6.95.) Thomas: Springfield, Ill. 1973.

The relevance of this book to the diagnosis of neurological deficit and its management in early childhood is not obvious. There is an ingenuous assumption, widespread in some professional circles, that the

practice of physiotherapy is made more rational and 'scientific', as well as more effective, if abnormalities of motor development are explained in terms of primitive reflexes.

This assumption is not well-founded. We simply do not know if Sherrington's cats and Tower's monkeys provide a good enough neurophysiological and neuropathological model for what happens to the damaged developing human brain. The fact that some schemes of physical treatment of children with, for example, cerebral palsy seem to be helpful is not proof that their underlying theory is sound.

Photographic illustrations of reflex behaviour are interesting and may help physiotherapists to understand cause and effect in motor behaviour of some handicapped children, but I reject as misleading and unhelpful talk of 'three levels of "reflexive" development'—apedal, quadrupedal, and bipedal—with its teleological underpinning.

JOHN WILSON

RESEARCH ON THE CEREBRAL CIRCULATION By J. S. Meyer, M. Reivich, H. Lechner, and O. Eichhorn. (Pp. 475; illustrated; \$35.50.) Thomas: Springfield, Ill. 1973.

The proceedings of the Salzburg Conferences on *Research on the Cerebral Circulation* are now a regular feature of the cerebral blood flow scene. This volume reports the 5th conference, held late in 1970. The basic theme is a medical one, cerebral ischaemia, in the widest sense and is regarded from many viewpoints, biochemical, physiological, epidemiological, and therapeutic to mention only some.

The book, like all conference proceedings, is somewhat of a curate's egg. The 45 papers vary in length (three to 25 pages) in format (from simple reviews to presentation of original and important data), and, inevitably, in quality of presentation. The discussion is recorded verbatim, which, although recalling the flavour of the meeting for those who attended, entails the publication of much gratuitous comment. There is no index.

The price of this volume and the delay in publication will deter most individual purchasers. This is a pity because there is much of interest to neurologists, general physicians, and workers in the field of the cerebral circulation.

J. DOUGLAS MILLER

A STUDY IN NEUROLINGUISTICS By S. Locke, D. Caplan, and L. Kellar. (Pp. 135; illustrated; \$9.75.) Thomas: Springfield, Ill. 1973.

The central and largest section of this book (80 pages) consists of the case reports of 14 patients: three with dementia, five with left hemisphere lesions, three with right hemisphere lesions, and three with schizo-