cellular sites of synthesis of myelin proteins. Paterson emphasises the similarity between immune events in experimental allergic encephalomyelitis and multiple sclerosis. He refers to unpublished evidence that fragments of myelin basic protein are normally present in the blood, disappear during the development of EAE and return in higher concentrations during recovery. His interesting speculation is that these fragments and antibodies directed against them may be important in modulating the immune responses against myelin basic protein. Eylar, Ishaque and Szymanska confirm that another myelin basic protein, P2, is the neuritogenic antigen which produces experimental allergic neuritis. P2 is a highly ordered protein whose neuritogenicity depends on its conformation: Eylar's preparation of rabbit P2 has a low neuritogenicity which can be restored by the addition of phosphatidylserine. In contradiction to the finding of another group P2 was demonstrated immunocytochemically in all, rather than a proportion of peripheral nerve myelin sheaths: some staining is also present in the rabbit spinal cord. Quarles describes the myelin glycoproteins: P0 is the major structural glycoprotein of peripheral nerve myelin. Its carbohydrate moiety is located in the intraperiod line (apposed external surfaces of the Schwann cell) but the P0 molecule also protrudes into the major dense line. Myelin-associated glycoprotein (MAG) is present in both the peripheral and central nervous system myelin, although there is less in the peripheral nerves. It is present in the cytoplasm of oligodendrocytes and Schwann cells but absent from compacted myelin. It is of considerable interest because the area of loss of MAG extends further into the white matter surrounding a multiple sclerosis plaque than the loss of myelin and myelin basic protein demonstrated by Luxol Fast Blue or antimyelin basic protein serum. MAG may have an important role in the interaction between the myelin producing cell and the axon. Hashim, the editor, briefly summarises a generation of work on the encephalitogenic determinants of myelin basic protein in different species. He then describes a series of unpublished experiments demonstrating that a synthetic peptide S42 will prevent and suppress EAE. This effect can be adoptively transferred to other animals with lymph node cells. Peptide S42 is postulated to mimic a portion of the myelin basic protein molecule which

induces suppressor T cells and is separate from the encephalitogenic determinant. Confirmation of this and the link between EAE and MS would raise interesting new possibilities for the treatment of multiple sclerosis. The conciseness of this volume and its low price (\$18) make up for the cheap typescript format and frequent typographical errors.

Such a book is not the best forum for original data but as an anthology of five personal reviews by respected workers it is an excellent buy for the libraries of neurobiologists.

RAC HUGHES

British National Formulary. (Pp 400; £3.80.) London: British Medical Association and The Pharmaceutical Society of Great Britain, 1981.

This should be one of the most important books in medicine for 1981. Since the second World War neurology has changed from a diagnostic to a therapeutic speciality. How do we deal with the explosion of factual pharmacology in treating our patients?

The main advantages of the previous BNF were its introductory sections on systemic pharmacology, with selective prescribing lists, convenient size and hard back format. This 1981 total revision has lost these advantages. It is an inch too big to fit most pockets, not very durable and contains complete rather than selective lists of available drugs, often in extremely small type. There are however many gains, not least in immediate access to a great deal of information about most of the products available in the United Kingdom. The association rather than separation of prescribers notes and lists of available drugs should help sensible and informed prescribing. The drug-lists can be overwhelming, however, as with the 25 different rubifacients described, 26 oral contraceptives, 31 compound antacids, 52 compound analgesics, and no less than 92 expectorant and cough medicines. All this duplicates MIMS, although without the advantage of giving exact drug costs. The text often repeats the same cautions, contra-indications and side effects with related drugs and gives little real advice on which of many similar drugs to select.

The value of the new BNF must, therefore, lie in the prescribers notes. How good are these? Unfortunately, there are many weaknesses in the central nervous system section and little or no use is made

of tables. The many different benzodiazepines are as numerous as previous generations of barbiturates, although they may be safer in clinical practice. Little guidance is given in individual drug selection. The anti-psychotic and anti-depressant drug sections are both good, although it is incorrect to say that tardive dyskinesia is usually irreversible on drug withdrawal, or that there are significant sex differences in the response to sedative tricyclics. Anti-emetics are recommended only when the cause of vomiting is known, probably a largely unattainable ideal. New drugs are courageously advocated although this may be sometimes premature, as with isometheptene which is said to be as effective but with less side effects than ergotamine in the treatment of migraine, without stressing the combination of isometheptene with paracetamol in one preparation and the possibility of paracetamol poisoning. The antiparkinsonian section is most curious. All the available anticholinergics are described as having similar actions, without noting their different sedative or stimulant effects. The use of anticholinergics is specially recommended in treating patients with postencephalitic Parkinsonism who are also taking anti-psychotic drugs, a combination I have never encountered. Amantadine is said, incorrectly, to benefit only a small proportion of patients. The described side effects of levodopa are ludicrous, and include abdominal pain, polyuria, incontinence and difficulty with micturition. A so-called slow release preparation of levodopa is described without comment that this ensures maximum gut decarboxylation and minimum brain dopamine availability.

Most students, house officers and registrars will find the new BNF a very convenient source of drug information. Neurologists and psychiatrists may have doubts about the coverage of their own speciality, but as a quick reference to the very many drugs available and their mode of action, the new BNF is far superior to both the old edition and to MIMS, and its constant use should improve our prescribing habits. It is not intended to be a substitute for more comprehensive books such as the excellent recent *Neurological Clinical Pharmacology* by Eady and Tyrer (MTP, 1980).

JD PARKES