changes in the cerebellum (see among others case III of Refsum reported by Cammermeyer,¹³ case III of Gordon and Hudson¹⁰ and the patient reported by Reese and Bareta¹⁴). It should be remembered that in essential tremor, no pathological changes in the cerebellum have so far been found. The present author has examined clinically two patients with Refsum's disease,^{15 16} and has studied the pathology of three cases; (ref 16, case III of Gordon and Hudson;¹⁰ case 15/65, unpublished necropsy, National Hospital Queen Square through the kindness of Professor LW Duchon). In none of these latter three cases was there any structural lesion in the cerebellum. In my view, the coordination difficulties mimicking cerebellar deficit that may occur in Refsum's disease are due to a combination of proprioceptive deafferentation associated with a disorder of movement similar to essential tremor.

P SALISACHS

Neurological Service,
Residencia de la Seguridad Social,
Badalona, Barcelona,
Spain

References

- 1 Refsum S. Refsum's disease. In: Vinken PJ, Bruyn GW, DeJong JMBV eds. "System Degenerations and Atrophies". Amsterdam: Elsevier. 1975:181-229.
- 2 Salisachs P. Charcot-Marie-Tooth disease associated with "Essential tremor". Report of seven cases and a review of the literature. *J Neurol Sci*, 1976;28:17-40.
- 3 Salisachs P, Codina A, Gimenez-Roldan S, Zarranz JJ. Charcot-Marie-Tooth associated with "essential tremor" and normal and/or slightly diminished motor conduction velocity. *Eur Neurol*, 1979;18:49-58.
- 4 Salisachs P. Unusual presentation of Charcot-Marie-Tooth disease. Incoordination with absent or minimal wasting. Report of two cases. *J Neurol Sci*, 1981;50:175-80.
- 5 Harding AE, Thomas PK. The clinical features of hereditary motor and sensory neuropathy. Types I and II. *Brain*, 1980, 103:259-80.
- 6 Findley LJ, Gresty MA, Halmagyi M. A novel method of recording arm movements. A survey of common abnormalities. *Arch Neurol*, 1981;38:38-42.
- 7 Hughes JT, Brownell B. Pathology of peroneal muscular atrophy (Charcot-Marie-Tooth disease). *J Neurol Neurosurg Psychiatry*, 1972;35:648-57.
- 8 Adams RD, Victor M. *Familial Tremor. Principles of Neurology*. New York: McGraw Hill, 1977;835.
- 9 Refsum S. Heredopathia atactica polyneuritiformis. In: Sobue I, ed. "Spinocerebellar degenerations". Baltimore: University Park Press, 1980;313-38.
- 10 Gordon N, Hudson REB. Refsum's syndrome. Heredopathia atactica polyneuritiformis. A report of three cases including a study of cardiac pathology. *Brain*, 1959;82:41-55.
- 11 Cammermeyer J. Refsum's disease. Neuropathological aspects. In: Vinken PJ, Bruyn GW, DeJong JMBV. *System degenerations and atrophies*. Amsterdam: Elsevier. 1975:231-61.
- 12 Duncan GB, Parker SW, Fisher M. Acute cerebellar infarction in the PICA territory. *Arch Neurol*, 1975;32:364-8.
- 13 Cammermeyer J. Neuropathological changes in hereditary neuropathies. Manifestation of the syndrome heredopathia atactica polyneuritiformis in the presence of interstitial hypertrophic neuritis. *J Neuropath Exp Neurol*, 1956;15:340-61.
- 14 Reese H, Bareta J. Heredopathia atactica polyneuritiformis. *J Neuropath Exp Neurol*, 1950;9:385-95.
- 15 Bonduelle M, Bouygues P, Lormeau G, Deloux G, Laudat PH, Wolf LM. Maladie de Refsum. Etude des lipides du serum et des urines. *Rev Neurol*, 1966;115:933-42.
- 16 Lapresle J, Man HX, Metrau R. Documents anatomiques concernant un cas de maladie de Refsum. *Rev Neurol*, 1974;130:103-10.

Address for correspondence: Paseo de Gracia 62, Barcelona-7, Spain.

Received 9 December 1981. Accepted 25 January 1982

Matters Arising

Sir: A recent article in your journal¹ concerned the differentiation of haemorrhage and traumatic puncture in blood-stained cerebrospinal fluid (CSF) by a variety of methods. Of interest to us was the use of spectrophotometric scanning of CSF; a rather elaborate methodology involving the use of extinction criteria was given.

We believe that for routine clinical use, the technique as described is overly complicated; we propose that it can be simplified without significant loss of accuracy, and with much more rapid results. The rationale for spectrophotometric analysis of CSF is as follows: a traumatic puncture results in the presence of red blood cells, and therefore haemo-

globin, in the CSF; intracranial haemorrhage (of more than a few hours duration) results in the breakdown of the erythrocytes, with the resultant presence of bilirubin. Both substances are well-defined spectrophotometrically, and are easily detected. Bilirubin shows a peak at 455 nm, and haemoglobin has a characteristic series of peaks at 412 nm (Soret Band), 541 nm, and 576 nm.

Since neither haemoglobin nor bilirubin are normally present in CSF, the presence of either or both substances can readily be determined by a simple scan from 400 to 650 nm. Using a scanning spectrophotometer available in many hospital biochemistry laboratories, results can be obtained within minutes of receiving the sample.

We suggest the following interpretations, which are compatible with those of Buruma *et al.* An intracranial haemorrhage is suggested by the appearance of a bilirubin peak, and a traumatic puncture by the presence of the characteristic triple peaks of oxyhaemoglobin. A graph with all peaks would suggest a traumatic tap with the concomitant presence of a haemorrhage.

We have run a number of scans on clear CSF, and have shown that it does not contain any substances which could be confused with either bilirubin or haemoglobin at the specified wavelengths.

Our interpretations are based on two suppositions: firstly, that the interval between the onset of the haemorrhage and lumbar puncture is more than two or three hours (which is usually the case); and secondly, that the original intracranial event is self-limiting, in that bleeding does not continue over a prolonged period of time. This is true of most such episodes.

We conclude that adequate and rapid differentiation of haemorrhage and traumatic puncture in CSF can be made by a spectrophotometric scan as described.

AI AIZENMAN

DB TONKS

Division of Clinical Chemistry
Montreal General Hospital
1650 Cedar
Montreal, Quebec
Canada, H3G 1A4

Reference

- 1 Buruma OJS, Janson HLF, Den Burgh FAJ, Bots GTH. Blood stained cerebrospinal fluid: traumatic puncture or haemorrhage? *J Neurol Neurosurg Psychiatry* 1981;44:144-7.