


# KEEP UP WITH CURRENT OPINION



Clear and concise  
reviews by leading  
authorities

Comprehensive  
bibliographies and  
annotated references

The best way to  
stay informed

**NO RISK GUARANTEED**  
you may cancel your subscription  
any time within 30 days of receiving your  
first issue and receive a full refund

## Edited by Lord Walton & RT Johnson

Current Opinion in NEUROLOGY & NEUROSURGERY vol 5, 1992, 6 issues

	EUROPE	USA & CAN	S AMERICA	REST OF WORLD
1 year (6 issues)	<input type="checkbox"/> £ 85	<input type="checkbox"/> \$145	<input type="checkbox"/> \$180	<input type="checkbox"/> £ 105
2 years (12 issues)	<input type="checkbox"/> £ 170	<input type="checkbox"/> \$290	<input type="checkbox"/> \$325	<input type="checkbox"/> £ 190
3 years (18 issues)	<input type="checkbox"/> £ 34.95	<input type="checkbox"/> \$49.95	<input type="checkbox"/> \$84.95	<input type="checkbox"/> £ 54.95
	ISA <input type="checkbox"/>	MasterCard <input type="checkbox"/>	Card No. _____	_____

Current Science Ltd, South Quay  
Current Science Ltd, North Quay  
Philadelphia, PA 19106, USA

REST OF THE WORLD Subscriptions  
Current Science Ltd, 34-40, Cleveland St  
London, W.1F 3AF

Name \_\_\_\_\_ Title \_\_\_\_\_ Signature \_\_\_\_\_ Date \_\_\_\_\_

Cheque enclosed  Bank transfer  Please bill me  
 I am an institutional subscriber

Current Science Ltd (Canada) add GST to the current rate. £ sterling must be drawn on a UK bank. **US\$ on a US bank.**  
 Payment by cheque must provide proof of status and may only subscribe at the discount rate for a maximum of 3 years until December 1992.

Address \_\_\_\_\_  
 \_\_\_\_\_  
 \_\_\_\_\_  
 Country \_\_\_\_\_

**OR YOU CAN FAX YOUR ORDER ON +44 (0)77 580 1911**

# THE MIGRAINE TRUST

## Research Grants and Fellowships

The Migraine Trust invites applications for Research Grants and/or Fellowships in the field of research into Migraine and Cluster Headaches.

*Application forms which should be submitted (6 copies) by not later than 31st January 1993 may be obtained from:*

**The Director, The Migraine Trust,  
45 Great Ormond Street, London WC1N 3HZ.  
Telephone Number: 071 278 2676.  
Fax Number 071 831 5174.**


For your copy of the

# BMJ

## Book Catalogue

**write now to**

**Books Marketing Department,  
British Medical Journal, BMA House,  
Tavistock Square, London WC1H 9JR.**



CURRENT ISSUES IN CANCER examines the present state of management of various cancers, and aims to increase understanding of the biology of these diseases, their treatments, and their impact on people.

Topics covered include:

- Molecular genetics of cancer
- Lung cancer
- Early breast cancer
- Testicular cancer
- Colorectal cancer
- Hodgkin's and Non-Hodgkin's lymphomas
- Biological therapy
- Radiotherapy
- Quality of life

This detailed but readable book provides valuable information for non-specialists who wish to know more about the present state of knowledge in cancer and its management, and a thought provoking text for all those who work with cancer patients.

*Published November 1992 147 pages ISBN 0 7279 0775 1  
UK £9.95; Overseas £12.00 BMA members: UK £8.95; Overseas £11.00  
Prices include postage, by air abroad*

**Available from: BRITISH MEDICAL JOURNAL,  
PO Box 295, London WC1H 9TE, medical booksellers or the  
BMJ bookshop in BMA House (book tokens accepted, UK  
only).**

Please send me ..... copy/ies of

**CURRENT ISSUES IN CANCER**

I enclose ..... BMA Membership No .....

Please make cheque payable to British Medical Journal

Debit my credit card (please tick box)

Visa  American Express  Mastercard

Card No .....

Card expiry date .....

Signature .....

NAME .....

ADDRESS .....

.....

.....

..... POSTCODE .....

Please send me a BMJ book catalogue

#### Epilim Oral Forms Prescribing Information

**Presentation 1.** Epilim 200 Enteric Coated. A lilac-coloured, enteric coated tablet containing 200mg Sodium Valproate B.P.2. Epilim 500 Enteric Coated. A lilac-coloured, enteric coated tablet containing 500mg Sodium Valproate B.P.3. Epilim 100mg Crushable Tablets. A white, scored tablet containing 100mg Sodium Valproate B.P.4. Epilim Syrup. A red, cherry-flavoured syrup containing 200mg Sodium Valproate B.P. per 5ml. 5. Epilim Liquid. A red, cherry-flavoured, sugar-free liquid containing 200mg Sodium Valproate B.P. per 5ml. **Indications** In the treatment of generalized, partial or other epilepsies. In women of childbearing age Epilim should be used only in severe cases or in those resistant to other treatment. **Dosage and Administration** To be taken with or after food. Epilim may be given twice daily. Enteric coated tablets should be swallowed whole. **Monotherapy Adults:** Start at 600mg daily increasing by 200mg at 3 day intervals until control is achieved. (Maximum dose 2500mg per day.) **Children over 20kg:** Initially 400mg/day with spaced increases until control is achieved. (Usually within the range 20-30mg/kg body weight per day.) **Children under 20kg:** 20mg/kg body weight per day; in severe cases may be increased but only when plasma valproic acid levels can be monitored. Above 40mg/kg per day monitor clinical chemistry and haematological parameters. **Combined Therapy** It may be necessary to raise the dose when used with anticonvulsants which induce liver enzyme activity. Dosage of barbiturates should be reduced if sedation is observed. Optimum dosage is mainly determined by seizure control and routine measurement of plasma levels is unnecessary. **Contra-indications, Warnings** **Contra-indications:** Active liver disease, family history of severe hepatic dysfunction, hypersensitivity to sodium valproate. **Side effects:** Liver dysfunction including hepatic failure resulting in fatalities has occurred in patients whose treatment included valproic acid or sodium valproate. Patients most at risk are children particularly those under the age of three and those with congenital metabolic or degenerative disorders, organic brain disease or severe seizure disorders associated with mental retardation. The incidents mainly occurred during the first 6 months of therapy and usually involved multiple anticonvulsant therapy. Monotherapy is to be preferred in this group of patients. Clinical symptoms are more helpful than laboratory investigations in the early stages of hepatic failure. Serious or fatal hepatotoxicity may be preceded by non-specific symptoms, usually of sudden onset, such as loss of seizure control, malaise, weakness, lethargy, oedema, anorexia, vomiting abdominal pain, drowsiness, jaundice. These are an indication for immediate withdrawal of the drug. Patients should be instructed to report any such signs to the clinician for investigations should they occur. Whilst it is difficult to establish which, if any, investigation is predictive, tests which reflect protein synthesis eg prothrombin time may be most relevant. Routine measurement of liver function should be undertaken before therapy and periodically during the first six months especially in those who seem most at risk, and those with a prior history of liver disease. Hyperammonaemia without hepatic damage may occur; it is usually transient, but may occasionally present clinically. If so Epilim should be discontinued. Valproic acid inhibits platelet aggregation. Thrombocytopenia has been reported. Prior to initiation of therapy and before surgery clinicians should assure themselves that there is no undue potential for bleeding complications. Spontaneous bruising or bleeding is an indication for withdrawal of medication. Pancreatitis, tremor, weight gain, transient hair loss, increased alertness, aggressiveness, hyperactivity, irregular periods, amenorrhoea, gynaecomastia, stupor and oedema have been reported. **Drug Interactions** Epilim may potentiate neuroleptics, monoamine oxidase inhibitors and other antidepressants. Caution is recommended when administering products which have anticoagulant properties eg warfarin and salicylates. Epilim does not induce liver enzymes and loss of efficacy of oral contraceptive agents does not appear to be a problem. Phenytoin levels should be monitored, particularly the free form. **Women of Childbearing Age** An increased incidence of congenital abnormalities (including facial dysmorphism, neural tube defects and multiple malformations) has been demonstrated in offspring born to mothers with epilepsy both untreated and treated, including those treated with sodium valproate. The incidence of neural tube defects in women receiving valproate during the first trimester has been estimated to be in the region of 1%. Pregnancies should be carefully screened by alpha-fetoprotein measurement and ultrasound and if indicated amniocentesis. In all pregnancies monotherapy is to be recommended and dosage reviewed. The benefits of antiepileptic therapy during pregnancy must be evaluated against the possible risks and patients should be informed of these and the need for screening. **Product Licence Numbers** Epilim 200 Enteric Coated 0623/0006. Epilim 500 Enteric Coated 0623/0005. Epilim 100mg Crushable Tablets 0623/0015. Epilim Syrup 0623/0004. Epilim Liquid 0623/0016. **NHS Cost (March 1992)** Epilim 200 Enteric Coated 100 tablets, £6.59. Epilim 500 Enteric Coated 100 tablets, £16.45. Epilim 100mg Crushable Tablets 100 tablets, £3.99. Epilim Syrup 300ml, £6.04. Epilim Liquid 300ml, £6.04. **Further information is available from:** Sanofi Winthrop Ltd., One Onslow Street, Guildford, Surrey GU1 4YS. Telephone: Guildford (0483) 505515. Telex: 859395. Fax: (0483) 35432. \*Registered trade mark.

**References:** 1. Chadwick DW. *Epilepsia* 1987; 28 (Suppl. 2): S12-S17. 2. Gillham R.A. et al. *Epilepsy Res* 1990; 7: 219-225.

# WHAT'S EPILIM<sup>®</sup> GOT TO DO WITH THE PRICE OF EGGS?

NATIONAL LIBRARY OF MEDICINE



NLM 01642346 0



67p for half a dozen Class 1 eggs, 75p for a pound of apples, that's a total of £1.42.

Easy for you to work out, but not so easy for an epileptic patient suffering the effects of cognitive dysfunction.

Epilim controls both partial and generalised seizures with *minimal cognitive side-effects*\* and

has been shown to produce fewer cognitive difficulties than carbamazepine and phenytoin.

That's what Epilim has to do with the price of eggs



sanofi  WINTHROP

1530 R148 2  
05-03-00 MAB

WORKING IN EPILEPSY