SHORT REPORT

Application of the polymerase chain reaction to monitor *Mycobacterium tuberculosis* DNA in the CSF of patients with tuberculous meningitis after antibiotic treatment

Juei-Jueng Lin, Horng-Jyh Harn

Abstract

The polymerase chain reaction (PCR) was used to detect *Mycobacterium tuberculosis* DNA in 29 CSF specimens from seven patients with tuberculous meningitis after the start of antituberculous chemotherapy. Ten of the 13 CSF specimens taken from these patients with an initial treatment of three weeks were positive for the PCR study. By contrast, only one of the other 16 CSF specimens taken from patients treated for more than three weeks was positive. This study shows that *M tuberculosis* DNA can exist in the CSF of a patient with tuberculous meningitis for three weeks after treatment and that PCR can still be a sensitive method to detect *M tuberculosis* DNA in the CSF after the start of treatment in patients with tuberculous meningitis.

(J Neurol Neurosurg Psychiatry 1995;59:175-177)

Keywords: antituberculous chemotherapy; *Mycobacterium tuberculosis*; polymerase chain reaction; tuberculosis meningitis

Early accurate diagnosis of tuberculous meningitis remains a challenge with conventional methods. A method of polymerase chain reaction (PCR) based on DNA amplification has the sensitivity to detect small numbers of mycobacteria present in clinical specimens^{1,2} and has proved to be the most sensitive method for an accurate diagnosis of patients with tuberculous meningitis.³⁻⁸ Although studies have successfully applied PCR for the diagnosis of patients with tuberculous meningitis, using this method to monitor *Mycobacterium tuberculosis* DNA in the CSF of patients with tuberculous meningitis after the start of antituberculous treatment has not yet been established.

Department of
Neurology
J-J Lin

Department of
Pathology,
Tri-Service General
Hospital and National
Defense Medical
Center, Taipei,
Taiwan, Republic of
China
H-J Harn

Correspondence to:
Dr Juei-Jueng Lin,
Department of Neurology,
Tri-Service General
Hospital No 8, Sec 3, Ting-
Chow Road, 107, Taipei,
Taiwan, Republic of China.

Received 20 December 1994
and in revised form
30 March 1995
Accepted 12 April 1995

Table 1 CSF studies and clinical outcome of patients with tuberculous meningitis

Patient No	Age (y)/sex	Time (days)†	CSF study				Culture	Clinical outcome	Duration of follow up
			WCC*	Protein (mg/dl)	Glucose (mg/dl)	PCR			
1	76/F	0	20	60	125/255‡	+	—	Complete recovery	8 m
		6	36	84	118/306	+	—		
		25	12	56	124/212	—	—		
		83	2	48	96/156	—	—		
2	20/M	0	80	186	24	+	—	1 Complete recovery except mild slowness of movement 2 Received VP shunt§	1 y
		4	101	279	33	+	—		
		10	54	164	48	+	—		
		20	16	86	59	—	—		
		28	18	80	56	—	—		
		40	2	97	64	—	—		
3	16/F	74	6	58	70	—	—	Mild weakness in right limbs¶	18 m
		0	386	312	20	+	—		
		13	450	306	38	+	—		
		20	280	214	34	+	—		
		28	120	98	46	—	—		
		50	50	86	78	—	—		
4	24/M	112	12	54	68	—	—	1 Vegetative state 2 Decorticate posture 3 Received VP shunt	1 y
		0	100	86	25	+	+		
		7	93	109	42	+	—		
		13	119	166	46	+	—		
		28	80	138	76	+	—		
		40	7	83	77	—	—		
5	8/F	106	16	60	80	—	—	Complete recovery	10 m
		0	146	104	32	+	—		
		14	66	74	66	+	—		
		28	12	48	60	—	—		
6	35/M	0	28	76	40	+	—	1 Complete recovery 2 Seizure attacks and well treated with antiepileptic drugs	2 y
		5	40	70	33	+	—		
		16	24	56	62	—	—		
		36	6	32	54	—	—		
7	22/M	92	2	36	78	—	—	1 Mild weakness in right extremities 2 Focal dystonia in right foot 3 Received VP shunt	2 y
		0	245	324	24	+	+		
		7	278	272	54	+	—		
		18	102	86	40	—	—		
		26	82	56	58	—	—		
		74	12	48	78	—	—		

*WCC = white cell count/mm³; †time CSF samples were taken from patients with tuberculous meningitis after the start of treatment. Day 0 means the CSF samples were taken from patients before treatment; ‡CSF glucose/blood glucose; §acute hydrocephalus after a ventriculoperitoneal shunt operation; ¶focal neurological deficit caused by cerebral infarction complicated by vasculitis.

Table 2 Results of applying the polymerase chain reaction to monitor *M tuberculosis* DNA in the CSF of patients with tuberculous meningitis, after the start of treatment

Time of CSF sampling after treatment	PCR result		Total
	Positive	Negative	
<3 weeks	10 (78%)	3	13
3–6 weeks	1	8	9
>6 weeks	0	7	7
Total	11 (38%)	18	29

Materials and methods

In our previous report,⁸ we prospectively applied PCR to detect *M tuberculosis* DNA in CSF specimens taken from 18 patients with clinically suspected tuberculous meningitis. Results showed that PCR was positive in 14. After the start of antituberculous treatment, we repeatedly sampled CSF specimens from these 14 patients with PCR positives at an irregular time to monitor the existence of *M tuberculosis* DNA. Four out of the 14 patients refused lumbar puncture again after the treatment, two were lost to follow up, and one died of sepsis. This study included seven patients with 29 CSF specimens (table 1).

Methods of extraction of DNA from CSF specimens, designation of primers, and procedures of PCR and Southern blot hybridisation were carried out as described in our previous publication.⁸ A sample was regarded as positive when DNA with a molecular weight expected for the amplified product (240 base pairs) was seen on autoradiography. Each CSF specimen was tested at least twice in different batches with the same procedures to assure the reproducibility of the results.

Results

This study showed that PCR could detect *M tuberculosis* DNA in 11 of 29 CSF specimens taken from these seven patients after the start of treatment. Ten of the 11 PCR positive CSF specimens were taken from patients with an initial treatment of three weeks (13 CSF specimens, 78%; table 2). Another PCR positive CSF specimen was taken from one patient on the 28th day after treatment. Specimens taken from patients with treatment of more than six weeks were all negative.

Table 1 summarises the detailed data of CSF specimens and outcome. Mycobacterial cultures of the 29 CSF specimens were all negative despite the fact that two patients (4 and 7) had positive CSF cultures of their specimens before the start of treatment. None of the CSF samples was positive by direct smear (Ziehl-Neelsen's stain). Each patient had, however, received antituberculous chemotherapy consisting of at least a triple regimen of isoniazid, rifampicin, and pyrazinamide (or ethionamide). As well as the triple regimen, patients 6 and 7 had also received streptomycin. Outcomes of six patients were good after follow up for eight months to two

years, four patients with complete resolution and two with residual mild neurological deficits. One was in a vegetative state.

Discussion

The PCR has been used to define novel diagnostic and epidemiological data for neurological diseases based on the fact that the method is rapid, accurate, and non-invasive.⁹ For identification of mycobacterial DNA in CSF specimens, PCR has proved to be the most sensitive method for an accurate diagnosis of tuberculous meningitis.^{3–8} Nevertheless, PCR assay has problems, especially when dealing with samples containing small amounts of mycobacterial bacilli. The study of Hance *et al* showed that when using hybridisation with a ³²P-labelled oligonucleotide, the sensitivity of the application of PCR for detection of *M tuberculosis* DNA in clinical specimens could be improved.^{10,11} Some studies showed that a nested PCR protocol could also improve the sensitivity of detecting mycobacterial DNA in clinical specimens.^{6,12,13} The study of Ostergaard *et al*, however, showed that Southern hybridisation on the nested PCR product did not give further PCR positive results.¹⁴ Therefore, regarding sensitivity, the nested PCR is superior to Southern hybridisation for the detection of *M tuberculosis* DNA in CSF specimens, due to fact that nested PCR is simpler and less time consuming. Because the detection of mycobacterial DNA in our previous report was done by PCR after Southern hybridisation,⁸ we therefore still used the same method in this study to have the advantage of a comparison between the sensitivity of PCR in detecting *M tuberculosis* DNA in the CSF of patients with tuberculous meningitis before and after antituberculous treatment.

After the start of treatment, mycobacterial cultures of CSF in patients with tuberculous meningitis rarely yield a positive result,¹⁵ which was the same as found in our study. Yet our study showed that PCR could persistently detect *M tuberculosis* DNA in the CSF of patients with tuberculous meningitis within the initial three week treatment period. After treatment for more than three weeks, the PCR rarely yielded a positive result. Because treatment was effective in this study (as shown by radiology and the resolution of clinical symptoms), the persistent presence of *M tuberculosis* DNA in CSF may indicate that antituberculous treatment induces the release of mycobacterial DNA in CSF. This persistence of positive PCR results was also found in patients with neurosyphilis in whom *Treponema pallidum* DNA was detectable in the CSF several months after effective treatment¹⁸ and patients with cerebral toxoplasmosis in whom *Toxoplasma gondii* in the blood could also be detected for up to 12 days after initiation of treatment.¹⁹

When compared with our previous report,⁸ this study does not indicate a significant difference of PCR sensitivity for applying the method to detect *M tuberculosis* DNA in the

CSF of patients with tuberculous meningitis before treatment (70% of sensitivity) and after an initial treatment of three weeks (78% of sensitivity). The sensitivity of applying PCR to detect *M. tuberculosis* DNA in the CSF of patients with tuberculous meningitis having the initial treatment for three weeks is also similar to that of other studies.³⁻⁸ Therefore, we advise that it is still valuable to apply PCR in patients with clinically suspected tuberculous meningitis who promptly receive a diagnostic antituberculous trial.

This study was supported in part by a National Science Council Research Grant of the republic of China (NSC 83-0412-B-016-050). We are indebted to Miss Hui-Chin Chen and Miss Man-Ch'ing Tang for their expertise in technical support.

- 1 Brisson-Noël A, Gicquel B, Lecossier D, Levy-Frébault V, Nassif X, Hance A. Rapid diagnosis of tuberculosis by amplification of mycobacterial DNA in clinical samples. *Lancet* 1989;ii:1069-77.
- 2 Brisson-Noël A, Aznar C, Chureau C, et al. Diagnosis of tuberculosis by DNA amplification in clinical practice evaluation. *Lancet* 1991;338:364-6.
- 3 Kaneko K, Onodera O, Miyatake T, Tsuji S. Rapid diagnosis of tuberculous meningitis by polymerase chain reaction (PCR). *Neurology* 1990;40:1617-8.
- 4 Shankar P, Manjunath N, Mohan KK, et al. Rapid diagnosis of tuberculous meningitis by polymerase chain reaction. *Lancet* 1991;337:5-7.
- 5 Donald PR, Victor TC, Jordaan AM, Schoeman JF, van Helden PD. Polymerase chain reaction in the diagnosis of tuberculous meningitis. *Scand J Infect Dis* 1993;25: 613-7.
- 6 Liu PYL, Shi ZY, Lau YJ, Hu BS. Rapid diagnosis of tuberculous meningitis by a simplified nested amplification protocol. *Neurology* 1994;44:1161-4.
- 7 Folgueira L, Delgado R, Palengue E, Noriega A. Polymerase chain reaction for rapid diagnosis of tuberculous meningitis in AIDS patients. *Neurology* 1994;44: 1336-8.
- 8 Lin JJ, Harn HJ, Hsu YD, Tsao WL, Lee HS, Lee WH. Rapid diagnosis of tuberculous meningitis by polymerase chain reaction assay of cerebrospinal fluid. *J Neurol* 1995;242:147-52.
- 9 Darnell RB. The polymerase chain reaction: application to nervous system disease. *Ann Neurol* 1993;34:513-23.
- 10 Hance AJ, Grandchamp B, Levy-Frébault V, et al. Detection and identification of mycobacteria by amplification of mycobacterial DNA. *Mol Microbiol* 1989;3: 843-9.
- 11 Brisson-Noël A, Aznar C, Chureau C, et al. Diagnosis of tuberculosis by DNA amplification in clinical practice evaluation. *Lancet* 1991;338:364-6.
- 12 Miyazaki Y, Koga H, Kohno S, Kaku M. Nested polymerase chain reaction for detection of *Mycobacterium tuberculosis* in clinical samples. *J Clin Microbiol* 1993; 31:2228-32.
- 13 Narita M, Matsuzono Y, Shibata M, Tagashi T. Nested amplification protocol for the detection of *Mycobacterium tuberculosis*. *Acta Paediatr Scand* 1992;81: 997-1001.
- 14 Ostergaard L, Nielsen AK, Black FT. DNA amplification on cerebrospinal fluid for diagnosis of cerebral toxoplasmosis among HIV-positive patients with signs or symptoms of neurological disease. *Scand J Infect Dis* 1993;25:227-37.
- 15 Kennedy DH, Fallon RJ. Tuberculous meningitis. *JAMA* 1979;24:264-8.
- 16 Noordhoek GT, Wolters FC, de Jonge MEJ, Van Embden JDA. Detection by polymerase chain reaction of *Treponema pallidum* DNA in cerebrospinal fluid from neurosyphilis patients before and after antibiotic treatment. *J Clin Microbiol* 1991;29:1976-84.
- 17 Dupouy-Camet J, de Souza SL, Maslo C, et al. Detection of *Toxoplasma gondii* in venous blood from AIDS patients by polymerase chain reaction. *J Clin Microbiol* 1993;31:1866-9.