- tis by protein analysis of cerebrospinal fluid. Particular relevance of the assays of ferritin and S-100. *J Neurol Sci* 1985;67:359-69.
- 79 Boucquoy D, Chalon MP, Sindic CJ, Lamy ME, Laterre C. Herpes simplex virus type 2 meningitis without genital lesions: an immunoblot study. *J Neurol* 1990;237:285-9.
 - 80 Felgenhauer K, Reiber H. The diagnostic significance of antibody specificity indices in multiple sclerosis and herpes virus induced diseases of the nervous system. *Clin Invest* 1992;70:28-37.
 - 81 Peter JB, Bowman RL. Intra-blood-brain barrier synthesis of IgG: comparison of IgG synthesis formulas in a computer model and in 1,629 consecutive specimens. *Neurology* 1992;42:510-5.
 - 82 Forsberg P, Kam Hansen S, Fryden A. Production of specific antibodies by cerebrospinal fluid lymphocytes in patients with herpes zoster, mumps meningitis and herpes simplex virus encephalitis. *Scand J Immunol* 1986;24:261-71.
 - 83 Mathiesen T, Linde A, Olding Stenkvist E, Wahren B. Specific IgG subclass reactivity in herpes simplex encephalitis. *J Neurol* 1988;235:400-6.
 - 84 van Loon AM, van der Logt JT, Heessen FW, Postma B, Peeters MF. Diagnosis of herpes simplex virus encephalitis by detection of virus-specific immunoglobulins A and G in serum and cerebrospinal fluid by using an antibody-capture enzyme-linked immunosorbent assay. *J Clin Microbiol* 1989;27:1983-7.
 - 85 Sindic CJ, Monteyne P, Laterre EC. The intrathecal synthesis of virus-specific oligoclonal IgG in multiple sclerosis. *J Neuroimmunol* 1994;54:75-80.
 - 86 Lakeman FD, Koga J, Whitley RJ. Detection of antigen to herpes simplex virus in cerebrospinal fluid from patients with herpes simplex encephalitis. *J Infect Dis* 1987;155:1172-8.
 - 87 Aurelius E, Forsgren M, Skoldenberg B, Strannegard O. Persistent intrathecal immune activation in patients with herpes simplex encephalitis. *J Infect Dis* 1993;168:1248-52.
 - 88 Lebon P, Boutin B, Dulac O, Ponsot G, Arthuis M. Interferon gamma in acute and subacute encephalitis. *BMJ* 1988;299:9-11.
 - 89 Aurelius E, Andersson B, Forsgren M, Skoldenberg B, Strannegard O. Cytokines and other markers of intrathecal immune response in patients with herpes simplex encephalitis. *J Infect Dis* 1994;170:678-81.
 - 90 Morawetz RB, Whitley RJ, Murphy DM. Experience with brain biopsy for suspected herpes encephalitis: a review of 40 consecutive cases. *Neurosurgery* 1983;12:654-7.
 - 91 Soong SJ, Watson NE, Caddell GR, Alford CA, Jr., Whitley RJ. Use of brain biopsy for diagnostic evaluation of patients with suspected herpes simplex encephalitis: a statistical model and its clinical implications. NIAID Collaborative Antiviral Study Group. *J Infect Dis* 1991;163:17-22.
 - 92 Whitley RJ, Soong SJ, Hirsch MS, et al. Herpes simplex encephalitis: vidarabine therapy and diagnostic problems. *N Engl J Med* 1981;304:313-8.
 - 93 Whitley RJ, Alford Jr CA, Hirsch MS, et al. Vidarabine versus acyclovir therapy of herpes simplex encephalitis. *N Engl J Med* 1986;314:144-9.
 - 94 van Landingham KE, Marsteller HB, Ross GW, Hayden FG. Relapse of herpes simplex encephalitis after conventional acyclovir therapy [published erratum *JAMA* 1988; 259:2547]. *JAMA* 1988;259:1051-3.
 - 95 Barthez MA, Billard C, Santini JJ, Ruchoux MM, Grangepon MC. Relapse of herpes simplex encephalitis. *Neuropediatrics* 1987;18:3-7.
 - 96 Gulliford MC, Chandrasekera CP, Cooper RA, Murphy RP. Acyclovir treatment of herpes simplex encephalitis: experience in a district hospital. *Postgrad Med J* 1987; 63:1037-41.

NEUROLOGICAL STAMP

Claude Bernard (1813-78)

The great French physiologist Claude Bernard was born in the same year as Livingstone. Bernard was the son of a poor wine grower who began writing plays to earn money but turned to medicine on the advice of a literary critic. He began his brilliant career by assisting Magendie and becoming involved in his research on spinal nerves. His works in neurophysiology included the description, origin, and function of the chorda tympani and studies of the function of the cervical sympathetic system. While investigating the function of the submandibular gland, he showed that the sympathetic nerve was the constrictor of the blood vessels and the chorda tympani the dilator. So the fundamental facts of vasomotor physiology became known. He demonstrated that simple reflex movements were due to the influence sensory roots exerted on the motor roots and that a puncture of the floor of the fourth ventricle caused diabetes. The *piqûre diabétique* has been interpreted by many as being the result of excessive stimulation of secretory nerves to the liver.

Claude Bernard was the first to demonstrate that curare blocked the nerve stimulation of muscle while the muscle itself remained directly excitable and he carried out important work on the action of other drugs, including the opium alkaloids.

Bernard's wife and daughters deserted him when he refused to give up experimental medicine for a more lucrative practice of his profession. Towards the end of his life he published his famous *Introduction à la médecine expérimentale* (1927). The book discusses the importance

of the constancy of the internal environment, refutes the notion of the "vital force" to explain life, and emphasises the need in planning experiments for a clear initial hypothesis which has then to be proved. Because of this work he was elected to the French Academy in 1869. In 1878 Charles Edouard Sequard was appointed to the Chair of Medicine at the College de France, in succession to Claude Bernard.

He has been philatelically honoured as a great physician and founder of modern physiology. His was a national funeral, the first ever granted to a scientist in France. France issued this stamp (Stanley Gibbons 648, Scott B89) in 1939. The surtax was used for supporting unemployed intellectuals.

L F HAAS