

Subject index

- 14-3-3 protein**, Hashimoto's encephalopathy mimicking Creutzfeldt-Jakob disease: brain biopsy findings, **601**
- 22q11 deletion**, Myoclonic movement disorder associated with microdeletion of chromosome 22q11, **600**
- α synuclein** Meta-analysis of α synuclein/ NACP polymorphism in Parkinson's disease in Japan, **350**
- acalculia**, Intraoperative mapping of the cortical areas involved in multiplication and subtraction: an electrostimulation study in a patient with a left parietal glioma, **733**
- acromegaly**, Pituitary tumours and acromegaly (Pierre Marie's disease), **394**
- acute dopaminergic challenge test**, Acute dopaminergic challenge tests to assess postural/kinetic tremor of different origin: a case report, **206**
- acute head injury**, Acute head injury for the neurologist, **i3**
Acute head injury for the neurologist, **ii3**
- acute transverse myelitis**, Bladder dysfunction in acute transverse myelitis: magnetic resonance imaging and neurophysiological and urodynamic correlations, **154**
- adhesion molecules**, High concentrations of sVCAM-1 and sICAM-1 in the cerebrospinal fluid of patients with intracerebral haemorrhage are associated with poor outcome, **346**
- affective disorder**, Enduring increased risk of developing depression and mania in patients with dementia, **40**
- akinet mutism**, Distinct neuropsychological characteristics in Creutzfeldt-Jakob disease, **686**
- alcohol drinking**, Chronic alcohol use and first symptomatic epileptic seizures, **495**
- alcoholism**, Chronic alcohol use and first symptomatic epileptic seizures, **495**
- alien hand syndrome**, Posterior alien hand syndrome after a right thalamic infarct, **447**
- Alzheimer's dementia**, We are how we eat?, **358**
- Alzheimer's disease**, Alterations in brain activation during cholinergic enhancement with rivastigmine in Alzheimer's disease, **665**
Association study of three polymorphisms of TGF- β 1 gene with Alzheimer's disease, **62**
Changes in appetite, food preference, and eating habits in frontotemporal dementia and Alzheimer's disease, **371**
Detection of grey matter loss in mild Alzheimer's disease with voxel based morphometry, **657**
Differentiation of dementia with Lewy bodies from Alzheimer's disease using a dopaminergic presynaptic ligand, **134**
Genetic variability in the insulin signalling pathway may contribute to the risk of late onset Alzheimer's disease, **261**
Goal setting and attainment in Alzheimer's disease patients treated with donepezil, **500**
Hippocampal atrophy and neocortical dysfunction in early Alzheimer's disease, **470**
Meaningful treatment outcomes in Alzheimer's disease, **471**
Morphometry of the corpus callosum in patients with questionable and mild dementia, **59**
Phosphorylated tau in cerebrospinal fluid as a marker for Creutzfeldt-Jakob disease, **79**
Relation between medial temporal atrophy and functional brain activity during memory processing in Alzheimer's disease: a combined MRI and SPECT study, **508**
Visual association test to detect early dementia of the Alzheimer type, **126**
- amantadine**, Drugs for Parkinson's disease, **607**
- amnesia**, Visual association test to detect early dementia of the Alzheimer type, **126**
- amygdala**, Measurement of amygdala T2 relaxation time in temporal lobe epilepsy, **753**
- amyloid β protein**, Association study of three polymorphisms of TGF- β 1 gene with Alzheimer's disease, **62**
- amyloid neuropathy**, "Sporadic" familial amyloidotic polyneuropathy in a German patient with B cell lymphocytic leukaemia, **86**
- amyotrophic lateral sclerosis**, Progressive sudomotor dysfunction in amyotrophic lateral sclerosis, **68**
- aneurysm**, Aneurysm of the extracranial internal carotid artery presenting as the syndrome of glossopharyngeal pain and syncope, **87**
Endovascular coiling versus neurosurgical clipping in patients with a ruptured basilar tip aneurysm, **591**
Thunderclap headache, reversible cerebral arterial vasoconstriction, and unruptured aneurysms, **96**
- angiography**, Feasibility and validity of transcranial duplex sonography in patients with acute stroke, **17**
- angular gyrus**, Intraoperative mapping of the cortical areas involved in multiplication and subtraction: an electrostimulation study in a patient with a left parietal glioma, **733**
- antibodies**, Myopathy with antibodies to the signal recognition particle: clinical and pathological features, **420**
- anticholinergic**, Drugs for Parkinson's disease, **607**
- anticoagulation**, Radiology of stroke, **96**
Transoesophageal echocardiography in selecting patients for anticoagulation after ischaemic stroke or transient ischaemic attack, **29**
- antiepileptic drugs**, Reproductive dysfunction in women with epilepsy: recommendations for evaluation and management, **121**
- apolipoprotein E**, The apolipoprotein E ϵ 2 allele and decline in episodic memory, **672**
- apomorphine**, Drugs for Parkinson's disease, **607**
- appetite**, Changes in appetite, food preference, and eating habits in frontotemporal dementia and Alzheimer's disease, **371**
- aqueductal stenosis**, Festination as the leading symptom of late onset idiopathic aqueductal stenosis, **599**
- arterial aneurysms**, Concurrent arterial aneurysms in brain arteriovenous malformations with haemorrhagic presentation, **294**
- assessment**, Sensitivity of clinical and behavioural tests of spatial neglect after right hemisphere stroke, **160**
- association study**, Genetic variability in the insulin signalling pathway may contribute to the risk of late onset Alzheimer's disease, **261**
- ataxia**, Acute attacks and brain stem signs in a patient with glutamic acid decarboxylase autoantibodies, **345**
- atlanto-axial instability**, Fatal instability following "odontoid sparing" transoral decompression of a periodontoid pseudotumour, **756**
- auditory cortex**, Electrophysiological evidence for a defect in the processing of temporal sound patterns in multiple sclerosis, **561**
- auditory evoked potential**, Electrophysiological evidence for a defect in the processing of temporal sound patterns in multiple sclerosis, **561**
- autoantibodies**, Neuromyotonia and myasthenia gravis without thymoma, **344**
- automatic voluntary dissociation**, Slowly progressive Foix-Chavany-Marie syndrome associated with chronic herpes simplex encephalitis, **203**
- autonomic dysfunction**, Progressive sudomotor dysfunction in amyotrophic lateral sclerosis, **68**
- autonomic**, High blood pressure and decreased heart rate variability in the Cuban epidemic neuropathy, **71**
- axonal neuropathy**, Six novel connexin32 (GJB1) mutations in X-linked Charcot-Marie-Tooth disease, **304**
- Babinski sign**, The Babinski sign, **360**
- bacterial meningitis**, Increased serum concentrations of tissue plasminogen activator correlate with an adverse clinical outcome in patients with bacterial meningitis, **456**
- basilar artery**, Endovascular coiling versus neurosurgical clipping in patients with a ruptured basilar tip aneurysm, **591**
- bed rest**, Effectiveness of bed rest after mild traumatic brain injury: a randomised trial of no versus six days of bed rest, **167**
- Behçet's disease**, An unusual case of Behçet's disease presenting with bilateral internal carotid artery occlusion, **343**
- bladder dysfunction**, Bladder dysfunction in acute transverse myelitis: magnetic resonance imaging and neurophysiological and urodynamic correlations, **154**
- BOLD**, Increase in focal concentration of deoxyhaemoglobin during neuronal activity in cerebral ischaemic patients, **182**
- Botox**, Dysport produces intrinsically more swallowing problems than Botox: unexpected results from a conversion factor study in cervical dystonia, **604**
- brachial plexus**, Inflammation and neuropathic attacks in hereditary brachial plexus neuropathy, **45**
- brain atrophy**, Intracranial cerebrospinal fluid measurement studies in suspected idiopathic normal pressure hydrocephalus, secondary normal pressure hydrocephalus, and brain atrophy, **552**
- brain biopsy**, Hashimoto's encephalopathy mimicking Creutzfeldt-Jakob disease: brain biopsy findings, **601**
- brain damage**, Neck muscle vibration induces lasting recovery in spatial neglect, **412**
- brain dysfunction**, Hippocampal atrophy and neocortical dysfunction in early Alzheimer's disease, **470**
- brain ischaemia**, Acute hyperperfusion syndrome complicating EC-IC bypass, **88**
- brain stem**, Acute attacks and brain stem signs in a patient with glutamic acid decarboxylase autoantibodies, **345**
Cognitive dysfunction after isolated brain stem insult. An underdiagnosed cause of long term morbidity, **191**

- brain**, Demyelination in the brain as a paraneoplastic disorder: candidates include some cases of seminoma and central nervous system lymphoma, **352**
- brainstem**, MRI findings in mesenrhombencephalitis due to *Listeria monocytogenes*, **775**
- cabergoline**, Long term tolerability of high dose ergoline derived dopamine agonist therapy for the treatment of Parkinson's disease, **602**
- CAG/polyglutamine**, Paradoxical absence of nuclear inclusion in cerebellar Purkinje cells of hereditary ataxias linked to CAG expansion, **450**
- camptocormia**, **cardiac arrest**, Propofol in myoclonus status epilepticus in comatose patients following cardiac resuscitation, **94**
- cardiac embolism**, Transoesophageal echocardiography in selecting patients for anticoagulation after ischaemic stroke or transient ischaemic attack, **29**
- cardiac rhythm**, Sex differences and lateral asymmetry in heart rate modulation in patients with temporal lobe epilepsy, **73**
- Cari Loder regime**, A randomised placebo controlled exploratory study of vitamin B-12, lofepramine, and L-phenylalanine (the "Cari Loder regime") in the treatment of multiple sclerosis, **246**
- carnitine**, Fulminant progression of hyperammonaemic encephalopathy after treatment with valproate in a patient with ureterosigmoidostomy, **90**
- carotid artery occlusion**, An unusual case of Behçet's disease presenting with bilateral internal carotid artery occlusion, **343**
- carotid stenosis**, Outcome, observer reliability, and patient preferences if CTA, MRA, or Doppler ultrasound were used, individually or together, instead of digital subtraction angiography before carotid endarterectomy, **21**
- carpal tunnel syndrome**, A randomised clinical trial of oral steroids in the treatment of carpal tunnel syndrome: a long term follow up, **710**
- case-control study**, Coffee and subarachnoid haemorrhage, **112**
Risk factors for aneurysmal subarachnoid haemorrhage: the Tromsø study, **185**
- case-control**, Chronic alcohol use and first symptomatic epileptic seizures, **495**
- catecholamine hypersensitivity**, Dermal microdialysis provides evidence for hypersensitivity to noradrenaline in patients with familial dysautonomia, **299**
- cauda equina lesion**, Sexual function in men with cauda equina lesions: a clinical and electromyographic study, **715**
- CAV-3 gene**, Familial isolated hyperCKaemia associated with a new mutation in the caveolin-3 [CAV-3] gene, **65**
- caveolin-3**, Familial isolated hyperCKaemia associated with a new mutation in the caveolin-3 [CAV-3] gene, **65**
- central motor conduction time**, Central motor conduction in patients with anti-ganglioside antibody associated neuropathy syndromes and hyperreflexia, **568**
- central nervous system pathways**, Joint position sense and vibration sense: anatomical organisation and assessment, **473**
- central nervous system vascular malformations**, Prevalence of adults with brain arteriovenous malformations: a community based study in Scotland using capture-recapture analysis, **547**
- central nervous system**, Herpes simplex encephalitis, **237**
- cerebellar disease**, Typical features of cerebellar ataxic gait, **310**
- cerebellar dysfunction**, Essential tremor and cerebellar dysfunction: abnormal ballistic movements, **400**
- cerebral arteriovenous malformations**, Concurrent arterial aneurysms in brain arteriovenous malformations with haemorrhagic presentation, **294**
- cerebral blood oxygenation**, Increase in focal concentration of deoxyhaemoglobin during neuronal activity in cerebral ischaemic patients, **182**
- cerebral infarction**, Feasibility and validity of transcranial duplex sonography in patients with acute stroke, **17**
Posterior alien hand syndrome after a right thalamic infarct, **447**
- cerebral pathology**, The palmomental reflex: a useful clinical sign?, **113**
- cerebral perfusion pressure**, Acute head injury for the neurologist, **i3**
Acute head injury for the neurologist, **ii3**
Raised intracranial pressure, **i23**
Raised intracranial pressure, **ii23**
- cerebral revascularisation**, Acute hyperperfusion syndrome complicating EC-IC bypass, **88**
- cerebrovascular disease**, Cerebral Whipple's disease with a stroke-like presentation and cerebrovascular pathology, **336**
- cerebral venous thrombosis**, Hyperthyroidism with increased factor VIII procoagulant protein as a predisposing factor for cerebral venous thrombosis, **458**
- cerebrospinal fluid**, An antibody to VacA of *Helicobacter pylori* in cerebrospinal fluid from patients with Guillain-Barre syndrome, **76**
High concentrations of sVCAM-1 and sICAM-1 in the cerebrospinal fluid of patients with intracerebral haemorrhage are associated with poor outcome, **346**
- Increased serum concentrations of tissue plasminogen activator correlate with an adverse clinical outcome in patients with bacterial meningitis, **456**
- cerebrovascular disease**, Executive dysfunction and depressive symptoms in cerebrovascular disease, **462**
Neurological presentation of Fabry's disease in a 52 year old man, **340**
- cerebrovascular events**, Patent foramen ovale, cerebrovascular risk, and complement, **351**
- certification**, Certification of deaths attributable to epilepsy, **751**
- cervical dystonia**, Dysport produces intrinsically more swallowing problems than Botox: unexpected results from a conversion factor study in cervical dystonia, **604**
Oropharyngeal swallowing in craniocervical dystonia, **406**
Unilateral globus pallidus internus stimulation improves delayed onset post-traumatic cervical dystonia with an ipsilateral focal basal ganglia lesion, **588**
- cervical spine**, Surgical disorders of the cervical spine: presentation and management of common disorders, **i34**
Surgical disorders of the cervical spine: presentation and management of common disorders, **ii34**
- cervical spondylosis**, Surgical disorders of the cervical spine: presentation and management of common disorders, **i34**
Surgical disorders of the cervical spine: presentation and management of common disorders, **ii34**
- cervical spondylotic myelopathy**, Age related shift in the primary sites of involvement in cervical spondylotic myelopathy from lower to upper levels, **316**
- chemokines**, Expression of chemokines in cerebrospinal fluid and serum of patients with chronic inflammatory demyelinating polyneuropathy, **320**
- child**, Minor head injury as cause and co-factor in the aetiology of stroke in childhood: a report of eight cases, **13**
- children**, Long term psychosocial outcomes after mild head injury in early childhood, **281**
- cholinergic enhancement**, Alterations in brain activation during cholinergic enhancement with rivastigmine in Alzheimer's disease, **665**
- chondroitin sulphate**, A Japanese case of steroid responsive myopathy with deficient chondroitin sulphate, **89**
- chorea**, Palatal tremor and cognitive decline in neuroferritinopathy, **91**
- chronic hydrocephalus**, Head drop and camptocormia, **1**
Is normal pressure hydrocephalus a valid concept in 2002? A reappraisal in five questions and proposal for a new designation of the syndrome as "chronic hydrocephalus", **9**
- chronic inflammatory demyelinating polyneuropathy**, Expression of chemokines in cerebrospinal fluid and serum of patients with chronic inflammatory demyelinating polyneuropathy, **320**
- chronic inflammatory demyelinating polyradiculoneuropathy**, Risk of relapse of Guillain-Barre syndrome or chronic inflammatory demyelinating polyradiculoneuropathy following immunisation, **348**
- circadian rhythm**, Bilateral paramedian thalamic syndrome: abnormal circadian wake-sleep and autonomic functions, **772**
- clinical outcome**, High concentrations of sVCAM-1 and sICAM-1 in the cerebrospinal fluid of patients with intracerebral haemorrhage are associated with poor outcome, **346**
- clinical signs**, The Babinski sign, **360**
The palmomental reflex: a useful clinical sign?, **113**
- clinical testing**, Joint position sense and vibration sense: anatomical organisation and assessment, **473**
- clinically isolated syndromes**, Progressive ventricular enlargement in patients with clinically isolated syndromes is associated with the early development of multiple sclerosis, **141**
- cocaine**, Cocaine induced hypokalaemic periodic paralysis, **92**
- cochlear implantation**, Cochlear implantation in a profoundly deaf patient with MELAS syndrome, **97**
- coffee**, Coffee and subarachnoid haemorrhage, **112**
- cognition disorders**, Apathy in Parkinson's disease, **636**
Cognitive slowing in Parkinson's disease resolves after practice, **524**
- cognition testing**, Testing cognitive function in elderly populations: the PROSPER study, **385**
- cognition**, Cognitive dysfunction after isolated brain stem insult. An underdiagnosed cause of long term morbidity, **191**
The apolipoprotein E ε2 allele and decline in episodic memory, **672**
- cognitive dysfunction**, CJD: are there distinct neuropsychological features?, **613**
- cognitive impairment**, Palatal tremor and cognitive decline in neuroferritinopathy, **91**
- coin rotation**, Lateralised motor control: hemispheric damage and the loss of dexterity, **574**
- coma**, Coma with focal neurological signs caused by *Datura stramonium* intoxication in a young man, **458**
- complications**, Neuropsychological and psychiatric complications in endoscopic third ventriculostomy, **460**
Neuropsychological and psychiatric complications in endoscopic third ventriculostomy, **460**

- computer communication networks**, The internet, **619**
- conduction block**, Age related shift in the primary sites of involvement in cervical spondylotic myelopathy from lower to upper levels, **316**
- Connexin32**, Six novel connexin32 (GJB1) mutations in X-linked Charcot-Marie-Tooth disease, **304**
- conversion disorder**, Functional weakness and sensory disturbance, **241**
- corpus callosum**, Morphometry of the corpus callosum in patients with questionable and mild dementia, **59**
Semi-quantitative analysis of corpus callosum injury using magnetic resonance imaging indicates clinical severity in patients with diffuse axonal injury, **289**
- cortical function**, The vegetative state, **355**
- cortical stimulations**, Intraoperative mapping of the cortical areas involved in multiplication and subtraction: an electrostimulation study in a patient with a left parietal glioma, **733**
- costs**, Home based management in multiple sclerosis: results of a randomised controlled trial, **250**
- cranial nerve palsy**, MRI findings in mesenrhombencephalitis due to *Listeria monocytogenes*, **775**
- craniotomy**, Superficial temporal artery pseudoaneurysm following craniotomy, **605**
- Creutzfeldt-Jakob disease**, CJD: are there distinct neuropsychological features?, **613**
Distinct neuropsychological characteristics in Creutzfeldt-Jakob disease, **686**
Hashimoto's encephalopathy mimicking Creutzfeldt-Jakob disease: brain biopsy findings, **601**
Neuropsychological and quantitative oculometric study of a case of sporadic Creutzfeldt-Jakob disease at predementia stage, **56**
- CSF pressure measurement**, Intracranial cerebrospinal fluid measurement studies in suspected idiopathic normal pressure hydrocephalus, secondary normal pressure hydrocephalus, and brain atrophy, **552**
- CSF tap test**, Comparison between the lumbar infusion and CSF tap tests to predict outcome after shunt surgery in suspected normal pressure hydrocephalus, **721**
- CT angiography**, Outcome, observer reliability, and patient preferences if CTA, MRA, or Doppler ultrasound were used, individually or together, instead of digital subtraction angiography before carotid endarterectomy, **21**
- Cuban epidemic neuropathy**, High blood pressure and decreased heart rate variability in the Cuban epidemic neuropathy, **71**
- cytokine**, Evidence for the role of demyelination, HLA-DR alleles, and cytokines in the pathogenesis of parvovirus B19 meningoencephalitis and its sequelae, **739**
- cytokines**, Cytokine profiles in HIV seropositive patients with tuberculous meningitis, **598**
- Datura stramonium intoxication**, Coma with focal neurological signs caused by *Datura stramonium* intoxication in a young man, **458**
- deafferentation syndrome**, The neural basis of Charles Bonnet hallucinations: a hypothesis, **535**
- deep brain stimulation**, Dual channel deep brain stimulation system (Kinetra) for Parkinson's disease and essential tremor: a prospective multicentre open label clinical study, **275**
Effect of chronic pallidal deep brain stimulation on off period dystonia and sensory symptoms in advanced Parkinson's disease, **395**
Effects of deep brain stimulation and levodopa on postural sway in Parkinson's disease, **267**
The effects of deep brain stimulation and levodopa on postural sway in subjects with Parkinson's disease, **240**
Unilateral globus pallidus internus stimulation improves delayed onset post-traumatic cervical dystonia with an ipsilateral focal basal ganglia lesion, **588**
- dementia**, Enduring increased risk of developing depression and mania in patients with dementia, **40**
Goal setting and attainment in Alzheimer's disease patients treated with donepezil, **500**
Morphometry of the corpus callosum in patients with questionable and mild dementia, **59**
Potentially reversible conditions in 1000 consecutive memory clinic patients, **390**
Visual association test to detect early dementia of the Alzheimer type, **126**
- dementia with Lewy bodies**, Differentiation of dementia with Lewy bodies from Alzheimer's disease using a dopaminergic presynaptic ligand, **134**
- demyelinating neuropathy**, Six novel connexin32 (GJB1) mutations in X-linked Charcot-Marie-Tooth disease, **304**
- demyelination**, Demyelination in the brain as a paraneoplastic disorder: candidates include some cases of seminoma and central nervous system lymphoma, **352**
Evidence for the role of demyelination, HLA-DR alleles, and cytokines in the pathogenesis of parvovirus B19 meningoencephalitis and its sequelae, **739**
- depression**, Cerebral haemodynamics and depression in the elderly, **34**
- depressive symptoms**, Executive dysfunction and depressive symptoms in cerebrovascular disease, **462**
Executive dysfunction and depressive symptoms in cerebrovascular disease, **462**
- dHMN-VII**, Clinical and genetic heterogeneity in peroneal muscular atrophy associated with vocal cord weakness, **762**
- diagnosis**, A relapsing-remitting type of ocular myasthenia gravis without typical muscle fatigability, **205**
How valid is the clinical diagnosis of Parkinson's disease in the community?, **529**
Thunderclap headache, reversible cerebral arterial vasoconstriction, and unruptured aneurysms, **96**
- diffuse axonal injury**, Semi-quantitative analysis of corpus callosum injury using magnetic resonance imaging indicates clinical severity in patients with diffuse axonal injury, **289**
- diffusion weighted imaging**, Diffusion weighted magnetic resonance imaging in a case of acute Wernicke's encephalopathy, **181**
- diffusion weighted MRI**, Don't throw in the towel! A case of reversible coma, **83**
- disability evaluation**, Comparison of the psychometric characteristics of the functional independence measure, 5 item Barthel index, and 10 item Barthel index in patients with stroke, **188**
- disseminated encephalomyelitis**, Optic neuromyelitis syndrome in Brazilian patients, **429**
- distal myopathy**, Distal myopathy with tubular aggregates: a new phenotype associated with multiple deletions in mitochondrial DNA?, **207**
- donepezil**, Goal setting and attainment in Alzheimer's disease patients treated with donepezil, **500**
- dopamine agonist**, Long term tolerability of high dose ergoline derived dopamine agonist therapy for the treatment of Parkinson's disease, **602**
- Doppler ultrasonography**, Outcome, observer reliability, and patient preferences if CTA, MRA, or Doppler ultrasound were used, individually or together, instead of digital subtraction angiography before carotid endarterectomy, **21**
- dual channel device**, Dual channel deep brain stimulation system (Kinetra) for Parkinson's disease and essential tremor: a prospective multicentre open label clinical study, **275**
- dysconjugacy**, Quantitative oculo-graphic characterisation of internuclear ophthalmoparesis in multiple sclerosis: the versional dysconjugacy index Z score, **51**
- dysphagia**, Oropharyngeal swallowing in craniocervical dystonia, **406**
- Dysport**, Dysport produces intrinsically more swallowing problems than Botox: unexpected results from a conversion factor study in cervical dystonia, **604**
- dystonia**, Effect of chronic pallidal deep brain stimulation on off period dystonia and sensory symptoms in advanced Parkinson's disease, **395**
Neuropathological findings in multiple system atrophy with dystonia, **460**
- eating patterns**, We are how we eat?, **358**
- eclampsia, hypertensive encephalopathy**, Don't throw in the towel! A case of reversible coma, **83**
- elderly people**, Testing cognitive function in elderly populations: the PROSPER study, **385**
- electromyography**, Sexual function in men with cauda equina lesions: a clinical and electromyographic study, **715**
- electrophysiology**, Oropharyngeal swallowing in craniocervical dystonia, **406**
- embolisation**, Endovascular coiling versus neurosurgical clipping in patients with a ruptured basilar tip aneurysm, **591**
- encephalitis**, Herpes simplex encephalitis, **237**
MRI findings in mesenrhombencephalitis due to *Listeria monocytogenes*, **775**
- encephalopathy**, Fulminant progression of hyperammonaemic encephalopathy after treatment with valproate in a patient with ureterosigmoidostomy, **90**
- endocrine disorders**, Reproductive dysfunction in women with epilepsy: recommendations for evaluation and management, **121**
- endoscopic third ventriculostomy**, Neuropsychological and psychiatric complications in endoscopic third ventriculostomy, **460**
Neuropsychological and psychiatric complications in endoscopic third ventriculostomy, **460**
- epidemiology**, Is multiple sclerosis a sexually transmitted infection?, **439**
Multiple sclerosis in Malta in 1999, **239**
Perceived health in a population based sample of victims of the 1956 polio epidemic in the Netherlands, **695**
- epilepsy**, Bilateral hemispheric alteration of memory processes in right medial temporal lobe epilepsy, **478**
Certification of deaths attributable to epilepsy, **751**
Chronic alcohol use and first symptomatic epileptic seizures, **495**
Clinically important change in quality of life in epilepsy, **116**
Head injury for neurologists, **i8**

- Head injury for neurologists, **ii8**
- Hypoglycaemia induced by phenytoin treatment for partial status epilepticus, **349**
- Measurement of amygdala T2 relaxation time in temporal lobe epilepsy, **753**
- Neuroimaging for epilepsy: quality and not just quantity is important, **612**
- Reproductive dysfunction in women with epilepsy: recommendations for evaluation and management, **121**
- Standard magnetic resonance imaging is inadequate for patients with refractory focal epilepsy, **643**
- The legacy of vigabatrin in a regional epilepsy clinic, **327**
- Topiramate induced manic episode, **208**
- Voxel based morphometry of grey matter abnormalities in patients with medically intractable temporal lobe epilepsy: effects of side of seizure onset and epilepsy duration, **648**
- epilepsy surgery**, Long term outcome of temporal lobe epilepsy surgery: analyses of 140 consecutive patients, **486**
- episodic memory**, Bilateral hemispheric alteration of memory processes in right medial temporal lobe epilepsy, **478**
- erectile function**, Sexual function in men with cauda equina lesions: a clinical and electromyographic study, **715**
- essential tremor**, Acute dopaminergic challenge tests to assess postural/kinetic tremor of different origin: a case report, **206**
- Dual channel deep brain stimulation system (Kinetra) for Parkinson's disease and essential tremor: a prospective multicentre open label clinical study, **275**
- Essential tremor and cerebellar dysfunction: abnormal ballistic movements, **400**
- event related potential**, Abnormal verbal event related potentials in mild cognitive impairment and incipient Alzheimer's disease, **377**
- executive dysfunction**, Executive dysfunction and depressive symptoms in cerebrovascular disease, **462**
- external ventricular drain**, Failure of regular external ventricular drain exchange to reduce cerebrospinal fluid infection: result of a randomised controlled trial, **759**
- extracranial-intracranial bypass**, Non-invasive assessment of extracranial-intracranial bypass grafts using advanced ultrasound technology, **594**
- extrastriate visual cortical area**, Cortical bricks and mortar, **472**
- Fabry's disease**, Neurological presentation of Fabry's disease in a 52 year old man, **340**
- facial weakness**, Chronic asymmetric progressive external ophthalmoplegia with right facial weakness: a unique presentation of mitochondrial myopathy, **95**
- factor VIII**, Hyperthyroidism with increased factor VIII procoagulant protein as a predisposing factor for cerebral venous thrombosis, **458**
- familial amyotrophic lateral sclerosis**, Detection of preclinical motor neurone loss in SOD1 mutation carriers using motor unit number estimation, **199**
- familial dysautonomia**, Dermal microdialysis provides evidence for hypersensitivity to noradrenaline in patients with familial dysautonomia, **299**
- FDOPA, 6-[¹⁸F]-fluoro-L-dopa** Differentiating multiple system atrophy from Parkinson's disease: contribution of striatal and mid-brain MRI volumetry and multi-tracer PET imaging, **517**
- festination**, Festination as the leading symptom of late onset idiopathic aqueductal stenosis, **599**
- fetal striatal transplantation**, Fetal striatal transplantation in Huntington's disease: time for a pause, **612**
- Filippi M, Comi G, editors**. Primary progressive multiple sclerosis. book reviews, **353**
- Fisher syndrome**, Quantitative cardiovascular autonomic function study in Fisher syndrome, **333**
- fMRI**, Alterations in brain activation during cholinergic enhancement with rivastigmine in Alzheimer's disease, **665**
- focal neurological deficits**, Coma with focal neurological signs caused by *Datura stramonium* intoxication in a young man, **458**
- Foix-Chavany-Marie syndrome**, Slowly progressive Foix-Chavany-Marie syndrome associated with chronic herpes simplex encephalitis, **203**
- FP-CIT single photon emission tomography**, Differentiation of dementia with Lewy bodies from Alzheimer's disease using a dopaminergic presynaptic ligand, **134**
- frontotemporal dementia**, Changes in appetite, food preference, and eating habits in frontotemporal dementia and Alzheimer's disease, **371**
- We are how we eat?, **358**
- functional brain imaging**, Functional reorganisation of memory after traumatic brain injury: a study with H₂¹⁵O positron emission tomography, **173**
- functional brain mapping**, Intraoperative mapping of the cortical areas involved in multiplication and subtraction: an electrostimulation study in a patient with a left parietal glioma, **733**
- functional MRI**, Bilateral hemispheric alteration of memory processes in right medial temporal lobe epilepsy, **478**
- functional neurosurgery**, Effect of chronic pallidal deep brain stimulation on off period dystonia and sensory symptoms in advanced Parkinson's disease, **395**
- functional symptoms**, Are functional motor and sensory symptoms really more frequent on the left? A systematic review, **578**
- functional weakness**, Functional weakness and sensory disturbance, **241**
- gait ataxia**, Typical features of cerebellar ataxic gait, **310**
- gait disturbance**, Festination as the leading symptom of late onset idiopathic aqueductal stenosis, **599**
- gait**, Relation between walking speed and muscle strength is affected by somatosensory loss in multiple sclerosis, **313**
- gaze**, Diagnostic value of nystagmus: spontaneous and induced ocular oscillations, **615**
- gender**, Sex differences and lateral asymmetry in heart rate modulation in patients with temporal lobe epilepsy, **73**
- gene polymorphisms**, Genetic variability in the insulin signalling pathway may contribute to the risk of late onset Alzheimer's disease, **261**
- genetic risk factors**, Genotype predisposition to leukoencephalopathy, **352**
- genetics**, Relative high frequency of the c.255delA parkin gene mutation in Spanish patients with autosomal recessive parkinsonism, **582**
- The genetic basis of Parkinson's disease, **363**
- Tuberous sclerosis presenting in late adult life, **436**
- gingival hyperplasia**, Phenobarbitone induced gingival hyperplasia, **601**
- glial fibrillary acidic protein**, Glial fibrillary acidic protein in late life major depressive disorder: an immunocytochemical study, **556**
- globus pallidus**, Effect of chronic pallidal deep brain stimulation on off period dystonia and sensory symptoms in advanced Parkinson's disease, **395**
- Unilateral globus pallidus internus stimulation improves delayed onset post-traumatic cervical dystonia with an ipsilateral focal basal ganglia lesion, **588**
- glossopharyngeal pain**, Aneurysm of the extracranial internal carotid artery presenting as the syndrome of glossopharyngeal pain and syncope, **87**
- glutamic acid decarboxylase autoantibodies**, Acute attacks and brain stem signs in a patient with glutamic acid decarboxylase autoantibodies, **345**
- Goal Attainment Scaling**, Goal setting and attainment in Alzheimer's disease patients treated with donepezil, **500**
- Meaningful treatment outcomes in Alzheimer's disease, **471**
- graft**, Non-invasive assessment of extracranial-intracranial bypass grafts using advanced ultrasound technology, **594**
- granuloma**, Peripheral nerve granuloma in a patient with tuberculosis, **769**
- guidelines**, Potentially reversible conditions in 1000 consecutive memory clinic patients, **390**
- Guillain-Barre syndrome**, An antibody to VacA of *Helicobacter pylori* in cerebrospinal fluid from patients with Guillain-Barre syndrome, **76**
- Central motor conduction in patients with anti-ganglioside antibody associated neuropathy syndromes and hyperreflexia, **568**
- Miller-Fisher syndrome and Hodgkin's disease, **344**
- Risk of relapse of Guillain-Barré syndrome or chronic inflammatory demyelinating polyradiculoneuropathy following immunisation, **348**
- Sulcal abnormalities on brain magnetic resonance imaging in the Guillain-Barré syndrome, **92**
- haemangioma**, Hemiageusia resulting from a cavernous haemangioma in the brain stem, **319**
- haemodynamics**, Cerebral haemodynamics and depression in the elderly, **34**
- haemorrhagic transformation**, Radiology of stroke, **96**
- hallucinations**, The neural basis of Charles Bonnet hallucinations: a hypothesis, **535**
- Hashimoto's encephalopathy**, Hashimoto's encephalopathy mimicking Creutzfeldt-Jakob disease: brain biopsy findings, **601**
- head injury**, Head injury for neurologists, **ii8**
- Head injury outcome prediction in the emergency department: a role for protein S-100B?, **542**
- Impact of early intervention on outcome following mild head injury in adults, **330**
- headache**, Hemiageusia resulting from a cavernous haemangioma in the brain stem, **319**
- health outcomes measurement**, Multiple Sclerosis Impact Scale (MSIS-29): reliability and validity in hospital based samples, **701**
- health status**, Perceived health in a population based sample of victims of the 1956 polio epidemic in the Netherlands, **695**
- heart rate variability**, High blood pressure and decreased heart rate variability in the Cuban epidemic neuropathy, **71**
- Helicobacter pylori**, An antibody to VacA of *Helicobacter pylori* in cerebrospinal fluid from patients with Guillain-Barre syndrome, **76**

- hemigeusia**, Hemiageusia resulting from a cavernous haemangioma in the brain stem, **319**
- hemiparesis**, Minor head injury as cause and co-factor in the aetiology of stroke in childhood: a report of eight cases, **13**
- hemiplegic migraine**, Prolonged hemiplegic migraine associated with unilateral hyperperfusion on perfusion weighted magnetic resonance imaging, **202**
- hereditary ataxias**, Paradoxical absence of nuclear inclusion in cerebellar Purkinje cells of hereditary ataxias linked to CAG expansion, **450**
- hereditary neuropathy**, Inflammation and neuropathic attacks in hereditary brachial plexus neuropathy, **45**
- herpes simplex encephalitis**, Slowly progressive Foix–Chavany–Marie syndrome associated with chronic herpes simplex encephalitis, **203**
- herpes simplex virus**, Herpes simplex encephalitis, **237**
- high frequency stimulation**, Parkinson's disease: neurosurgery at an earlier stage?, **778**
- hippocampal dysfunction**, Bilateral hemispheric alteration of memory processes in right medial temporal lobe epilepsy, **478**
- hippocampus, atrophy**, Detection of grey matter loss in mild Alzheimer's disease with voxel based morphometry, **657**
- hippocampus**, Voxel based morphometry of grey matter abnormalities in patients with medically intractable temporal lobe epilepsy: effects of side of seizure onset and epilepsy duration, **648**
- HLA-DRB1**, Evidence for the role of demyelination, HLA-DR alleles, and cytokines in the pathogenesis of parvovirus B19 meningoencephalitis and its sequelae, **739**
- HMSN-IC**, Clinical and genetic heterogeneity in peroneal muscular atrophy associated with vocal cord weakness, **762**
- Hodgkin's disease** Miller-Fisher syndrome and Hodgkin's disease, **344**
- Holmes' tremor**, Vim thalamotomy for Holmes' tremor secondary to midbrain tumour, **453**
- home based management**, Home based management in multiple sclerosis: results of a randomised controlled trial, **250**
- human fetal tissue**, Unilateral transplantation of human primary fetal tissue in four patients with Huntington's disease: NEST-UK safety report ISRCTN no 36485475, **678**
- human immunodeficiency virus**, Cytokine profiles in HIV seropositive patients with tuberculous meningitis, **598**
- Huntington disease**, Fetal striatal transplantation in Huntington's disease: time for a pause, **612**
- hydrocephalus**, Festination as the leading symptom of late onset idiopathic aqueductal stenosis, **599**
- Hydrocephalus and shunts: what the neurologist should know**, **ii17**
- Hydrocephalus and shunts: what the neurologist should know**, **ii17**
- Neuropsychological and psychiatric complications in endoscopic third ventriculostomy, **460**
- hyperammonaemia**, Fulminant progression of hyperammonaemic encephalopathy after treatment with valproate in a patient with ureterosigmoidostomy, **90**
- hyperCKaemia**, Familial isolated hyperCKaemia associated with a new mutation in the caveolin-3 (CAV-3) gene, **65**
- hyperhidrosis**, Mexiletine on segmental hyperhidrosis, **351**
- hyperhidrosis**, Mexiletine on segmental hyperhidrosis, **351**
- hypertension**, Coffee and subarachnoid haemorrhage, **112**
- High blood pressure and decreased heart rate variability in the Cuban epidemic neuropathy, **71**
- Risk factors for aneurysmal subarachnoid haemorrhage: the Tromsø study, **185**
- hyperthyroidism**, Hyperthyroidism with increased factor VIII procoagulant protein as a predisposing factor for cerebral venous thrombosis, **458**
- hypokalaemia**, Cocaine induced hypokalaemic periodic paralysis, **92**
- immunisation**, Risk of relapse of Guillain-Barré syndrome or chronic inflammatory demyelinating polyradiculoneuropathy following immunisation, **348**
- infection**, Failure of regular external ventricular drain exchange to reduce cerebrospinal fluid infection: result of a randomised controlled trial, **759**
- inflammation**, Hashimoto's encephalopathy mimicking Creutzfeldt-Jakob disease: brain biopsy findings, **601**
- Inflammation and neuropathic attacks in hereditary brachial plexus neuropathy, **45**
- information**, PubMed: <http://www.pubmed.org>, **746**
- insulin**, Genetic variability in the insulin signalling pathway may contribute to the risk of late onset Alzheimer's disease, **261**
- interferon beta**, Differential effects of three interferon betas on neutralising antibodies in patients with multiple sclerosis: a follow up study in an independent laboratory, **148**
- Neutralising antibodies to interferon beta during the treatment of multiple sclerosis, **465**
- Neutralising antibodies to the beta interferons, **110**
- internal carotid artery**, Aneurysm of the extracranial internal carotid artery presenting as the syndrome of glossopharyngeal pain and syncope, **87**
- internet**, Don't slip through the net, **611**
- PubMed: <http://www.pubmed.org>, **746**
- The internet, **619**
- internuclear ophthalmoparesis**, Quantitative oculographic characterisation of internuclear ophthalmoparesis in multiple sclerosis: the versional dysconjugacy index Z score, **51**
- intervention**, Impact of early intervention on outcome following mild head injury in adults, **330**
- intervertebral disc prolapse**, Surgical disorders of the thoracic and lumbar spine: a guide for neurologists, **ii42**
- Surgical disorders of the thoracic and lumbar spine: a guide for neurologists, **ii42**
- intra-arterial angiography**, Outcome, observer reliability, and patient preferences if CTA, MRA, or Doppler ultrasound were used, individually or together, instead of digital subtraction angiography before carotid endarterectomy, **21**
- intracerebral haemorrhage**, High concentrations of sVCAM-1 and sICAM-1 in the cerebrospinal fluid of patients with intracerebral haemorrhage are associated with poor outcome, **346**
- intracranial aneurysm**, Subarachnoid haemorrhage and intracranial aneurysms: what neurologists need to know, **ii28**
- Subarachnoid haemorrhage and intracranial aneurysms: what neurologists need to know, **ii28**
- intracranial arteriovenous malformations**, Prevalence of adults with brain arteriovenous malformations: a community based study in Scotland using capture-recapture analysis, **547**
- intracranial haemorrhage**, Concurrent arterial aneurysms in brain arteriovenous malformations with haemorrhagic presentation, **294**
- intracranial pressure**, Acute head injury for the neurologist, **i3**
- Acute head injury for the neurologist, **ii3**
- intrathecal phenol**, Managing severe lower limb spasticity in multiple sclerosis: does intrathecal phenol have a role?, **705**
- investigative techniques**, Some neurological signs, **110**
- involuntary movement**, Posterior alien hand syndrome after a right thalamic infarct, **447**
- ipsilateral**, Lateralised motor control: hemispheric damage and the loss of dexterity, **574**
- ischaemia**, Glial fibrillary acidic protein in late life major depressive disorder: an immunocytochemical study, **556**
- ischaemic nerve block**, Peripheral mechanisms in tremor after traumatic neck injury, **585**
- ischaemic stroke**, Duration, frequency, recency, and type of migraine and the risk of ischaemic stroke in women of childbearing age, **747**
- Posterior alien hand syndrome after a right thalamic infarct, **447**
- joint position sense**, Joint position sense and vibration sense: anatomical organisation and assessment, **473**
- Lambert-Eaton myasthenic syndrome**, Difference in distribution of muscle weakness between myasthenia gravis and the Lambert-Eaton myasthenic syndrome, **766**
- late positive component**, Abnormal verbal event related potentials in mild cognitive impairment and incipient Alzheimer's disease, **377**
- late-life depression**, Glial fibrillary acidic protein in late life major depressive disorder: an immunocytochemical study, **556**
- laterality**, Are functional motor and sensory symptoms really more frequent on the left? A systematic review, **578**
- L-dopa**, Acute dopaminergic challenge tests to assess postural/kinetic tremor of different origin: a case report, **206**
- Drugs for Parkinson's disease, **607**
- legal aspects**, The vegetative state, **355**
- leukoaraiosis**, Genotype predisposition to leukoaraiosis, **352**
- levetiracetam**, Amelioration of spinal myoclonus with levetiracetam, **457**
- levodopa**, Effects of deep brain stimulation and levodopa on postural sway in Parkinson's disease, **267**
- Listeria**, MRI findings in mesenrhombencephalitis due to *Listeria monocytogenes*, **775**
- literature search**, PubMed: <http://www.pubmed.org>, **746**
- lofepramine**, A randomised placebo controlled exploratory study of vitamin B-12, lofepramine, and L-phenylalanine (the "Cari Loder regime") in the treatment of multiple sclerosis, **246**
- longitudinal study**, Quantitative cardiovascular autonomic function study in Fisher syndrome, **333**
- The apolipoprotein E ε2 allele and decline in episodic memory, **672**
- lower limb spasticity**, Managing severe lower limb spasticity in multiple sclerosis: does intrathecal phenol have a role?, **705**
- L-phenylalanine**, A randomised placebo controlled exploratory study of vitamin B-12, lofepramine, and L-phenylalanine (the "Cari Loder regime") in the treatment of multiple sclerosis, **246**
- lumbar infusion test**, Comparison between the lumbar infusion and CSF tap tests to predict outcome after shunt surgery in suspected normal pressure hydrocephalus, **721**
- lumbar spine disorders**, Surgical disorders of the thoracic and lumbar spine: a guide for neurologists, **ii42**
- Surgical disorders of the thoracic and lumbar spine: a guide for neurologists, **ii42**

- lumbar stenosis**, Surgical disorders of the thoracic and lumbar spine: a guide for neurologists, **i42**
Surgical disorders of the thoracic and lumbar spine: a guide for neurologists, **ii42**
- lumbosacral nerve root compression**, Diagnostic value of history and physical examination in patients suspected of lumbosacral nerve root compression, **604**
- lymphoma**, "Sporadic" familial amyloidotic polyneuropathy in a German patient with B cell lymphocytic leukaemia, **86**
- macular hole**, The neural basis of Charles Bonnet hallucinations: a hypothesis, **535**
- magnetic resonance imaging**, Sulcal abnormalities on brain magnetic resonance imaging in the Guillain-Barré syndrome, **92**
Bladder dysfunction in acute transverse myelitis: magnetic resonance imaging and neurophysiological and urodynamic correlations, **154**
Diffusion weighted magnetic resonance imaging in a case of acute Wernicke's encephalopathy, **181**
Lesion responsible for WEMINO syndrome confirmed by magnetic resonance imaging, **204**
Measurement of amygdala T2 relaxation time in temporal lobe epilepsy, **753**
MRI findings in mesenrhombencephalitis due to *Listeria monocytogenes*, **775**
Prolonged hemiplegic migraine associated with unilateral hyperperfusion on perfusion weighted magnetic resonance imaging, **202**
Semiquantitative analysis of corpus callosum injury using magnetic resonance imaging indicates clinical severity in patients with diffuse axonal injury, **289**
Standard magnetic resonance imaging is inadequate for patients with refractory focal epilepsy, **643**
Voxel based morphometry of grey matter abnormalities in patients with medically intractable temporal lobe epilepsy: effects of side of seizure onset and epilepsy duration, **648**
- manic episode**, Topiramate induced manic episode, **208**
- medial longitudinal fasciculus**, Lesion responsible for WEMINO syndrome confirmed by magnetic resonance imaging, **204**
- medial temporal lobe atrophy**, Hippocampal atrophy and neocortical dysfunction in early Alzheimer's disease, **470**
- MELAS syndrome**, Cochlear implantation in a profoundly deaf patient with MELAS syndrome, **97**
Cochlear implantation in a profoundly deaf patient with MELAS syndrome, **97**
- memory**, Functional reorganisation of memory after traumatic brain injury: a study with H₂¹⁵O positron emission tomography, **173**
Potentially reversible conditions in 1000 consecutive memory clinic patients, **390**
The apolipoprotein E ε2 allele and decline in episodic memory, **672**
- meningoencephalitis**, Evidence for the role of demyelination, HLA-DR alleles, and cytokines in the pathogenesis of parvovirus B19 meningoencephalitis and its sequelae, **739**
- mexiletine**, Mexiletine on segmental hyperhidrosis, **351**
- microdialysis**, Dermal microdialysis provides evidence for hypersensitivity to noradrenaline in patients with familial dysautonomia, **299**
- microtubule associated protein**, Phosphorylated tau in cerebrospinal fluid as a marker for Creutzfeldt-Jakob disease, **79**
- midbrain tumour**, Vim thalamotomy for Holmes' tremor secondary to midbrain tumour, **453**
- MIDD syndrome**, Cochlear implantation in a profoundly deaf patient with MELAS syndrome, **97**
- migraine**, Duration, frequency, recency, and type of migraine and the risk of ischaemic stroke in women of childbearing age, **747**
- mild head injury**, Impact of early intervention on outcome following mild head injury in adults, **239**
Long term psychosocial outcomes after mild head injury in early childhood, **281**
- mild traumatic brain injury**, Effectiveness of bed rest after mild traumatic brain injury: a randomised trial of no versus six days of bed rest, **167**
Impact of early intervention on outcome following mild head injury in adults, **239**
Prediction of post-traumatic complaints after mild traumatic brain injury: early symptoms and biochemical markers, **727**
- Miller Fisher syndrome**, Central motor conduction in patients with anti-ganglioside antibody associated neuropathy syndromes and hyperreflexia, **568**
Miller-Fisher syndrome and Hodgkin's disease, **344**
- minimally conscious state**, The vegetative state, **355**
- minor traumatic head injury**, Minor head injury as cause and co-factor in the aetiology of stroke in childhood: a report of eight cases, **13**
- mismatch negativity**, Electrophysiological evidence for a defect in the processing of temporal sound patterns in multiple sclerosis, **561**
- mitochondrial cytopathy**, Ondine's curse in a woman with Leber's hereditary optic neuropathy, **347**
- mitochondrial DNA mutation**, Cochlear implantation in a profoundly deaf patient with MELAS syndrome, **97**
- mitochondrial myopathy**, Chronic asymmetric progressive external ophthalmoplegia with right facial weakness: a unique presentation of mitochondrial myopathy, **95**
- motivation**, Apathy in Parkinson's disease, **636**
- motor asymmetry**, Lateralised motor control: hemispheric damage and the loss of deftness, **574**
- motor control**, Lateralised motor control: hemispheric damage and the loss of deftness, **574**
- motor neurone loss**, Detection of preclinical motor neurone loss in SOD1 mutation carriers using motor unit number estimation, **199**
- motor symptoms**, Are functional motor and sensory symptoms really more frequent on the left? A systematic review, **578**
- motor unit number estimation**, Detection of preclinical motor neurone loss in SOD1 mutation carriers using motor unit number estimation, **199**
- movement analysis**, Typical features of cerebellar ataxic gait, **310**
- MR angiography**, Outcome, observer reliability, and patient preferences if CTA, MRA, or Doppler ultrasound were used, individually or together, instead of digital subtraction angiography before carotid endarterectomy, **21**
- multicentre trial**, Dual channel deep brain stimulation system (Kinetra) for Parkinson's disease and essential tremor: a prospective multicentre open label clinical study, **275**
- multifocal motor neuropathy**, Central motor conduction in patients with anti-ganglioside antibody associated neuropathy syndromes and hyperreflexia, **568**
- multiple mtDNA deletions**, Distal myopathy with tubular aggregates: a new phenotype associated with multiple deletions in mitochondrial DNA?, **207**
- Multiple Sclerosis Impact Scale (MSIS-29)**, Multiple Sclerosis Impact Scale (MSIS-29): reliability and validity in hospital based samples, **701**
- multiple sclerosis**, A randomised placebo controlled exploratory study of vitamin B-12, lofepramine, and L-phenylalanine (the "Cari Loder regime") in the treatment of multiple sclerosis, **246**
Diagnostic value of nystagmus: spontaneous and induced ocular oscillations, **615**
Differential effects of three interferon betas on neutralising antibodies in patients with multiple sclerosis: a follow up study in an independent laboratory, **148**
Electrophysiological evidence for a defect in the processing of temporal sound patterns in multiple sclerosis, **561**
Home based management in multiple sclerosis: results of a randomised controlled trial, **250**
Infection and multiple sclerosis—a new hypothesis?, **358**
Is multiple sclerosis a sexually transmitted infection?, **439**
Managing severe lower limb spasticity in multiple sclerosis: does intrathecal phenol have a role?, **705**
Multiple Sclerosis Impact Scale (MSIS-29): reliability and validity in hospital based samples, **701**
Multiple sclerosis in Malta in 1999, **239**
Multiple sclerosis in Malta in 1999: an update, **256**
Neutralising antibodies to interferon beta during the treatment of multiple sclerosis, **465**
Neutralising antibodies to the beta interferons, **110**
Optic neuromyelitis syndrome in Brazilian patients, **429**
Progressive ventricular enlargement in patients with clinically isolated syndromes is associated with the early development of multiple sclerosis, **141**
Quantitative oculographic characterisation of internuclear ophthalmoparesis in multiple sclerosis: the versional dysconjugacy index Z score, **51**
Relation between walking speed and muscle strength is affected by somatosensory loss in multiple sclerosis, **313**
- multiple system atrophy**, Differentiating multiple system atrophy from Parkinson's disease: contribution of striatal and midbrain MRI volumetry and multi-tracer PET imaging, **517**
Neuropathological findings in multiple system atrophy with dystonia, **460**
Sympathetic cardiac denervation in Parkinson's disease and pure autonomic failure but not in multiple system atrophy, **776**
- muscle metabolism**, Muscle tissue oxygenation as a functional tool in the follow up of dermatomyositis, **93**
- muscular weakness**, Difference in distribution of muscle weakness between myasthenia gravis and the Lambert-Eaton myasthenic syndrome, **766**
- myasthenia gravis**, A relapsing-remitting type of ocular myasthenia gravis without typical muscle fatigability, **205**
Difference in distribution of muscle weakness between myasthenia gravis and the Lambert-Eaton myasthenic syndrome, **766**
- myelitis**, Bladder dysfunction in acute transverse myelitis: magnetic resonance imaging and neurophysiological and urodynamic correlations, **154**
- myoclonic movement disorder**, Myoclonic movement disorder associated with microdeletion of chromosome 22q11, **600**
- myoclonus**, Amelioration of spinal myoclonus with levetiracetam, **457**

- Propofol in myoclonus status epilepticus in comatose patients following cardiac resuscitation, **94**
- myopathy**, A Japanese case of steroid responsive myopathy with deficient chondroitin sulphate, **89**
Tubular aggregate myopathy with abnormal pupils and skeletal deformities, **324**
- myositis**, Myopathy with antibodies to the signal recognition particle: clinical and pathological features, **420**
- mysathenia gravis**, Neuromyotonia and myasthenia gravis without thymoma, **344**
- N400**, Abnormal verbal event related potentials in mild cognitive impairment and incipient Alzheimer's disease, **377**
- natural toxins**, Envenomation and consumption of poisonous seafood, **103**
- near-infrared spectroscopy**, Muscle tissue oxygenation as a functional tool in the follow up of dermatomyositis, **93**
- neck injury**, Peripheral mechanisms in tremor after traumatic neck injury, **585**
- neck muscle vibration**, Neck muscle vibration induces lasting recovery in spatial neglect, **412**
- neglect**, Sensitivity of clinical and behavioural tests of spatial neglect after right hemisphere stroke, **160**
- neural transplant**, Unilateral transplantation of human primary fetal tissue in four patients with Huntington's disease: NEST-UK safety report ISRCTN no 36485475, **678**
- neuroferritinopathy**, Palatal tremor and cognitive decline in neuroferritinopathy, **91**
- neuroimaging**, Functional reorganisation of memory following traumatic brain injury: a study with H₂¹⁵O PET, **111**
Neuroimaging for epilepsy: quality and not just quantity is important, **612**
- neurological picture**, Eugene Jamot (1879–1937), **656**
- neurological stamp**, Harvey Williams Cushing (1869–1939), **596**
Hideyo Noguchi (1876–1928), **147**
- neuromyotonia**, Neuromyotonia and myasthenia gravis without thymoma, **344**
- Neuronline**, PubMed: <http://www.pubmed.org>, **746**
Don't slip through the net, **611**
- neuropathology**, Neuropathological findings in multiple system atrophy with dystonia, **460**
The vegetative state, **355**
- neuropathy**, Peripheral nerve granuloma in a patient with tuberculosis, **769**
- neuropsychological tests**, Visual association test to detect early dementia of the Alzheimer type, **126**
- neuropsychological tools**, CJD: are there distinct neuropsychological features?, **613**
- neuropsychology**, Cognitive dysfunction after isolated brain stem insult. An underdiagnosed cause of long term morbidity, **191**
Distinct neuropsychological characteristics in Creutzfeldt-Jakob disease, **686**
Neuropsychological sequelae of bilateral posteroventral pallidotomy, **444**
- neurosurgery**, Neurosurgery and the neurologist, **ii**
Neurosurgery and the neurologist, **ii1**
Standard magnetic resonance imaging is inadequate for patients with refractory focal epilepsy, **643**
- neutralising antibodies**, Differential effects of three interferon betas on neutralising antibodies in patients with multiple sclerosis: a follow up study in an independent laboratory, **148**
Neutralising antibodies to interferon beta during the treatment of multiple sclerosis, **465**
Neutralising antibodies to the beta interferons, **110**
- NIRS**, Increase in focal concentration of deoxyhaemoglobin during neuronal activity in cerebral ischaemic patients, **182**
- normal pressure hydrocephalus**, Comparison between the lumbar infusion and CSF tap tests to predict outcome after shunt surgery in suspected normal pressure hydrocephalus, **721**
Head drop and camptocormia, **1**
Intracranial cerebrospinal fluid measurement studies in suspected idiopathic normal pressure hydrocephalus, secondary normal pressure hydrocephalus, and brain atrophy, **552**
Is normal pressure hydrocephalus a valid concept in 2002? A reappraisal in five questions and proposal for a new designation of the syndrome as "chronic hydrocephalus", **9**
- novel antiepileptic drug**, Topiramate induced manic episode, **208**
- novel mutations**, An adult Japanese Sanfilippo A patient with novel compound heterozygous S347F and D444G mutations in the sulphamidase gene, **777**
- nuclear inclusion**, Paradoxical absence of nuclear inclusion in cerebellar Purkinje cells of hereditary ataxias linked to CAG expansion, **450**
- nystagmus**, Diagnostic value of nystagmus: spontaneous and induced ocular oscillations, **615**
- oculopharyngeal myopathy**, A Japanese case of steroid responsive myopathy with deficient chondroitin sulphate, **89**
- ophthalmoplegia**, Chronic asymmetric progressive external ophthalmoplegia with right facial weakness: a unique presentation of mitochondrial myopathy, **95**
- opsoclonus**, Diagnostic value of nystagmus: spontaneous and induced ocular oscillations, **615**
- optic atrophy**, Ondine's curse in a woman with Leber's hereditary optic neuropathy, **347**
- optic neuromyelitis**, Optic neuromyelitis syndrome in Brazilian patients, **429**
- orbitofrontal cortex**, Changes in appetite, food preference, and eating habits in frontotemporal dementia and Alzheimer's disease, **371**
- outcome prediction**, Head injury outcome prediction in the emergency department: a role for protein S-100B?, **542**
- oxygen consumption measurement**, Muscle tissue oxygenation as a functional tool in the follow up of dermatomyositis, **93**
- palatal tremor**, Palatal tremor and cognitive decline in neuroferritinopathy, **91**
- pallidotomy**, Bilateral lesions restricted to the posteroventral pallidum are unlikely to provoke corticobulbar syndrome and psychic akinesia, **98**
Neuropsychological sequelae of bilateral posteroventral pallidotomy, **444**
- palmomental reflex**, The palmomental reflex: a useful clinical sign?, **113**
- paralysis**, Cocaine induced hypokalaemic periodic paralysis, **92**
- paraneoplastic disorder**, Demyelination in the brain as a paraneoplastic disorder: candidates include some cases of seminoma and central nervous system lymphoma, **352**
- parkin gene**, Relative high frequency of the c.255delA parkin gene mutation in Spanish patients with autosomal recessive parkinsonism, **582**
- Parkinson's disease**, Acute dopaminergic challenge tests to assess postural/kinetic tremor of different origin: a case report, **206**
Apathy in Parkinson's disease, **636**
Bilateral lesions restricted to the posteroventral pallidum are unlikely to provoke corticobulbar syndrome and psychic akinesia, **98**
Cognitive slowing in Parkinson's disease resolves after practice, **524**
Differentiating multiple system atrophy from Parkinson's disease: contribution of striatal and midbrain MRI volumetry and multi-tracer PET imaging, **517**
Drugs for Parkinson's disease, **607**
Dual channel deep brain stimulation system (Kinetra) for Parkinson's disease and essential tremor: a prospective multicentre open label clinical study, **275**
Effect of chronic pallidal deep brain stimulation on off period dystonia and sensory symptoms in advanced Parkinson's disease, **395**
Effects of deep brain stimulation and levodopa on postural sway in Parkinson's disease, **267**
How valid is the clinical diagnosis of Parkinson's disease in the community?, **529**
Long term tolerability of high dose ergoline derived dopamine agonist therapy for the treatment of Parkinson's disease, **602**
Meta-analysis of α synuclein/ NACP polymorphism in Parkinson's disease in Japan, **350**
Neuropsychological sequelae of bilateral posteroventral pallidotomy, **444**
Parkinson's disease: neurosurgery at an earlier stage?, **778**
Relative high frequency of the c.255delA parkin gene mutation in Spanish patients with autosomal recessive parkinsonism, **582**
Sympathetic cardiac denervation in Parkinson's disease and pure autonomic failure but not in multiple system atrophy, **776**
The effects of deep brain stimulation and levodopa on postural sway in subjects with Parkinson's disease, **240**
The genetic basis of Parkinson's disease, **363**
The Parkinson's disease sleep scale: a new instrument for assessing sleep and nocturnal disability in Parkinson's disease, **629**
- paroxysmal discharges**, The neural basis of Charles Bonnet hallucinations: a hypothesis, **535**
- parvovirus B19**, Evidence for the role of demyelination, HLA-DR alleles, and cytokines in the pathogenesis of parvovirus B19 meningoencephalitis and its sequelae, **739**
- patent foramen ovale**, Patent foramen ovale, cerebrovascular risk, and complement, **351**
- patient information**, Impact of early intervention on outcome following mild head injury in adults, **239**
- patient perspective**, Meaningful treatment outcomes in Alzheimer's disease, **471**
- perfusion weighted imaging**, Prolonged hemiplegic migraine associated with unilateral hyperperfusion on perfusion weighted magnetic resonance imaging, **202**
- pergolide**, Long term tolerability of high dose ergoline derived dopamine agonist therapy for the treatment of Parkinson's disease, **602**
- Periodic limb movement syndrome**, Periodic limb movement syndrome, **428**
- peripheral receptors**, Joint position sense and vibration sense: anatomical organisation and assessment, **473**

- peroneal muscular atrophy**, Clinical and genetic heterogeneity in peroneal muscular atrophy associated with vocal cord weakness, **762**
- phenobarbitone**, Phenobarbitone induced gingival hyperplasia, **601**
- phenytoin**, Hypoglycaemia induced by phenytoin treatment for partial status epilepticus, **349**
- physical disability**, Characteristics of patients with persistent severe disability and medically unexplained neurological symptoms: a pilot study, **307**
- physical signs**, Some neurological signs, **110**
- Pierre Marie's disease**, Pituitary tumours and acromegaly (Pierre Marie's disease), **394**
- pituitary tumours**, Pituitary tumours and acromegaly (Pierre Marie's disease), **394**
- poisons**, Envenomation and consumption of poisonous seafood, **103**
- polymerase chain reaction**, Cerebral Whipple's disease with a stroke-like presentation and cerebrovascular pathology, **336**
- polymorphism**, Association study of three polymorphisms of TGF- β 1 gene with Alzheimer's disease, **62**
Meta-analysis of α synuclein/ NACP polymorphism in Parkinson's disease in Japan, **350**
- positron emission tomography**, Differentiating multiple system atrophy from Parkinson's disease: contribution of striatal and midbrain MRI volumetry and multi-tracer PET imaging, **517**
- posterior reversible encephalopathy syndrome**, Don't throw in the towel! A case of reversible coma, **83**
- postganglionic sympathetic nerve**, Sympathetic cardiac denervation in Parkinson's disease and pure autonomic failure but not in multiple system atrophy, **776**
- postpoliomyelitis syndrome**, Perceived health in a population based sample of victims of the 1956 polio epidemic in the Netherlands, **695**
- post-traumatic complaints**, Prediction of post-traumatic complaints after mild traumatic brain injury: early symptoms and biochemical markers, **727**
- postural instability**, The effects of deep brain stimulation and levodopa on postural sway in subjects with Parkinson's disease, **240**
- posture control**, Effects of deep brain stimulation and levodopa on postural sway in Parkinson's disease, **267**
- practice effects**, Testing cognitive function in elderly populations: the PROSPER study, **385**
- prediction**, Prediction of post-traumatic complaints after mild traumatic brain injury: early symptoms and biochemical markers, **727**
- prevalence**, Multiple sclerosis in Malta in 1999, **239**
Multiple sclerosis in Malta in 1999: an update, **256**
Prevalence of adults with brain arteriovenous malformations: a community based study in Scotland using capture-recapture analysis, **547**
- primary visual cortex**, Cortical bricks and mortar, **472**
- PRNP genotype** CJD: are there distinct neuropsychological features?, **613**
- propofol**, Propofol in myoclonus status epilepticus in comatose patients following cardiac resuscitation, **94**
- pseudotumour**, Fatal instability following "odontoid sparing" transoral decompression of a periodontoid pseudotumour, **756**
- psychometric methods**, Multiple Sclerosis Impact Scale (MSIS-29): reliability and validity in hospital based samples, **701**
- psychosis**, Topiramate induced manic episode, **208**
- psychosocial outcome**, Long term psychosocial outcomes after mild head injury in early childhood, **281**
- pupil**, Tubular aggregate myopathy with abnormal pupils and skeletal deformities, **324**
- pure autonomic failure**, Sympathetic cardiac denervation in Parkinson's disease and pure autonomic failure but not in multiple system atrophy, **776**
- Purkinje cell**, Paradoxical absence of nuclear inclusion in cerebellar Purkinje cells of hereditary ataxias linked to CAG expansion, **450**
- quality of life**, Clinically important change in quality of life in epilepsy, **116**
Home based management in multiple sclerosis: results of a randomised controlled trial, **250**
- quantitative autonomic function study**, Quantitative cardiovascular autonomic function study in Fisher syndrome, **333**
- raised intracranial pressure**, Raised intracranial pressure, **i23**
Raised intracranial pressure, **ii23**
- regional cerebral blood flow**, Relation between medial temporal atrophy and functional brain activity during memory processing in Alzheimer's disease: a combined MRI and SPECT study, **508**
- rehabilitation**, Cognitive dysfunction after isolated brain stem insult. An underdiagnosed cause of long term morbidity, **191**
Cognitive neuroscience and brain rehabilitation: a promise kept, **357**
- Neck muscle vibration induces lasting recovery in spatial neglect, **412**
Viewing less to see better, **195**
- relapse**, Risk of relapse of Guillain-Barré syndrome or chronic inflammatory demyelinating polyradiculoneuropathy following immunisation, **348**
- reliability**, Comparison of the psychometric characteristics of the functional independence measure, 5 item Barthel index, and 10 item Barthel index in patients with stroke, **188**
Testing cognitive function in elderly populations: the PROSPER study, **385**
- respiratory failure**, Ondine's curse in a woman with Leber's hereditary optic neuropathy, **347**
- responsiveness**, Comparison of the psychometric characteristics of the functional independence measure, 5 item Barthel index, and 10 item Barthel index in patients with stroke, **188**
- reversible conditions**, Potentially reversible conditions in 1000 consecutive memory clinic patients, **390**
- rheumatoid arthritis**, Surgical disorders of the cervical spine: presentation and management of common disorders, **134**
Surgical disorders of the cervical spine: presentation and management of common disorders, **ii34**
- rhythmic isometric exercise**, Muscle tissue oxygenation as a functional tool in the follow up of dermatomyositis, **93**
- risk factors**, Coffee and subarachnoid haemorrhage, **112**
Concurrent arterial aneurysms in brain arteriovenous malformations with haemorrhagic presentation, **294**
Patent foramen ovale, cerebrovascular risk, and complement, **351**
Risk factors for aneurysmal subarachnoid haemorrhage: the Tromsø study, **185**
- rivastigmine**, Alterations in brain activation during cholinergic enhancement with rivastigmine in Alzheimer's disease, **665**
- S-100B protein**, Head injury outcome prediction in the emergency department: a role for protein S-100B?, **542**
- saccade**, Neuropsychological and quantitative oculometric study of a case of sporadic Creutzfeldt-Jakob disease at predementia stage, **56**
- saccadic velocity**, Neuropsychological and quantitative oculometric study of a case of sporadic Creutzfeldt-Jakob disease at predementia stage, **56**
- safety trial**, Unilateral transplantation of human primary fetal tissue in four patients with Huntington's disease: NEST-UK safety report ISRCTN no 36485475, **678**
- Sanfilippo A**, An adult Japanese Sanfilippo A patient with novel compound heterozygous S347F and D444G mutations in the sulphamidase gene, **777**
- sciatica**, Diagnostic value of history and physical examination in patients suspected of lumbosacral nerve root compression, **604**
- seizure outcome**, Long term outcome of temporal lobe epilepsy surgery: analyses of 140 consecutive patients, **486**
- sensory disturbance**, Functional weakness and sensory disturbance, **241**
- sensory symptoms**, Are functional motor and sensory symptoms really more frequent on the left? A systematic review, **578**
- sexual function**, Sexual function in men with cauda equina lesions: a clinical and electromyographic study, **715**
- sexually transmitted infection**, Infection and multiple sclerosis—a new hypothesis?, **358**
Is multiple sclerosis a sexually transmitted infection?, **439**
- shunts**, Hydrocephalus and shunts: what the neurologist should know, **i17**
Hydrocephalus and shunts: what the neurologist should know, **ii17**
- signal recognition particle**, Myopathy with antibodies to the signal recognition particle: clinical and pathological features, **420**
- single fibre electromyography**, A relapsing-remitting type of ocular myasthenia gravis without typical muscle fatigability, **205**
- sleep disorder**, Periodic limb movement syndrome, **428**
- sleep**, Bilateral paramedian thalamic syndrome: abnormal circadian wake-sleep and autonomic functions, **772**
The Parkinson's disease sleep scale: a new instrument for assessing sleep and nocturnal disability in Parkinson's disease, **629**
- smoking**, Risk factors for aneurysmal subarachnoid haemorrhage: the Tromsø study, **185**
- social phobia**, Social phobia in spasmodic torticollis: some conceptual issues, **461**
- SOD1 mutation**, Detection of preclinical motor neurone loss in SOD1 mutation carriers using motor unit number estimation, **199**
- somatosensory loss**, Relation between walking speed and muscle strength is affected by somatosensory loss in multiple sclerosis, **313**
- spasmodic torticollis**, Social phobia in spasmodic torticollis: some conceptual issues, **461**
- spatial attention**, Viewing less to see better, **195**
- spatial neglect**, Cognitive neuroscience and brain rehabilitation: a promise kept, **357**

- Neck muscle vibration induces lasting recovery in spatial neglect, **412**
- spatial representation**, Viewing less to see better, **195**
- SPECT**, Relation between medial temporal atrophy and functional brain activity during memory processing in Alzheimer's disease: a combined MRI and SPECT study, **508**
- spinal degenerative disease**, Surgical disorders of the cervical spine: presentation and management of common disorders, **i34**
Surgical disorders of the cervical spine: presentation and management of common disorders, **ii34**
- spinal evoked potential**, Age related shift in the primary sites of involvement in cervical spondylotic myelopathy from lower to upper levels, **316**
- spinal tumours**, Surgical disorders of the thoracic and lumbar spine: a guide for neurologists, **i42**
Surgical disorders of the thoracic and lumbar spine: a guide for neurologists, **ii42**
- sporadic**, CJD: are there distinct neuropsychological features?, **613**
- sport diving**, Patent foramen ovale, cerebrovascular risk, and complement, **351**
- status epilepticus**, Hypoglycaemia induced by phenytoin treatment for partial status epilepticus, **349**
- steroids**, A Japanese case of steroid responsive myopathy with deficient chondroitin sulphate, **89**
A randomised clinical trial of oral steroids in the treatment of carpal tunnel syndrome: a long term follow up, **710**
- stroke**, Cerebral Whipple's disease with a stroke-like presentation and cerebrovascular pathology, **336**
Cognitive neuroscience and brain rehabilitation: a promise kept, **357**
Concurrent arterial aneurysms in brain arteriovenous malformations with haemorrhagic presentation, **294**
Feasibility and validity of transcranial duplex sonography in patients with acute stroke, **17**
Increase in focal concentration of deoxyhaemoglobin during neuronal activity in cerebral ischaemic patients, **182**
Minor head injury as cause and co-factor in the aetiology of stroke in childhood: a report of eight cases, **13**
Radiology of stroke, **96**
Sensitivity of clinical and behavioural tests of spatial neglect after right hemisphere stroke, **160**
Transoesophageal echocardiography in selecting patients for anticoagulation after ischaemic stroke or transient ischaemic attack, **29**
- subarachnoid haemorrhage**, Coffee and subarachnoid haemorrhage, **112**
Endovascular coiling versus neurosurgical clipping in patients with a ruptured basilar tip aneurysm, **591**
Risk factors for aneurysmal subarachnoid haemorrhage: the Tromsø study, **185**
Subarachnoid haemorrhage and intracranial aneurysms: what neurologists need to know, **i28**
Subarachnoid haemorrhage and intracranial aneurysms: what neurologists need to know, **ii28**
- subthalamic nucleus**, Parkinson's disease: neurosurgery at an earlier stage?, **778**
- sudden death**, Certification of deaths attributable to epilepsy, **751**
- sudomotor function**, Progressive sudomotor dysfunction in amyotrophic lateral sclerosis, **68**
- surgical treatment**, Surgical treatment of temporal lobe epilepsy, **470**
- survival analysis**, Enduring increased risk of developing depression and mania in patients with dementia, **40**
- swallowing**, Dysport produces intrinsically more swallowing problems than Botox: unexpected results from a conversion factor study in cervical dystonia, **604**
Oropharyngeal swallowing in craniocervical dystonia, **406**
- syncope**, Aneurysm of the extracranial internal carotid artery presenting as the syndrome of glossopharyngeal pain and syncope, **87**
- task complexity**, Lateralised motor control: hemispheric damage and the loss of deftness, **574**
- taste**, Hemiageusia resulting from a cavernous haemangioma in the brain stem, **319**
- temporal artery pseudoaneurysm**, Superficial temporal artery pseudoaneurysm following craniotomy, **605**
- temporal lobe epilepsy**, Long term outcome of temporal lobe epilepsy surgery: analyses of 140 consecutive patients, **486**
Sex differences and lateral asymmetry in heart rate modulation in patients with temporal lobe epilepsy, **73**
Surgical treatment of temporal lobe epilepsy, **470**
- temporal lobectomy**, Long term outcome of temporal lobe epilepsy surgery: analyses of 140 consecutive patients, **486**
Surgical treatment of temporal lobe epilepsy, **470**
- TGF-β1**, Association study of three polymorphisms of TGF-β1 gene with Alzheimer's disease, **62**
- thalamus**, Bilateral paramedian thalamic syndrome: abnormal circadian wake-sleep and autonomic functions, **772**
- thear eminence**, The palmomental reflex: a useful clinical sign?, **113**
- thiamine**, Diffusion weighted magnetic resonance imaging in a case of acute Wernicke's encephalopathy, **181**
- thoracic spine disorders**, Surgical disorders of the thoracic and lumbar spine: a guide for neurologists, **i42**
Surgical disorders of the thoracic and lumbar spine: a guide for neurologists, **ii42**
- thunderclap headache**, Thunderclap headache, reversible cerebral arterial vasoconstriction, and unruptured aneurysms, **96**
- tissue type plasminogen activator**, Increased serum concentrations of tissue plasminogen activator correlate with an adverse clinical outcome in patients with bacterial meningitis, **456**
PubMed: <http://www.pubmed.org>, **746**
- topiramate**, Topiramate induced manic episode, **208**
- transcranial Doppler**, Cerebral haemodynamics and depression in the elderly, **34**
- transoesophageal echocardiography**, Transoesophageal echocardiography in selecting patients for anticoagulation after ischaemic stroke or transient ischaemic attack, **29**
- transoral surgery**, Fatal instability following "odontoid sparing" transoral decompression of a periodontoid pseudotumour, **756**
- traumatic brain injury**, Functional reorganisation of memory after traumatic brain injury: a study with H₂¹⁵O positron emission tomography, **173**
Functional reorganisation of memory following traumatic brain injury: a study with H₂¹⁵O PET, **111**
Head injury for neurologists, **i8**
Head injury for neurologists, **ii8**
Impact of early intervention on outcome following mild head injury in adults, **330**
- tremor**, Peripheral mechanisms in tremor after traumatic neck injury, **585**
- triphasic pattern**, Essential tremor and cerebellar dysfunction: abnormal ballistic movements, **400**
- Tropheryma whippelii**, Cerebral Whipple's disease with a stroke-like presentation and cerebrovascular pathology, **336**
- tropical medicine**, Envenomation and consumption of poisonous seafood, **103**
- tuberculosis**, Peripheral nerve granuloma in a patient with tuberculosis, **769**
- tuberculous meningitis**, Cytokine profiles in HIV seropositive patients with tuberculous meningitis, **598**
- tuberous sclerosis**, Tuberous sclerosis presenting in late adult life, **436**
- tubular aggregates**, Distal myopathy with tubular aggregates: a new phenotype associated with multiple deletions in mitochondrial DNA?, **207**
Tubular aggregate myopathy with abnormal pupils and skeletal deformities, **324**
- tumefactive plaque, multiple sclerosis**, Tumefactive multiple sclerosis plaque, **85**
- ultrasound**, Feasibility and validity of transcranial duplex sonography in patients with acute stroke, **17**
Non-invasive assessment of extracranial-intracranial bypass grafts using advanced ultrasound technology, **594**
- unexplained disability**, Characteristics of patients with persistent severe disability and medically unexplained neurological symptoms: a pilot study, **307**
- ureterosigmoidostomy**, Fulminant progression of hyperammonaemic encephalopathy after treatment with valproate in a patient with ureterosigmoidostomy, **90**
- urodynamics**, Bladder dysfunction in acute transverse myelitis: magnetic resonance imaging and neurophysiological and urodynamic correlations, **154**
- VacA antibody**, An antibody to VacA of *Helicobacter pylori* in cerebrospinal fluid from patients with Guillain-Barre syndrome, **76**
- validity**, Comparison of the psychometric characteristics of the functional independence measure, 5 item Barthel index, and 10 item Barthel index in patients with stroke, **188**
- valproate**, Fulminant progression of hyperammonaemic encephalopathy after treatment with valproate in a patient with ureterosigmoidostomy, **90**
- variant Creutzfeldt-Jakob disease**, Irregular presence of abnormal prion protein in appendix in variant Creutzfeldt-Jakob disease, **597**
- vasculitis**, Inflammation and neuropathic attacks in hereditary brachial plexus neuropathy, **45**
- vegetative state**, The vegetative state, **355**
- ventricles**, Progressive ventricular enlargement in patients with clinically isolated syndromes is associated with the early development of multiple sclerosis, **141**
- verbal memory**, Abnormal verbal event related potentials in mild cognitive impairment and incipient Alzheimer's disease, **377**
- vertebral osteomyelitis**, Surgical disorders of the thoracic and lumbar spine: a guide for neurologists, **i42**

- Surgical disorders of the thoracic and lumbar spine: a guide for neurologists, **ii42**
- vibration sense**, Joint position sense and vibration sense: anatomical organisation and assessment, **473**
- vigabatrin**, The legacy of vigabatrin in a regional epilepsy clinic, **327**
- Vim thalamotomy**, Vim thalamotomy for Holmes' tremor secondary to midbrain tumour, **453**
- visual field defects**, The legacy of vigabatrin in a regional epilepsy clinic, **327**
- visual hallucinations**, Cortical bricks and mortar, **472**
- visuospatial neglect**, Viewing less to see better, **195**
- vitamin B-12**, A randomised placebo controlled exploratory study of vitamin B-12, lofepramine, and L-phenylalanine (the "Cari Loder regime") in the treatment of multiple sclerosis, **246**
- vocal cord weakness**, Clinical and genetic heterogeneity in peroneal muscular atrophy associated with vocal cord weakness, **762**
- voxel based morphometry**, Detection of grey matter loss in mild Alzheimer's disease with voxel based morphometry, **657**
- Voxel based morphometry of grey matter abnormalities in patients with medically intractable temporal lobe epilepsy: effects of side of seizure onset and epilepsy duration, **648**
- WEMINO syndrome**, Lesion responsible for WEMINO syndrome confirmed by magnetic resonance imaging, **204**
- Wernicke's encephalopathy**, Diffusion weighted magnetic resonance imaging in a case of acute Wernicke's encephalopathy, **181**
- Whipple's disease**, Cerebral Whipple's disease with a stroke-like presentation and cerebrovascular pathology, **336**
- working memory**, Cognitive slowing in Parkinson's disease resolves after practice, **524**
- X-linked Charcot-Marie-Tooth disease**, Six novel connexin32 (GJB1) mutations in X-linked Charcot-Marie-Tooth disease, **304**