

008

**ALEMTUZUMAB: LONG TERM FOLLOW-UP IN A SINGLE CENTRE COHORT**

Mark Willis, Katharine Harding, Mark Wardle, Trevor Pickersgill, Valentina Tomassini, Neil Robertson. *Institute of Psychological Medicine & Clinical Neuroscience, Cardiff University*

10.1136/jnnp-2014-309236.8

**Introduction** Alemtuzumab is an anti-CD52 monoclonal antibody, approved for the treatment of relapsing multiple sclerosis (RMS). However, it has significant side effects and long-term follow-up is required to provide accurate information for patient management and counselling.

**Methods** Patients were followed prospectively within an active surveillance program and standardised data collected on a regional database. Data on patient demographics, adverse events, disability trajectories (EDSS), relapses and retreatment intervals were analysed.

**Results** 96 patients with highly active RMS were identified (69F:27M). Mean follow-up post treatment was 5.1 years. Most patients received 2 annual cycles but 27 (28.1%) received 3, 7 (7.3%) received 4 and 2 (2.1%) received 5. Pre- and post-treatment annualised relapse rates were 2.1 and 0.4 respectively. Mean change in EDSS from baseline was 0.44. 8 patients

developed secondary progression a mean of 2.8 years post treatment. A range of autoimmune disorders (AID), notably thyroid disease was identified in 45 (47%).

**Discussion** Our data supports the effectiveness of alemtuzumab in patients with RMS in reducing relapse rates. However, despite a paucity of inflammatory disease activity, disability accumulation continued in some patients. Overall retreatment rate was 37.5% and AID, whilst common, could be effectively identified through routine surveillance processes and managed without serious consequence.