

## **Additional References**

- s1. Das I, Krzyzosiak A, Schneider K, et al. Preventing proteostasis diseases by selective inhibition of a phosphatase regulatory subunit. *Science* 2015;348:239-242.
- s2. Jerath NU, Shy ME. Hereditary motor and sensory neuropathies: Understanding molecular pathogenesis could lead to future treatment strategies. *Biochimica et biophysica acta* 2015;1852:667-678.
- s3. Devaux JJ, Scherer SS. Altered ion channels in an animal model of Charcot-Marie-Tooth disease type IA. *The Journal of neuroscience* 2005;25:1470-1480.
- s4. Bai Y, Ianokova E, Pu Q, et al. Effect of an R69C mutation in the myelin protein zero gene on myelination and ion channel subtypes. *Archives of neurology* 2006;63:1787-1794.
- s5. Rosberg MR, Alvarez S, Krarup C, Moldovan M. An oral NaV1.8 blocker improves motor function in mice completely deficient of myelin protein P0. *Neuroscience letters* 2016;632:33-38.
- s6. Nave KA, Sereda MW, Ehrenreich H. Mechanisms of disease: inherited demyelinating neuropathies--from basic to clinical research. *Nature clinical practice Neurology* 2007;3:453-464.
- s7. Mo Z, Zhao X, Liu H, et al. Aberrant GlyRS-HDAC6 interaction linked to axonal transport deficits in Charcot-Marie-Tooth neuropathy. *Nature communications* 2018;9:1007.
- s8. Benoy V, Van Helleputte L, Prior R, et al. HDAC6 is a therapeutic target in mutant GARS-induced Charcot-Marie-Tooth disease. *Brain* 2018;141:673-687.
- s9. Helleputte L, Benoy V, Van Den Bosch L. The role of histone deacetylase 6 (HDAC6) in neurodegeneration. *Research and Reports in Biology* 2014;5:1-13.
- s10. He W, Bai G, Zhou H, et al. CMT2D neuropathy is linked to the neomorphic binding activity of glycyl-tRNA synthetase. *Nature* 2015;526:710-714.
- s11. Kalmar B, Innes A, Wanisch K, et al. Mitochondrial deficits and abnormal mitochondrial retrograde axonal transport play a role in the pathogenesis of mutant Hsp27-induced Charcot Marie Tooth Disease. *Human molecular genetics* 2017;26:3313-3326.
- s12. Rocha AG, Franco A, Krezel AM, et al. MFN2 agonists reverse mitochondrial defects in preclinical models of Charcot-Marie-Tooth disease type 2A. *Science* 2018;360:336-341.
- s13. Jia Li. MC, Rajasekhar NVS Suragani, R Scott Pearsall, and Ravindra, Kumar. ACE-083, A Locally-Acting GDF/Activin Ligand Trap, Augments Dorsiflexor Muscle Function in a

Murine Model of Charcot-Marie-Tooth (CMT) Disease. In: Peripheral Nerve Society Annual Meeting. Sitges, Spain, 2017.

s14. Rodino-Klapac LR, Haidet AM, Kota J, Handy C, Kaspar BK, Mendell JR. Inhibition of myostatin with emphasis on follistatin as a therapy for muscle disease. *Muscle & nerve* 2009;39:283-296.

s15. Pearsall R, Widrick J, Cotton E, et al. ACE-083 increases muscle hypertrophy and strength in C57BL/6 mice. *Neuromuscular Disorders*;25:S218.

s16. Mulivor A, Sako D, Cannell M, et al. G.P.205: A modified cysteine knot ligand trap of the TGF $\beta$  superfamily, ACE-083, increases muscle mass locally in a mouse model of Duchenne muscular dystrophy. *Neuromuscular Disorders* 2014;24:878.

s17. Pearsall RS WJ, Sako D, Davies M, et al. ACE-083, a Locally-Acting TGF- $\beta$  Superfamily Ligand Trap, Increases Muscle Mass and Strength in a Mouse Model of ALS. In: MDA Clinical Conference; 2016, 2016.

s18. Li J, Cannell M, Suragani R, Pearsall R, Kumar R. Ace-083, a Locally-Acting Gdf/Activin Ligand Trap, Augments Dorsiflexor Muscle Function in a Murine Model of Charcot-Marie-Tooth (Cmt) Disease. *Journal of the Peripheral Nervous System* 2017;22:331-331.

s19. Glasser CE, Gartner MR, Wilson D, Miller B, Sherman ML, Attie KM. Locally acting ACE-083 increases muscle volume in healthy volunteers. *Muscle & nerve* 2018;57:921-926.

s20. Russo FB, Cugola FR, Fernandes IR, Pignatari GC, Beltrao-Braga PC. Induced pluripotent stem cells for modeling neurological disorders. *World journal of transplantation* 2015;5:209-221.

s21. Saporta MA, Dang V, Volfson D, et al. Axonal Charcot-Marie-Tooth disease patient-derived motor neurons demonstrate disease-specific phenotypes including abnormal electrophysiological properties. *Experimental neurology* 2015;263:190-199.

s22. Juneja M, Azmi A, Baets J, et al. PFN2 and GAMT as common molecular determinants of axonal Charcot-Marie-Tooth disease. *Journal of neurology, neurosurgery, and psychiatry* 2018.

s23. Guo W, Naujock M, Fumagalli L, et al. HDAC6 inhibition reverses axonal transport defects in motor neurons derived from FUS-ALS patients. *Nature communications* 2017;8:861.

- s24. Amoroso MW, Croft GF, Williams DJ, et al. Accelerated high-yield generation of limb-innervating motor neurons from human stem cells. *The Journal of neuroscience* 2013;33:574-586.
- s25. Kim JY, Woo SY, Hong YB, et al. HDAC6 Inhibitors Rescued the Defective Axonal Mitochondrial Movement in Motor Neurons Derived from the Induced Pluripotent Stem Cells of Peripheral Neuropathy Patients with HSPB1 Mutation. *Stem cells international* 2016;2016:9475981.
- s26. Haidar M, Timmerman V. Autophagy as an Emerging Common Pathomechanism in Inherited Peripheral Neuropathies. *Frontiers in molecular neuroscience* 2017;10:143.
- s27. Kitani-Morii F, Imamura K, Kondo T, et al. iPSC model for demyelinating Charcot-Marie-Tooth disease. *Journal of the Neurological Sciences* 2017;381:856-857.
- s28. Shi L, Huang L, He R, et al. Modeling the Pathogenesis of Charcot-Marie-Tooth Disease Type 1A Using Patient-Specific iPSCs. *Stem cell reports* 2018;10:120-133.
- s29. Kitani-Morii F, Imamura K, Kondo T, et al. Analysis of neural crest cells from Charcot-Marie-Tooth disease patients demonstrates disease-relevant molecular signature. *Neuroreport* 2017;28:814-821.
- s30. Chambers SM, Fasano CA, Papapetrou EP, Tomishima M, Sadelain M, Studer L. Highly efficient neural conversion of human ES and iPS cells by dual inhibition of SMAD signaling. *Nature biotechnology* 2009;27:275-280.
- s31. Lee G, Chambers SM, Tomishima MJ, Studer L. Derivation of neural crest cells from human pluripotent stem cells. *Nature protocols* 2010;5:688-701.
- s32. Jiang X, Gwye Y, McKeown SJ, Bronner-Fraser M, Lutzko C, Lawlor ER. Isolation and characterization of neural crest stem cells derived from in vitro-differentiated human embryonic stem cells. *Stem cells and development* 2009;18:1059-1070.
- s33. Kreitzer FR, Salomonis N, Sheehan A, et al. A robust method to derive functional neural crest cells from human pluripotent stem cells. *American journal of stem cells* 2013;2:119-131.
- s34. Liu Q, Swistowski A, Zeng X. Human neural crest stem cells derived from human pluripotent stem cells. *Methods in molecular biology* 2014;1210:79-90.
- s35. Kim HS, Lee J, Lee DY, et al. Schwann Cell Precursors from Human Pluripotent Stem Cells as a Potential Therapeutic Target for Myelin Repair. *Stem cell reports* 2017;8:1714-1726.

- s36. Vigo T, Nobbio L, Hummelen PV, et al. Experimental Charcot-Marie-Tooth type 1A: a cDNA microarrays analysis. *Molecular and cellular neurosciences* 2005;28:703-714.
- s37. Duan RS, Jin T, Yang X, Mix E, Adem A, Zhu J. Apolipoprotein E deficiency enhances the antigen-presenting capacity of Schwann cells. *Glia* 2007;55:772-776.
- s38. Garcia-Mateo N, Ganfornina MD, Montero O, Gijon MA, Murphy RC, Sanchez D. Schwann cell-derived Apolipoprotein D controls the dynamics of post-injury myelin recognition and degradation. *Frontiers in cellular neuroscience* 2014;8:374.
- s39. Hunter M, Angelicheva D, Tournev I, et al. NDRG1 interacts with APO A-I and A-II and is a functional candidate for the HDL-C QTL on 8q24. *Biochemical and biophysical research communications* 2005;332:982-992.
- s40. LeBlanc AC, Foldvari M, Spencer DF, et al. The apolipoprotein A-I gene is actively expressed in the rapidly myelinating avian peripheral nerve. *The Journal of cell biology* 1989;109:1245-1256.
- s41. Poliani PL, Wang Y, Fontana E, et al. TREM2 sustains microglial expansion during aging and response to demyelination. *The Journal of clinical investigation* 2015;125:2161-2170.
- s42. Kohl B, Fischer S, Groh J, Wessig C, Martini R. MCP-1/CCL2 modifies axon properties in a PMP22-overexpressing mouse model for Charcot-Marie-tooth 1A neuropathy. *The American journal of pathology* 2010;176:1390-1399.
- s43. Adebola AA, Di Castri T, He CZ, et al. Neurofilament light polypeptide gene N98S mutation in mice leads to neurofilament network abnormalities and a Charcot-Marie-Tooth Type 2E phenotype. *Human molecular genetics* 2015;24:2163-2174.
- s44. Cartoni R, Arnaud E, Medard JJ, et al. Expression of mitofusin 2(R94Q) in a transgenic mouse leads to Charcot-Marie-Tooth neuropathy type 2A. *Brain* 2010;133:1460-1469.
- s45. Misko A, Jiang S, Wegorzewska I, Milbrandt J, Baloh RH. Mitofusin 2 is necessary for transport of axonal mitochondria and interacts with the Miro/Milton complex. *The Journal of neuroscience* 2010;30:4232-4240.
- s46. Bannerman P, Burns T, Xu J, Miers L, Pleasure D. Mice Hemizygous for a Pathogenic Mitofusin-2 Allele Exhibit Hind Limb/Foot Gait Deficits and Phenotypic Perturbations in Nerve and Muscle. *PloS one* 2016;11:e0167573.

- s47. Detmer SA, Vande Velde C, Cleveland DW, Chan DC. Hindlimb gait defects due to motor axon loss and reduced distal muscles in a transgenic mouse model of Charcot-Marie-Tooth type 2A. *Human molecular genetics* 2008;17:367-375.
- s48. Vallat JM, Sindou P, Preux PM, et al. Ultrastructural PMP22 expression in inherited demyelinating neuropathies. *Annals of neurology* 1996;39:813-817.
- s49. Li J, Bai Y, Ghandour K, et al. Skin biopsies in myelin-related neuropathies: bringing molecular pathology to the bedside. *Brain* 2005;128:1168-1177.
- s50. Bouhy D, Juneja M, Katona I, et al. A knock-in/knock-out mouse model of HSPB8-associated distal hereditary motor neuropathy and myopathy reveals toxic gain-of-function of mutant Hspb8. *Acta neuropathologica* 2018;135:131-148.
- s51. Sociali G, Visigalli D, Prukop T, et al. Tolerability and efficacy study of P2X7 inhibition in experimental Charcot-Marie-Tooth type 1A (CMT1A) neuropathy. *Neurobiology of disease* 2016;95:145-157.
- s52. Magyar JP, Martini R, Ruelicke T, et al. Impaired differentiation of Schwann cells in transgenic mice with increased PMP22 gene dosage. *The Journal of neuroscience* 1996;16:5351-5360.
- s53. Verhamme C, King RH, ten Asbroek AL, et al. Myelin and axon pathology in a long-term study of PMP22-overexpressing mice. *Journal of neuropathology and experimental neurology* 2011;70:386-398.
- s54. Robertson AM, Perea J, McGuigan A, et al. Comparison of a new pmp22 transgenic mouse line with other mouse models and human patients with CMT1A. *Journal of anatomy* 2002;200:377-390.
- s55. Perea J, Robertson A, Tolmachova T, et al. Induced myelination and demyelination in a conditional mouse model of Charcot-Marie-Tooth disease type 1A. *Human molecular genetics* 2001;10:1007-1018.
- s56. Giese KP, Martini R, Lemke G, Soriano P, Schachner M. Mouse P0 gene disruption leads to hypomyelination, abnormal expression of recognition molecules, and degeneration of myelin and axons. *Cell* 1992;71:565-576.
- s57. Runker AE, Kobsar I, Fink T, et al. Pathology of a mouse mutation in peripheral myelin protein P0 is characteristic of a severe and early onset form of human Charcot-Marie-Tooth type 1B disorder. *The Journal of cell biology* 2004;165:565-573.

- s58. Wrabetz L, D'Antonio M, Pennuto M, et al. Different intracellular pathomechanisms produce diverse Myelin Protein Zero neuropathies in transgenic mice. *The Journal of neuroscience* 2006;26:2358-2368.
- s59. Pennuto M, Tinelli E, Malaguti M, et al. Ablation of the UPR-mediator CHOP restores motor function and reduces demyelination in Charcot-Marie-Tooth 1B mice. *Neuron* 2008;57:393-405.
- s60. Saporta MA, Shy BR, Patzko A, et al. MpzR98C arrests Schwann cell development in a mouse model of early-onset Charcot-Marie-Tooth disease type 1B. *Brain* 2012;135:2032-2047.
- s61. Patzko A, Bai Y, Saporta MA, et al. Curcumin derivatives promote Schwann cell differentiation and improve neuropathy in R98C CMT1B mice. *Brain* 2012;135:3551-3566.
- s62. Zhu H, Guariglia S, Yu RY, et al. Mutation of SIMPLE in Charcot-Marie-Tooth 1C alters production of exosomes. *Molecular biology of the cell* 2013;24:1619-1637, S1611-1613.
- s63. Lee SM, Sha D, Mohammed AA, et al. Motor and sensory neuropathy due to myelin infolding and paranodal damage in a transgenic mouse model of Charcot-Marie-Tooth disease type 1C. *Human molecular genetics* 2013;22:1755-1770.
- s64. Schneider-Maunoury S, Topilko P, Seitandou T, et al. Disruption of Krox-20 results in alteration of rhombomeres 3 and 5 in the developing hindbrain. *Cell* 1993;75:1199-1214.
- s65. Nagarajan R, Svaren J, Le N, Araki T, Watson M, Milbrandt J. EGR2 mutations in inherited neuropathies dominant-negatively inhibit myelin gene expression. *Neuron* 2001;30:355-368.
- s66. Suter U, Moskow JJ, Welcher AA, et al. A leucine-to-proline mutation in the putative first transmembrane domain of the 22-kDa peripheral myelin protein in the trembler-J mouse. *Proceedings of the National Academy of Sciences of the United States of America* 1992;89:4382-4386.
- s67. Rangaraju S, Verrier JD, Madorsky I, Nicks J, Dunn WA, Jr., Notterpek L. Rapamycin activates autophagy and improves myelination in explant cultures from neuropathic mice. *The Journal of neuroscience* 2010;30:11388-11397.
- s68. Suh JG, Ichihara N, Saigoh K, et al. An in-frame deletion in peripheral myelin protein-22 gene causes hypomyelination and cell death of the Schwann cells in the new Trembler mutant mice. *Neuroscience* 1997;79:735-744.

- s69. Isaacs AM, Davies KE, Hunter AJ, et al. Identification of two new Pmp22 mouse mutants using large-scale mutagenesis and a novel rapid mapping strategy. *Human molecular genetics* 2000;9:1865-1871.
- s70. Strickland AV, Rebelo AP, Zhang F, et al. Characterization of the mitofusin 2 R94W mutation in a knock-in mouse model. *Journal of the peripheral nervous system* 2014;19:152-164.
- s71. Li J, Lancaster E, Yousaf A, et al. A Rat Model of Cmt2a Develops a Progressive Neuropathy. *Journal of the Peripheral Nervous System* 2017;22:331-331.
- s72. Zhao C, Takita J, Tanaka Y, et al. Charcot-Marie-Tooth disease type 2A caused by mutation in a microtubule motor KIF1Bbeta. *Cell* 2001;105:587-597.
- s73. Poitelon Y, Kozlov S, Devaux J, et al. Behavioral and molecular exploration of the AR-CMT2A mouse model Lmna (R298C/R298C). *Neuromolecular medicine* 2012;14:40-52.
- s74. Seburn KL, Nangle LA, Cox GA, Schimmel P, Burgess RW. An active dominant mutation of Glycyl-tRNA synthetase causes neuropathy in a Charcot-Marie-Tooth 2D mouse model. *Neuron* 2006;51:715-726.
- s75. Achilli F, Bros-Facer V, Williams HP, et al. An ENU-induced mutation in mouse glycyl-tRNA synthetase (GARS) causes peripheral sensory and motor phenotypes creating a model of Charcot-Marie-Tooth type 2D peripheral neuropathy. *Disease models & mechanisms* 2009;2:359-373.
- s76. Seo AJ, Shin YH, Lee SJ, et al. A novel adenoviral vector-mediated mouse model of Charcot-Marie-Tooth type 2D (CMT2D). *Journal of molecular histology* 2014;45:121-128.
- s77. Dequen F, Filali M, Lariviere RC, Perrot R, Hisanaga S, Julien JP. Reversal of neuropathy phenotypes in conditional mouse model of Charcot-Marie-Tooth disease type 2E. *Human molecular genetics* 2010;19:2616-2629.
- s78. Shen H, Barry DM, Dale JM, Garcia VB, Calcutt NA, Garcia ML. Muscle pathology without severe nerve pathology in a new mouse model of Charcot-Marie-Tooth disease type 2E. *Human molecular genetics* 2011;20:2535-2548.
- s79. Lee J, Jung SC, Joo J, et al. Overexpression of mutant HSP27 causes axonal neuropathy in mice. *Journal of biomedical science* 2015;22:43.
- s80. Bouhy D, Geuens T, De Winter V, et al. Characterization of New Transgenic Mouse Models for Two Charcot-Marie-Tooth-Causing HspB1 Mutations using the Rosa26 Locus. *Journal of neuromuscular diseases* 2016;3:183-200.

- s81. Srivastava AK, Rensch SR, Naiman NE, et al. Mutant HSPB1 overexpression in neurons is sufficient to cause age-related motor neuronopathy in mice. *Neurobiology of disease* 2012;47:163-173.
- s82. Barneo-Munoz M, Juarez P, Civera-Tregon A, et al. Lack of GDAP1 induces neuronal calcium and mitochondrial defects in a knockout mouse model of charcot-marie-tooth neuropathy. *PLoS genetics* 2015;11:e1005115.
- s83. Niemann A, Huber N, Wagner KM, et al. The Gdap1 knockout mouse mechanistically links redox control to Charcot-Marie-Tooth disease. *Brain* 2014;137:668-682.
- s84. Sanbe A, Marunouchi T, Abe T, et al. Phenotype of cardiomyopathy in cardiac-specific heat shock protein B8 K141N transgenic mouse. *The Journal of biological chemistry* 2013;288:8910-8921.
- s85. Zhang R, Zhang F, Li X, et al. A novel transgenic mouse model of Chinese Charcot-Marie-Tooth disease type 2L. *Neural regeneration research* 2014;9:413-419.
- s86. Sabblah TT, Nandini S, Ledray AP, et al. A novel mouse model carrying a human cytoplasmic dynein mutation shows motor behavior deficits consistent with Charcot-Marie-Tooth type 2O disease. *Scientific reports* 2018;8:1739.
- s87. Bogdanik LP, Sleight JN, Tian C, et al. Loss of the E3 ubiquitin ligase LRSAM1 sensitizes peripheral axons to degeneration in a mouse model of Charcot-Marie-Tooth disease. *Disease models & mechanisms* 2013;6:780-792.
- s88. M. Gu WG, C. Luan, Z. Yu, Y. Chen, S. Dang, Y. Kuang, Z. Wang. The Dhtkd1 Tyr486\* knock-in mouse model recapitulates some phenotypes of Charcot-Marie-Tooth disease type 2Q. In: ASHG. Boston, 2013.
- s89. Balastik M, Ferraguti F, Pires-da Silva A, et al. Deficiency in ubiquitin ligase TRIM2 causes accumulation of neurofilament light chain and neurodegeneration. *Proceedings of the National Academy of Sciences of the United States of America* 2008;105:12016-12021.
- s90. Lee S, Panthi S, Jo HW, et al. Anatomical distributional defects in mutant genes associated with dominant intermediate Charcot-Marie-Tooth disease type C in an adenovirus-mediated mouse model. *Neural regeneration research* 2017;12:486-492.
- s91. Mones S, Bordignon B, Fontes M. Connexin 32 is involved in mitosis. *Glia* 2012;60:457-464.



- s92. Mones S, Bordignon B, Peiretti F, et al. CamKII inhibitors reduce mitotic instability, connexon anomalies and progression of the in vivo behavioral phenotype in transgenic animals expressing a mutated Gjb1 gene. *Frontiers in neuroscience* 2014;8:151.
- s93. Nelles E, Butzler C, Jung D, et al. Defective propagation of signals generated by sympathetic nerve stimulation in the liver of connexin32-deficient mice. *Proceedings of the National Academy of Sciences of the United States of America* 1996;93:9565-9570.
- s94. Kagiava A, Sargiannidou I, Theophilidis G, et al. Intrathecal gene therapy rescues a model of demyelinating peripheral neuropathy. *Proceedings of the National Academy of Sciences of the United States of America* 2016;113:E2421-2429.
- s95. Son D, Kang PJ, Yun W, You S. Generation of induced pluripotent stem cell (iPSC) line from a 36-year-old Charcot-Marie-Tooth disease patient with GJB1 mutation (CMTX). *Stem cell research* 2017;21:9-12.
- s96. Sainio MT, Ylikallio E, Mäenpää L, et al. Absence of NEFL in patient-specific neurons in early-onset Charcot-Marie-Tooth neuropathy. *Neurology Genetics* 2018;doi:10.1212/NXG.0000000000000244.