

Supplementary Table 4. Characteristic of models for patients with diabetes

Study sample	Sample size	Outcomes	Follow-up	Baseline age	Number of events	Model	Sensitivity (%)	Specificity (%)	PPV (%)	NPV (%)	Accuracy (%)	AUC
Type 2 diabetes patients from the Kaiser Permanente Northern California (KPNC) Diabetes Registry, Oakland, CA, USA ¹	29,961	dementia	10 years	≥60	5173	age, education, microvascular disease, diabetic foot, cerebrovascular disease, cardiovascular disease, acute metabolic event, depression						0.74
Patients with type 2 diabetes from pathways study cohort from Washington, USA	2,413	dementia	mean 6.7 years	≥60	328	age, education, microvascular disease, diabetic foot, cerebrovascular disease, cardiovascular disease, acute metabolic event, depression						0.75
Clinical Practice Research Datalink	133,176	dementia		≥60 years		Age, Male, Myocardial infarction, Congestive heart failure, Coronary and peripheral vascular disease, Cerebrovascular disease, Chronic pulmonary disease, Rheumatologic						0.86

Supplementary Table 4. Characteristic of models for patients with diabetes

Study sample	Sample size	Outcomes	Follow-up	Baseline age	Number of events	Model	Sensitivity (%)	Specificity (%)	PPV (%)	NPV (%)	Accuracy (%)	AUC
(CPRD) 2						disease, Peptic ulcer disease, Renal disease/end stage renal disease, Mild liver disease/moderate or severe liver disease, Any malignancy including lymphoma and leukemia/metastatic solid tumor, Epilepsy, Hyperlipidemia, Parkinson's disease, Cardiac disease ASCVD, Glaucoma, Transplantation, Thyroid disorder, Gout, Crohn's and ulcerative disease, Pain and inflammation / pain, Depression, Psychotic illness, Bipolar disorders, Anxiety and tension						
National Diabetes Care Management Program (NDCMP) ³	18,360	dementia	mean 8.09 years	50–94 years	853	age, sex, duration of type 2 diabetes, BMI, FPG, HbA1c, stroke, postural hypertension, hypoglycemia, coronary artery disease, anti-diabetes medications (3 years)						0.82
						age, sex, duration of type 2 diabetes, BMI, FPG, HbA1c, stroke, postural hypertension, hypoglycemia, coronary artery disease, anti-diabetes medications (5 years)						0.79
						age, sex, duration of type 2 diabetes, BMI, FPG, HbA1c, stroke, postural hypertension, hypoglycemia, coronary artery disease, anti-diabetes medications (10 years)						0.76
	9,180	dementia	mean 8.09 years	50–94 years	375	age, sex, duration of type 2 diabetes, BMI, FPG, HbA1c, stroke, postural hypertension, hypoglycemia, coronary artery disease, anti-diabetes medications (3 years)						0.84

Supplementary Table 4. Characteristic of models for patients with diabetes

Study sample	Sample size	Outcomes	Follow-up	Baseline age	Number of events	Model	Sensitivity (%)	Specificity (%)	PPV (%)	NPV (%)	Accuracy (%)	AUC
						age, sex, duration of type 2 diabetes, BMI, FPG, HbA1c, stroke, postural hypertension, hypoglycemia, coronary artery disease, anti-diabetes medications (5 years)						0.80
						age, sex, duration of type 2 diabetes, BMI, FPG, HbA1c, stroke, postural hypertension, hypoglycemia, coronary artery disease, anti-diabetes medications (10 years)						0.75

Abbreviations: ASCVD: arteriosclerotic cardiovascular disease; BMI: body mass index; FPG: Fasting Plasma Glucose.

1. Exalto LG, Biessels GJ, Karter AJ, *et al.* Risk score for prediction of 10 year dementia risk in individuals with type 2 diabetes: A cohort study. *The Lancet Diabetes and Endocrinology* 2013; 1(3): 183-90.
2. Mehta HB, Mehta V, Tsai CL, *et al.* Development and Validation of the RxDx-Dementia Risk Index to Predict Dementia in Patients with Type 2 Diabetes and Hypertension. *J Alzheimers Dis* 2016; 49(2): 423-32.
3. Li CI, Li TC, Liu CS, *et al.* Risk score prediction model for dementia in patients with type 2 diabetes. *Eur J Neurol* 2018.