

SUPPLEMENTAL REFERENCES

60. Kahana MJ, Seelig D, Madsen JR. Theta returns. *Curr Opin Neurobiol.* 2001;11:739–744.
61. Burgess N, Maguire EA, O'Keefe J. The human hippocampus and spatial and episodic memory. *Neuron.* 2002;35:625–641.
62. Buzsáki G. Theta oscillations in the hippocampus. *Neuron.* 2002;33:325–340.
63. Hasselmo ME, Bodelón C, Wyble BP. A proposed function for hippocampal theta rhythm: separate phases of encoding and retrieval enhance reversal of prior learning. *Neural Comput.* 2002;14:793–817.
64. Kirk IJ, Mackay JC. The role of theta-range oscillations in synchronising and integrating activity in distributed mnemonic networks. *Cortex J Devoted Study Nerv Syst Behav.* 2003;39:993–1008.
65. Buzsáki G. Theta rhythm of navigation: link between path integration and landmark navigation, episodic and semantic memory. *Hippocampus.* 2005;15:827–840.
66. Hasselmo ME. What is the function of hippocampal theta rhythm?--Linking behavioral data to phasic properties of field potential and unit recording data. *Hippocampus.* 2005;15:936–949.
67. Stewart M, Fox SE. Do septal neurons pace the hippocampal theta rhythm? *Trends Neurosci.* 1990;13:163–168.
68. Bland BH, Oddie SD. Theta band oscillation and synchrony in the hippocampal formation and associated structures: the case for its role in sensorimotor integration. *Behav Brain Res.* 2001;127:119–136.
69. Petsche H, Gogolak G, Vanzwieten PA. RHYTHMICITY OF SEPTAL CELL DISCHARGES AT VARIOUS LEVELS OF RETICULAR EXCITATION. *Electroencephalogr Clin Neurophysiol.* 1965;19:25–33.
70. Gogolák G, Stumpf C, Petsche H, Sterc J. The firing pattern of septal neurons and the form of the hippocampal theta wave. *Brain Res.* 1968;7:201–207.
71. Lamour Y, Dutar P, Jobert A. Septo-hippocampal and other medial septum-diagonal band neurons: electrophysiological and pharmacological properties. *Brain Res.* 1984;309:227–239.
72. Brazhnik ES, Vinogradova OS. Control of the neuronal rhythmic bursts in the septal pacemaker of theta-rhythm: effects of anaesthetic and anticholinergic drugs. *Brain Res.* 1986;380:94–106.
73. Vinogradova OS. Expression, control, and probable functional significance of the neuronal theta-rhythm. *Prog Neurobiol.* 1995;45:523–583.

74. Brazhnik ES, Fox SE. Intracellular recordings from medial septal neurons during hippocampal theta rhythm. *Exp Brain Res*. 1997;114:442–453.
75. Dragoi G, Carpi D, Recce M, Csicsvari J, Buzsáki G. Interactions between hippocampus and medial septum during sharp waves and theta oscillation in the behaving rat. *J Neurosci Off J Soc Neurosci*. 1999;19:6191–6199.
76. Leung LS, Shen B. Glutamatergic synaptic transmission participates in generating the hippocampal EEG. *Hippocampus*. 2004;14:510–525.
77. Koenig J, Linder AN, Leutgeb JK, Leutgeb S. The spatial periodicity of grid cells is not sustained during reduced theta oscillations. *Science*. 2011;332:592–595.
78. Freund TF, Gulyás AI. Inhibitory control of GABAergic interneurons in the hippocampus. *Can J Physiol Pharmacol*. 1997;75:479–487.
79. Brandon MP, Bogaard AR, Libby CP, Connerney MA, Gupta K, Hasselmo ME. Reduction of theta rhythm dissociates grid cell spatial periodicity from directional tuning. *Science*. 2011;332:595–599.
80. Mizumori SJ, Perez GM, Alvarado MC, Barnes CA, McNaughton BL. Reversible inactivation of the medial septum differentially affects two forms of learning in rats. *Brain Res*. 1990;528:12–20.
81. M'Harzi M, Jarrard LE. Effects of medial and lateral septal lesions on acquisition of a place and cue radial maze task. *Behav Brain Res*. 1992;49:159–165.
82. Asaka Y, Griffin AL, Berry SD. Reversible septal inactivation disrupts hippocampal slow-wave and unit activity and impairs trace conditioning in rabbits (*Oryctolagus cuniculus*). *Behav Neurosci*. 2002;116:434–442.
83. Wang Y, Romani S, Lustig B, Leonardo A, Pastalkova E. Theta sequences are essential for internally generated hippocampal firing fields. *Nat Neurosci*. 2015;18:282–288.
84. Tóth K, Borhegyi Z, Freund TF. Postsynaptic targets of GABAergic hippocampal neurons in the medial septum-diagonal band of Broca complex. *J Neurosci Off J Soc Neurosci*. 1993;13:3712–3724.
85. Wang X-J. Pacemaker neurons for the theta rhythm and their synchronization in the septohippocampal reciprocal loop. *J Neurophysiol*. 2002;87:889–900.
86. Gaykema RP, van der Kuil J, Hersh LB, Luiten PG. Patterns of direct projections from the hippocampus to the medial septum-diagonal band complex: anterograde tracing with Phaseolus vulgaris leucoagglutinin combined with immunohistochemistry of choline acetyltransferase. *Neuroscience*. 1991;43:349–360.
87. Hasselmo ME, Schnell E. Laminar selectivity of the cholinergic suppression of synaptic transmission in rat hippocampal region CA1: computational modeling and brain slice physiology. *J Neurosci Off J Soc Neurosci*. 1994;14:3898–3914.

88. Rokers B, Mercado E, Allen MT, Myers CE, Gluck MA. A connectionist model of septohippocampal dynamics during conditioning: closing the loop. *Behav Neurosci*. 2002;116:48–62.
89. Manseau F, Goutagny R, Danik M, Williams S. The hippocamptoseptal pathway generates rhythmic firing of GABAergic neurons in the medial septum and diagonal bands: an investigation using a complete septohippocampal preparation in vitro. *J Neurosci Off J Soc Neurosci*. 2008;28:4096–4107.
90. Hao S, Tang B, Wu Z, Ure K, Sun Y, Tao H, Gao Y, Patel AJ, Curry DJ, Samaco RC, Zoghbi HY, Tang J. Forniceal deep brain stimulation rescues hippocampal memory in Rett syndrome mice. *Nature*. 2015;526:430–434.
91. Pohodich AE, Yalamanchili H, Raman AT, Wan Y-W, Gundry M, Hao S, Jin H, Tang J, Liu Z, Zoghbi HY. Forniceal deep brain stimulation induces gene expression and splicing changes that promote neurogenesis and plasticity. *eLife*. 2018;7. doi:10.7554/eLife.34031.
92. McNaughton BL, Barnes CA. Physiological identification and analysis of dentate granule cell responses to stimulation of the medial and lateral perforant pathways in the rat. *J Comp Neurol*. 1977;175:439–454.
93. McNaughton BL. Evidence for two physiologically distinct perforant pathways to the fascia dentata. *Brain Res*. 1980;199:1–19.
94. Kramis R, Vanderwolf CH. Frequency-specific RSA-like hippocampal patterns elicited by septal, hypothalamic, and brain stem electrical stimulation. *Brain Res*. 1980;192:383–398.
95. McNaughton N, Ruan M, Woodnorth M-A. Restoring theta-like rhythmicity in rats restores initial learning in the Morris water maze. *Hippocampus*. 2006;16:1102–1110.
96. Scarlett D, Dypvik AT, Bland BH. Comparison of spontaneous and septally driven hippocampal theta field and theta-related cellular activity. *Hippocampus*. 2004;14:99–106.
97. Stumpf C. THE FAST COMPONENT IN THE ELECTRICAL ACTIVITY OF RABBIT'S HIPPOCAMPUS. *Electroencephalogr Clin Neurophysiol*. 1965;18:477–486.
98. Ball GG, Gray JA. Septal self-stimulation and hippocampal activity. *Physiol Behav*. 1971;6:547–549.
99. Wishart TB, Bland BH, Vanderwolf CH, Altman JL. Electroencephalographic correlates of behaviors elicited by electrical stimulation of the septum: seizure induced feeding. *Behav Biol*. 1973;9:763–769.
100. Bland BH. The Medial Septum: Node of the Ascending Brainstem Hippocampal Synchronizing Pathways. In: *The Behavioral Neuroscience of the Septal Region*. Springer, New York, NY; 2000: 115–145.
101. Defrance JF, Stanley JC, Marchand JE, Chronister RB. Cholinergic mechanisms and short-term potentiation. *Ciba Found Symp*. 1977;:109–126.

102. Lee DJ, Gurkoff GG, Izadi A, Seidl SE, Echeverri A, Melnik M, Berman RF, Ekstrom AD, Muizelaar JP, Lyeth BG, Shahlaie K. Septohippocampal Neuromodulation Improves Cognition after Traumatic Brain Injury. *J Neurotrauma*. 2015;32:1822–1832.
103. Lee DJ, Gurkoff GG, Izadi A, Berman RF, Ekstrom AD, Muizelaar JP, Lyeth BG, Shahlaie K. Medial septal nucleus theta frequency deep brain stimulation improves spatial working memory after traumatic brain injury. *J Neurotrauma*. 2013;30:131–139.
104. Kahana MJ. The cognitive correlates of human brain oscillations. *J Neurosci Off J Soc Neurosci*. 2006;26:1669–1672.
105. Lisman J, Buzsáki G. A neural coding scheme formed by the combined function of gamma and theta oscillations. *Schizophr Bull*. 2008;34:974–980.
106. Lisman JE, Idiart MA. Storage of 7 +/- 2 short-term memories in oscillatory subcycles. *Science*. 1995;267:1512–1515.
107. Jensen O. Maintenance of multiple working memory items by temporal segmentation. *Neuroscience*. 2006;139:237–249.
108. Bikbaev A, Manahan-Vaughan D. Relationship of hippocampal theta and gamma oscillations to potentiation of synaptic transmission. *Front Neurosci*. 2008;2:56–63.
109. Mehta MR, Lee AK, Wilson MA. Role of experience and oscillations in transforming a rate code into a temporal code. *Nature*. 2002;417:741–746.
110. Shirvalkar PR, Rapp PR, Shapiro ML. Bidirectional changes to hippocampal theta-gamma comodulation predict memory for recent spatial episodes. *Proc Natl Acad Sci U S A*. 2010;107:7054–7059.
111. Buño W, Garcia-Sanchez JL, Garcia-Austt E. Reset of hippocampal rhythmical activities by afferent stimulation. *Brain Res Bull*. 1978;3:21–28.
112. García-Sánchez JL, Buño W, Fuentes J, García-Austt E. Non-rhythmical hippocampal units, theta rhythm and afferent stimulation. *Brain Res Bull*. 1978;3:213–219.
113. Williams JM, Givens B. Stimulation-induced reset of hippocampal theta in the freely performing rat. *Hippocampus*. 2003;13:109–116.
114. Bland BH, Bird J, Jackson J, Natsume K. Medial septal modulation of the ascending brainstem hippocampal synchronizing pathways in the freely moving rat. *Hippocampus*. 2006;16:11–19.
115. Mamad O, McNamara HM, Reilly RB, Tsanov M. Medial septum regulates the hippocampal spatial representation. *Front Behav Neurosci*. 2015;9:166.
116. Larson J, Lynch G. Induction of synaptic potentiation in hippocampus by patterned stimulation involves two events. *Science*. 1986;232:985–988.
117. Diamond DM, Rose GM. Stress impairs LTP and hippocampal-dependent memory. *Ann N Y Acad Sci*. 1994;746:411–414.

118. Adey WR. Hippocampal states and functional relations with corticosubcortical systems in attention and learning. *Prog Brain Res.* 1967;27:228–245.
119. Givens B. Stimulus-evoked resetting of the dentate theta rhythm: relation to working memory. *Neuroreport.* 1996;8:159–163.
120. Vinogradova OS, Brazhnik ES, Kichigina VF, Stafekhina VS. Modulation of the reaction of hippocampal neurons to sensory stimuli by cholinergic substances. *Neurosci Behav Physiol.* 1996;26:113–124.
121. Weiler HT, Hasenöhrl RU, van Landeghem AA, van Landeghem M, Brankack J, Huston JP, Haas HL. Differential modulation of hippocampal signal transfer by tuberomammillary nucleus stimulation in freely moving rats dependent on behavioral state. *Synapse NY N.* 1998;28:294–301.
122. Krnjević K, Ropert N. Electrophysiological and pharmacological characteristics of facilitation of hippocampal population spikes by stimulation of the medial septum. *Neuroscience.* 1982;7:2165–2183.
123. Kiss J, Patel AJ, Freund TF. Distribution of septohippocampal neurons containing parvalbumin or choline acetyltransferase in the rat brain. *J Comp Neurol.* 1990;298:362–372.
124. Lewis PR, Shute CC. The cholinergic limbic system: projections to hippocampal formation, medial cortex, nuclei of the ascending cholinergic reticular system, and the subfornical organ and supra-optic crest. *Brain J Neurol.* 1967;90:521–540.
125. Köhler C, Chan-Palay V, Wu JY. Septal neurons containing glutamic acid decarboxylase immunoreactivity project to the hippocampal region in the rat brain. *Anat Embryol (Berl).* 1984;169:41–44.
126. Griffith WH, Matthews RT. Electrophysiology of AChE-positive neurons in basal forebrain slices. *Neurosci Lett.* 1986;71:169–174.
127. Freund TF, Antal M. GABA-containing neurons in the septum control inhibitory interneurons in the hippocampus. *Nature.* 1988;336:170–173.
128. Markram H, Segal M. Long-lasting facilitation of excitatory postsynaptic potentials in the rat hippocampus by acetylcholine. *J Physiol.* 1990;427:381–393.
129. Pepeu G, Blandina P. The acetylcholine, GABA, glutamate triangle in the rat forebrain. *J Physiol Paris.* 1998;92:351–355.
130. Jones GA, Norris SK, Henderson Z. Conduction velocities and membrane properties of different classes of rat septohippocampal neurons recorded in vitro. *J Physiol.* 1999;517 (Pt 3):867–877.
131. Manns ID, Mainville L, Jones BE. Evidence for glutamate, in addition to acetylcholine and GABA, neurotransmitter synthesis in basal forebrain neurons projecting to the entorhinal cortex. *Neuroscience.* 2001;107:249–263.

132. Colom LV, Castaneda MT, Reyna T, Hernandez S, Garrido-Sanabria E. Characterization of medial septal glutamatergic neurons and their projection to the hippocampus. *Synap N Y N*. 2005;58:151–164.
133. Gritti I, Henny P, Galloni F, Mainville L, Mariotti M, Jones BE. Stereological estimates of the basal forebrain cell population in the rat, including neurons containing choline acetyltransferase, glutamic acid decarboxylase or phosphate-activated glutaminase and colocalizing vesicular glutamate transporters. *Neuroscience*. 2006;143:1051–1064.
134. Lecourtier L, de Vasconcelos AP, Leroux E, Cosquer B, Geiger K, Lithfous S, Cassel J-C. Septohippocampal pathways contribute to system consolidation of a spatial memory: sequential implication of GABAergic and cholinergic neurons. *Hippocampus*. 2011;21:1277–1289.
135. Sotty F, Danik M, Manseau F, Laplante F, Quirion R, Williams S. Distinct electrophysiological properties of glutamatergic, cholinergic and GABAergic rat septohippocampal neurons: novel implications for hippocampal rhythmicity. *J Physiol*. 2003;551:927–943.
136. Simon AP, Poindessous-Jazat F, Dutar P, Epelbaum J, Bassant M-H. Firing properties of anatomically identified neurons in the medial septum of anesthetized and unanesthetized restrained rats. *J Neurosci Off J Soc Neurosci*. 2006;26:9038–9046.
137. Huh CYL, Goutagny R, Williams S. Glutamatergic neurons of the mouse medial septum and diagonal band of Broca synaptically drive hippocampal pyramidal cells: relevance for hippocampal theta rhythm. *J Neurosci Off J Soc Neurosci*. 2010;30:15951–15961.
138. Morris NP, Harris SJ, Henderson Z. Parvalbumin-immunoreactive, fast-spiking neurons in the medial septum/diagonal band complex of the rat: intracellular recordings in vitro. *Neuroscience*. 1999;92:589–600.
139. Leranath C, Carpi D, Buzsaki G, Kiss J. The entorhino-septo-supramammillary nucleus connection in the rat: morphological basis of a feedback mechanism regulating hippocampal theta rhythm. *Neuroscience*. 1999;88:701–718.
140. Hajszan T, Alreja M, Leranath C. Intrinsic vesicular glutamate transporter 2-immunoreactive input to septohippocampal parvalbumin-containing neurons: novel glutamatergic local circuit cells. *Hippocampus*. 2004;14:499–509.
141. Manseau F, Danik M, Williams S. A functional glutamatergic neurone network in the medial septum and diagonal band area. *J Physiol*. 2005;566:865–884.
142. Peterson GM, Williams LR, Varon S, Gage FH. Loss of GABAergic neurons in medial septum after fimbria-fornix transection. *Neurosci Lett*. 1987;76:140–144.
143. Papp EC, Hajos N, Acsády L, Freund TF. Medial septal and median raphe innervation of vasoactive intestinal polypeptide-containing interneurons in the hippocampus. *Neuroscience*. 1999;90:369–382.
144. Gulyás AI, Görcs TJ, Freund TF. Innervation of different peptide-containing neurons in the hippocampus by GABAergic septal afferents. *Neuroscience*. 1990;37:31–44.

145. Acsády L, Halasy K, Freund TF. Calretinin is present in non-pyramidal cells of the rat hippocampus--III. Their inputs from the median raphe and medial septal nuclei. *Neuroscience*. 1993;52:829–841.
146. Takács VT, Freund TF, Gulyás AI. Types and synaptic connections of hippocampal inhibitory neurons reciprocally connected with the medial septum. *Eur J Neurosci*. 2008;28:148–164.
147. Yamano M, Luiten PG. Direct synaptic contacts of medial septal efferents with somatostatin immunoreactive neurons in the rat hippocampus. *Brain Res Bull*. 1989;22:993–1001.
148. Colom LV, Bland BH. Medial septal cell interactions in relation to hippocampal field activity and the effects of atropine. *Hippocampus*. 1991;1:15–30.
149. Smythe JW, Colom LV, Bland BH. The extrinsic modulation of hippocampal theta depends on the coactivation of cholinergic and GABA-ergic medial septal inputs. *Neurosci Biobehav Rev*. 1992;16:289–308.
150. Brazhnik ES, Fox SE. Action potentials and relations to the theta rhythm of medial septal neurons in vivo. *Exp Brain Res*. 1999;127:244–258.
151. Yoder RM, Pang KCH. Involvement of GABAergic and cholinergic medial septal neurons in hippocampal theta rhythm. *Hippocampus*. 2005;15:381–392.
152. Monmaur P, Collet A, Puma C, Frankel-Kohn L, Sharif A. Relations between acetylcholine release and electrophysiological characteristics of theta rhythm: a microdialysis study in the urethane-anesthetized rat hippocampus. *Brain Res Bull*. 1997;42:141–146.
153. Xu C, Datta S, Wu M, Alreja M. Hippocampal theta rhythm is reduced by suppression of the H-current in septohippocampal GABAergic neurons. *Eur J Neurosci*. 2004;19:2299–2309.
154. Borhegyi Z, Varga V, Szilágyi N, Fabo D, Freund TF. Phase segregation of medial septal GABAergic neurons during hippocampal theta activity. *J Neurosci Off J Soc Neurosci*. 2004;24:8470–8479.
155. Bassant M-H, Simon A, Poindessous-Jazat F, Csaba Z, Epelbaum J, Dournaud P. Medial septal GABAergic neurons express the somatostatin sst2A receptor: functional consequences on unit firing and hippocampal theta. *J Neurosci Off J Soc Neurosci*. 2005;25:2032–2041.
156. Varga V, Hangya B, Kránitz K, Ludányi A, Zemankovics R, Katona I, Shigemoto R, Freund TF, Borhegyi Z. The presence of pacemaker HCN channels identifies theta rhythmic GABAergic neurons in the medial septum. *J Physiol*. 2008;586:3893–3915.
157. Kermer P, Naumann T, Bender R, Frotscher M. Fate of GABAergic septohippocampal neurons after fimbria-fornix transection as revealed by in situ hybridization for glutamate decarboxylase mRNA and parvalbumin immunocytochemistry. *J Comp Neurol*. 1995;362:385–399.

158. Hangya B, Borhegyi Z, Szilágyi N, Freund TF, Varga V. GABAergic neurons of the medial septum lead the hippocampal network during theta activity. *J Neurosci Off J Soc Neurosci*. 2009;29:8094–8102.
159. Tóth K, Freund TF, Miles R. Disinhibition of rat hippocampal pyramidal cells by GABAergic afferents from the septum. *J Physiol*. 1997;500 (Pt 2):463–474.
160. Kaifosh P, Lovett-Barron M, Turi GF, Reardon TR, Losonczy A. Septo-hippocampal GABAergic signaling across multiple modalities in awake mice. *Nat Neurosci*. 2013;16:1182–1184.
161. Talley EM, Cribbs LL, Lee JH, Daud A, Perez-Reyes E, Bayliss DA. Differential distribution of three members of a gene family encoding low voltage-activated (T-type) calcium channels. *J Neurosci Off J Soc Neurosci*. 1999;19:1895–1911.
162. Gangadharan G, Shin J, Kim S-W, Kim A, Paydar A, Kim D-S, Miyazaki T, Watanabe M, Yanagawa Y, Kim J, Kim Y-S, Kim D, Shin H-S. Medial septal GABAergic projection neurons promote object exploration behavior and type 2 theta rhythm. *Proc Natl Acad Sci U S A*. 2016;113:6550–6555.
163. Milner TA, Loy R, Amaral DG. An anatomical study of the development of the septo-hippocampal projection in the rat. *Brain Res*. 1983;284:343–371.
164. Amaral DG, Kurz J. An analysis of the origins of the cholinergic and noncholinergic septal projections to the hippocampal formation of the rat. *J Comp Neurol*. 1985;240:37–59.
165. Houser CR, Crawford GD, Barber RP, Salvaterra PM, Vaughn JE. Organization and morphological characteristics of cholinergic neurons: an immunocytochemical study with a monoclonal antibody to choline acetyltransferase. *Brain Res*. 1983;266:97–119.
166. Frotscher M, Léránth C. Cholinergic innervation of the rat hippocampus as revealed by choline acetyltransferase immunocytochemistry: a combined light and electron microscopic study. *J Comp Neurol*. 1985;239:237–246.
167. Matthews DA, Salvaterra PM, Crawford GD, Houser CR, Vaughn JE. An immunocytochemical study of choline acetyltransferase-containing neurons and axon terminals in normal and partially deafferented hippocampal formation. *Brain Res*. 1987;402:30–43.
168. Wainer BH, Bolam JP, Freund TF, Henderson Z, Totterdell S, Smith AD. Cholinergic synapses in the rat brain: a correlated light and electron microscopic immunohistochemical study employing a monoclonal antibody against choline acetyltransferase. *Brain Res*. 1984;308:69–76.
169. Léránth C, Frotscher M. GABAergic input of cholecystokinin-immunoreactive neurons in the hilar region of the rat hippocampus. An electron microscopic double immunostaining study. *Histochemistry*. 1987;86:287–290.
170. Cobb SR, Davies CH. Cholinergic modulation of hippocampal cells and circuits. *J Physiol*. 2005;562:81–88.

171. Nyakas C, Luiten PG, Spencer DG, Traber J. Detailed projection patterns of septal and diagonal band efferents to the hippocampus in the rat with emphasis on innervation of CA1 and dentate gyrus. *Brain Res Bull.* 1987;18:533–545.
172. Descarries L, Gisiger V, Steriade M. Diffuse transmission by acetylcholine in the CNS. *Prog Neurobiol.* 1997;53:603–625.
173. Vizi ES, Kiss JP. Neurochemistry and pharmacology of the major hippocampal transmitter systems: synaptic and nonsynaptic interactions. *Hippocampus.* 1998;8:566–607.
174. Descarries L. The hypothesis of an ambient level of acetylcholine in the central nervous system. *J Physiol Paris.* 1998;92:215–220.
175. Zoli M, Picciotto MR, Ferrari R, Cocchi D, Changeux JP. Increased neurodegeneration during ageing in mice lacking high-affinity nicotine receptors. *EMBO J.* 1999;18:1235–1244.
176. Fabian-Fine R, Skehel P, Errington ML, Davies HA, Sher E, Stewart MG, Fine A. Ultrastructural distribution of the alpha7 nicotinic acetylcholine receptor subunit in rat hippocampus. *J Neurosci Off J Soc Neurosci.* 2001;21:7993–8003.
177. Hill JA, Zoli M, Bourgeois JP, Changeux JP. Immunocytochemical localization of a neuronal nicotinic receptor: the beta 2-subunit. *J Neurosci Off J Soc Neurosci.* 1993;13:1551–1568.
178. Alkondon M, Pereira EF, Eisenberg HM, Albuquerque EX. Choline and selective antagonists identify two subtypes of nicotinic acetylcholine receptors that modulate GABA release from CA1 interneurons in rat hippocampal slices. *J Neurosci Off J Soc Neurosci.* 1999;19:2693–2705.
179. Alkondon M, Albuquerque EX. Nicotinic acetylcholine receptor alpha7 and alpha4beta2 subtypes differentially control GABAergic input to CA1 neurons in rat hippocampus. *J Neurophysiol.* 2001;86:3043–3055.
180. Graham AJ, Ray MA, Perry EK, Jaros E, Perry RH, Volsen SG, Bose S, Evans N, Lindstrom J, Court JA. Differential nicotinic acetylcholine receptor subunit expression in the human hippocampus. *J Chem Neuroanat.* 2003;25:97–113.
181. Bell KA, Shim H, Chen C-K, McQuiston AR. Nicotinic excitatory postsynaptic potentials in hippocampal CA1 interneurons are predominantly mediated by nicotinic receptors that contain $\alpha 4$ and $\beta 2$ subunits. *Neuropharmacology.* 2011;61:1379–1388.
182. Tang A-H, Karson MA, Nagode DA, McIntosh JM, Uebele VN, Renger JJ, Klugmann M, Milner TA, Alger BE. Nerve terminal nicotinic acetylcholine receptors initiate quantal GABA release from perisomatic interneurons by activating axonal T-type (Cav3) Ca^{2+} channels and Ca^{2+} release from stores. *J Neurosci Off J Soc Neurosci.* 2011;31:13546–13561.
183. Yamasaki M, Matsui M, Watanabe M. Preferential localization of muscarinic M1 receptor on dendritic shaft and spine of cortical pyramidal cells and its anatomical evidence for volume transmission. *J Neurosci Off J Soc Neurosci.* 2010;30:4408–4418.

184. Cea-del Rio CA, Lawrence JJ, Tricoire L, Erdelyi F, Szabo G, McBain CJ. M3 muscarinic acetylcholine receptor expression confers differential cholinergic modulation to neurochemically distinct hippocampal basket cell subtypes. *J Neurosci Off J Soc Neurosci*. 2010;30:6011–6024.
185. Cea-del Rio CA, Lawrence JJ, Erdelyi F, Szabo G, McBain CJ. Cholinergic modulation amplifies the intrinsic oscillatory properties of CA1 hippocampal cholecystokinin-positive interneurons. *J Physiol*. 2011;589:609–627.
186. Raiteri M, Leardi R, Marchi M. Heterogeneity of presynaptic muscarinic receptors regulating neurotransmitter release in the rat brain. *J Pharmacol Exp Ther*. 1984;228:209–214.
187. Levey AI, Edmunds SM, Koliatsos V, Wiley RG, Heilman CJ. Expression of m1-m4 muscarinic acetylcholine receptor proteins in rat hippocampus and regulation by cholinergic innervation. *J Neurosci Off J Soc Neurosci*. 1995;15:4077–4092.
188. Hájos N, Papp EC, Acsády L, Levey AI, Freund TF. Distinct interneuron types express m2 muscarinic receptor immunoreactivity on their dendrites or axon terminals in the hippocampus. *Neuroscience*. 1998;82:355–376.
189. Szabó GG, Holderith N, Gulyás AI, Freund TF, Hájos N. Distinct synaptic properties of perisomatic inhibitory cell types and their different modulation by cholinergic receptor activation in the CA3 region of the mouse hippocampus. *Eur J Neurosci*. 2010;31:2234–2246.
190. Rouse ST, Levey AI. Muscarinic acetylcholine receptor immunoreactivity after hippocampal commissural/associational pathway lesions: evidence for multiple presynaptic receptor subtypes. *J Comp Neurol*. 1997;380:382–394.
191. Zhang Z, Reboreda A, Alonso A, Barker PA, Séguéla P. TRPC channels underlie cholinergic plateau potentials and persistent activity in entorhinal cortex. *Hippocampus*. 2011;21:386–397.
192. Wu M, Shanabrough M, Leranath C, Alreja M. Cholinergic excitation of septohippocampal GABA but not cholinergic neurons: implications for learning and memory. *J Neurosci Off J Soc Neurosci*. 2000;20:3900–3908.
193. Wu M, Hajszan T, Leranath C, Alreja M. Nicotine recruits a local glutamatergic circuit to excite septohippocampal GABAergic neurons. *Eur J Neurosci*. 2003;18:1155–1168.
194. Frazier CJ, Buhler AV, Weiner JL, Dunwiddie TV. Synaptic potentials mediated via alpha-bungarotoxin-sensitive nicotinic acetylcholine receptors in rat hippocampal interneurons. *J Neurosci Off J Soc Neurosci*. 1998;18:8228–8235.
195. McQuiston AR, Madison DV. Nicotinic receptor activation excites distinct subtypes of interneurons in the rat hippocampus. *J Neurosci Off J Soc Neurosci*. 1999;19:2887–2896.
196. Ji D, Dani JA. Inhibition and disinhibition of pyramidal neurons by activation of nicotinic receptors on hippocampal interneurons. *J Neurophysiol*. 2000;83:2682–2690.

197. Kawai H, Zago W, Berg DK. Nicotinic alpha 7 receptor clusters on hippocampal GABAergic neurons: regulation by synaptic activity and neurotrophins. *J Neurosci Off J Soc Neurosci*. 2002;22:7903–7912.
198. Wanaverbecq N, Semyanov A, Pavlov I, Walker MC, Kullmann DM. Cholinergic axons modulate GABAergic signaling among hippocampal interneurons via postsynaptic alpha 7 nicotinic receptors. *J Neurosci Off J Soc Neurosci*. 2007;27:5683–5693.
199. Gu Z, Yakel JL. Timing-dependent septal cholinergic induction of dynamic hippocampal synaptic plasticity. *Neuron*. 2011;71:155–165.
200. Margiotta JF, Berg DK, Dionne VE. Cyclic AMP regulates the proportion of functional acetylcholine receptors on chicken ciliary ganglion neurons. *Proc Natl Acad Sci U S A*. 1987;84:8155–8159.
201. Sargent PB. The diversity of neuronal nicotinic acetylcholine receptors. *Annu Rev Neurosci*. 1993;16:403–443.
202. Vijayaraghavan S, Pugh PC, Zhang ZW, Rathouz MM, Berg DK. Nicotinic receptors that bind alpha-bungarotoxin on neurons raise intracellular free Ca²⁺. *Neuron*. 1992;8:353–362.
203. Sharma G, Vijayaraghavan S. Modulation of presynaptic store calcium induces release of glutamate and postsynaptic firing. *Neuron*. 2003;38:929–939.
204. Williams JH, Kauer JA. Properties of carbachol-induced oscillatory activity in rat hippocampus. *J Neurophysiol*. 1997;78:2631–2640.
205. Cobb SR, Bulters DO, Suchak S, Riedel G, Morris RG, Davies CH. Activation of nicotinic acetylcholine receptors patterns network activity in the rodent hippocampus. *J Physiol*. 1999;518:131–140.
206. Griguoli M, Scuri R, Ragozzino D, Cherubini E. Activation of nicotinic acetylcholine receptors enhances a slow calcium-dependent potassium conductance and reduces the firing of stratum oriens interneurons. *Eur J Neurosci*. 2009;30:1011–1022.
207. Madison DV, Lancaster B, Nicoll RA. Voltage clamp analysis of cholinergic action in the hippocampus. *J Neurosci Off J Soc Neurosci*. 1987;7:733–741.
208. Lawrence JJ, Statland JM, Grinspan ZM, McBain CJ. Cell type-specific dependence of muscarinic signalling in mouse hippocampal stratum oriens interneurons. *J Physiol*. 2006;570:595–610.
209. Kahle JS, Cotman CW. Carbachol depresses synaptic responses in the medial but not the lateral perforant path. *Brain Res*. 1989;482:159–163.
210. Foster TC, Deadwyler SA. Acetylcholine modulates averaged sensory evoked responses and perforant path evoked field potentials in the rat dentate gyrus. *Brain Res*. 1992;587:95–101.

211. Hasselmo ME, Schnell E, Barkai E. Dynamics of learning and recall at excitatory recurrent synapses and cholinergic modulation in rat hippocampal region CA3. *J Neurosci Off J Soc Neurosci*. 1995;15:5249–5262.
212. Qian J, Saggau P. Presynaptic inhibition of synaptic transmission in the rat hippocampus by activation of muscarinic receptors: involvement of presynaptic calcium influx. *Br J Pharmacol*. 1997;122:511–519.
213. Seeger T, Alzheimer C. Muscarinic activation of inwardly rectifying K(+) conductance reduces EPSPs in rat hippocampal CA1 pyramidal cells. *J Physiol*. 2001;535:383–396.
214. Kunitake A, Kunitake T, Stewart M. Differential modulation by carbachol of four separate excitatory afferent systems to the rat subiculum in vitro. *Hippocampus*. 2004;14:986–999.
215. Dasari S, Gullidge AT. M1 and M4 receptors modulate hippocampal pyramidal neurons. *J Neurophysiol*. 2011;105:779–792.
216. Jerusalinsky D, Kornisiuk E, Izquierdo I. Cholinergic neurotransmission and synaptic plasticity concerning memory processing. *Neurochem Res*. 1997;22:507–515.
217. Segal M, Auerbach JM. Muscarinic receptors involved in hippocampal plasticity. *Life Sci*. 1997;60:1085–1091.
218. Leung LS, Shen B, Rajakumar N, Ma J. Cholinergic activity enhances hippocampal long-term potentiation in CA1 during walking in rats. *J Neurosci Off J Soc Neurosci*. 2003;23:9297–9304.
219. Ovsepian SV, Anwyl R, Rowan MJ. Endogenous acetylcholine lowers the threshold for long-term potentiation induction in the CA1 area through muscarinic receptor activation: in vivo study. *Eur J Neurosci*. 2004;20:1267–1275.
220. Shinoe T, Matsui M, Taketo MM, Manabe T. Modulation of synaptic plasticity by physiological activation of M1 muscarinic acetylcholine receptors in the mouse hippocampus. *J Neurosci Off J Soc Neurosci*. 2005;25:11194–11200.
221. Dorralp S, Leung LS. Cholinergic modulation of hippocampal CA1 basal-dendritic long-term potentiation. *Neurobiol Learn Mem*. 2008;90:382–388.
222. Isaac JTR, Buchanan KA, Muller RU, Mellor JR. Hippocampal place cell firing patterns can induce long-term synaptic plasticity in vitro. *J Neurosci Off J Soc Neurosci*. 2009;29:6840–6850.
223. Buchanan KA, Petrovic MM, Chamberlain SEL, Marrison NV, Mellor JR. Facilitation of long-term potentiation by muscarinic M(1) receptors is mediated by inhibition of SK channels. *Neuron*. 2010;68:948–963.
224. Jo J, Son GH, Winters BL, Kim MJ, Whitcomb DJ, Dickinson BA, Lee Y-B, Futai K, Amici M, Sheng M, Collingridge GL, Cho K. Muscarinic receptors induce LTD of NMDAR EPSCs via a mechanism involving hippocalcin, AP2 and PSD-95. *Nat Neurosci*. 2010;13:1216–1224.

225. Sugisaki E, Fukushima Y, Tsukada M, Aihara T. Cholinergic modulation on spike timing-dependent plasticity in hippocampal CA1 network. *Neuroscience*. 2011;192:91–101.
226. Fujii S, Sumikawa K. Nicotine accelerates reversal of long-term potentiation and enhances long-term depression in the rat hippocampal CA1 region. *Brain Res*. 2001;894:340–346.
227. Ge S, Dani JA. Nicotinic acetylcholine receptors at glutamate synapses facilitate long-term depression or potentiation. *J Neurosci Off J Soc Neurosci*. 2005;25:6084–6091.
228. Gu Z, Lamb PW, Yakel JL. Cholinergic coordination of presynaptic and postsynaptic activity induces timing-dependent hippocampal synaptic plasticity. *J Neurosci Off J Soc Neurosci*. 2012;32:12337–12348.
229. Wainer BH, Steininger TL, Roback JD, Burke-Watson MA, Mufson EJ, Kordower J. Ascending cholinergic pathways: functional organization and implications for disease models. *Prog Brain Res*. 1993;98:9–30.
230. Bender F, Gorbati M, Cadavieco MC, Denisova N, Gao X, Holman C, Korotkova T, Ponomarenko A. Theta oscillations regulate the speed of locomotion via a hippocampus to lateral septum pathway. *Nat Commun*. 2015;6:8521.
231. Gorelova N, Reiner PB. Role of the afterhyperpolarization in control of discharge properties of septal cholinergic neurons in vitro. *J Neurophysiol*. 1996;75:695–706.
232. Apartis E, Poindessous-Jazat FR, Lamour YA, Bassant MH. Loss of rhythmically bursting neurons in rat medial septum following selective lesion of septohippocampal cholinergic system. *J Neurophysiol*. 1998;79:1633–1642.
233. Lee MG, Chrobak JJ, Sik A, Wiley RG, Buzsáki G. Hippocampal theta activity following selective lesion of the septal cholinergic system. *Neuroscience*. 1994;62:1033–1047.
234. Vandecasteele M, Varga V, Berényi A, Papp E, Barthó P, Venance L, Freund TF, Buzsáki G. Optogenetic activation of septal cholinergic neurons suppresses sharp wave ripples and enhances theta oscillations in the hippocampus. *Proc Natl Acad Sci U S A*. 2014;111:13535–13540.
235. Materi LM, Semba K. Inhibition of synaptically evoked cortical acetylcholine release by intracortical glutamate: involvement of GABAergic neurons. *Eur J Neurosci*. 2001;14:38–46.
236. Clement EA, Richard A, Thwaites M, Ailon J, Peters S, Dickson CT. Cyclic and sleep-like spontaneous alternations of brain state under urethane anaesthesia. *PLoS One*. 2008;3:e2004.
237. Chapman CA, Lacaille JC. Cholinergic induction of theta-frequency oscillations in hippocampal inhibitory interneurons and pacing of pyramidal cell firing. *J Neurosci Off J Soc Neurosci*. 1999;19:8637–8645.

238. Lengyel M, Huhn Z, Erdi P. Computational theories on the function of theta oscillations. *Biol Cybern.* 2005;92:393–408.
239. Greenstein YJ, Pavlides C, Winson J. Long-term potentiation in the dentate gyrus is preferentially induced at theta rhythm periodicity. *Brain Res.* 1988;438:331–334.
240. Huerta PT, Lisman JE. Heightened synaptic plasticity of hippocampal CA1 neurons during a cholinergically induced rhythmic state. *Nature.* 1993;364:723–725.
241. Huerta PT, Lisman JE. Bidirectional synaptic plasticity induced by a single burst during cholinergic theta oscillation in CA1 in vitro. *Neuron.* 1995;15:1053–1063.
242. Hölscher C, Anwyl R, Rowan MJ. Stimulation on the positive phase of hippocampal theta rhythm induces long-term potentiation that can be depotentiated by stimulation on the negative phase in area CA1 in vivo. *J Neurosci Off J Soc Neurosci.* 1997;17:6470–6477.
243. Hyman JM, Wyble BP, Goyal V, Rossi CA, Hasselmo ME. Stimulation in hippocampal region CA1 in behaving rats yields long-term potentiation when delivered to the peak of theta and long-term depression when delivered to the trough. *J Neurosci Off J Soc Neurosci.* 2003;23:11725–11731.
244. Henderson Z, Lu CB, Janzso G, Matto N, McKinley CE, Yanagawa Y, Halasy K. Distribution and role of Kv3.1b in neurons in the medial septum diagonal band complex. *Neuroscience.* 2010;166:952–969.
245. Sun Y, Nguyen AQ, Nguyen JP, Le L, Saur D, Choi J, Callaway EM, Xu X. Cell-type-specific circuit connectivity of hippocampal CA1 revealed through Cre-dependent rabies tracing. *Cell Rep.* 2014;7:269–280.
246. Vanderwolf CH. Hippocampal electrical activity and voluntary movement in the rat. *Electroencephalogr Clin Neurophysiol.* 1969;26:407–418.
247. McFarland WL, Teitelbaum H, Hedges EK. Relationship between hippocampal theta activity and running speed in the rat. *J Comp Physiol Psychol.* 1975;88:324–328.
248. Czurkó A, Hirase H, Csicsvari J, Buzsáki G. Sustained activation of hippocampal pyramidal cells by “space clamping” in a running wheel. *Eur J Neurosci.* 1999;11:344–352.
249. McNaughton BL, Barnes CA, O’Keefe J. The contributions of position, direction, and velocity to single unit activity in the hippocampus of freely-moving rats. *Exp Brain Res.* 1983;52:41–49.
250. Ekstrom AD, Meltzer J, McNaughton BL, Barnes CA. NMDA receptor antagonism blocks experience-dependent expansion of hippocampal “place fields.” *Neuron.* 2001;31:631–638.
251. Fuhrmann F, Justus D, Sosulina L, Kaneko H, Beutel T, Friedrichs D, Schoch S, Schwarz MK, Fuhrmann M, Remy S. Locomotion, Theta Oscillations, and the Speed-Correlated Firing of Hippocampal Neurons Are Controlled by a Medial Septal Glutamatergic Circuit. *Neuron.* 2015;86:1253–1264.

252. Robinson J, Manseau F, Ducharme G, Amilhon B, Vigneault E, El Mestikawy S, Williams S. Optogenetic Activation of Septal Glutamatergic Neurons Drive Hippocampal Theta Rhythms. *J Neurosci Off J Soc Neurosci*. 2016;36:3016–3023.
253. Goutagny R, Manseau F, Jackson J, Danik M, Williams S. In vitro activation of the medial septum-diagonal band complex generates atropine-sensitive and atropine-resistant hippocampal theta rhythm: an investigation using a complete septohippocampal preparation. *Hippocampus*. 2008;18:531–535.
254. Segal M. Brain stem afferents to the rat medial septum. *J Physiol*. 1976;261:617–631.
255. Wilson CL, Motter BC, Lindsley DB. Influences of hypothalamic stimulation upon septal and hippocampal electrical activity in the cat. *Brain Res*. 1976;107:55–68.
256. Dutar P, Bassant MH, Senut MC, Lamour Y. The septohippocampal pathway: structure and function of a central cholinergic system. *Physiol Rev*. 1995;75:393–427.
257. Witter MP, Groenewegen HJ. Chapter 4 The subiculum: cytoarchitectonically a simple structure, but hodologically complex. In: Storm-Mathisen J, Zimmer J, Ottersen OP, eds. *Progress in Brain Research*. Elsevier; 1990: 47–58.
258. O'Mara SM, Commins S, Anderson M, Gigg J. The subiculum: a review of form, physiology and function. *Prog Neurobiol*. 2001;64:129–155.
259. Cooper DC, Moore SJ, Staff NP, Spruston N. Psychostimulant-Induced Plasticity of Intrinsic Neuronal Excitability in Ventral Subiculum. *J Neurosci*. 2003;23:9937–9946.
260. Jarsky T, Mady R, Kennedy B, Spruston N. Distribution of bursting neurons in the CA1 region and the subiculum of the rat hippocampus. *J Comp Neurol*. 2008;506:535–547.
261. Prida LM de la, Suarez F, Pozo MA. Electrophysiological and morphological diversity of neurons from the rat subicular complex in vitro. *Hippocampus*. 2003;13:728–744.
262. Staff NP, Jung H-Y, Thiagarajan T, Yao M, Spruston N. Resting and Active Properties of Pyramidal Neurons in Subiculum and CA1 of Rat Hippocampus. *J Neurophysiol*. 2000;84:2398–2408.
263. Taube JS. Electrophysiological properties of neurons in the rat subiculum in vitro. *Exp Brain Res*. 1993;96:304–318.
264. Canteras NS, Swanson LW. Projections of the ventral subiculum to the amygdala, septum, and hypothalamus: A PHAL anterograde tract-tracing study in the rat. *J Comp Neurol*. 1992;324:180–194.
265. Namura S, Takada M, Kikuchi H, Mizuno N. Topographical organization of subicular neurons projecting to subcortical regions. *Brain Res Bull*. 1994;35:221–231.
266. Verwer RWH, Meijer RJ, Uum HFMV, Witter MP. Collateral projections from the rat hippocampal formation to the lateral and medial prefrontal cortex. *Hippocampus*. 1997;7:397–402.

267. van Groen T, Wyss JM. The connections of presubiculum and parasubiculum in the rat. *Brain Res.* 1990;518:227–243.
268. Witter MP. Connections of the subiculum of the rat: Topography in relation to columnar and laminar organization. *Behav Brain Res.* 2006;174:251–264.
269. Naber PA, Witter MP. Subicular efferents are organized mostly as parallel projections: A double-labeling, retrograde-tracing study in the rat. *J Comp Neurol.* 1998;393:284–297.
270. Swanson LW, Sawchenko PE, Cowan WM. Evidence for collateral projections by neurons in Ammon's horn, the dentate gyrus, and the subiculum: a multiple retrograde labeling study in the rat. *J Neurosci.* 1981;1:548–559.
271. Donovan MK, Wyss JM. Evidence for some collateralization between cortical and diencephalic efferent axons of the rat subicular cortex. *Brain Res.* 1983;259:181–192.
272. Stewart M. Antidromic and orthodromic responses by subicular neurons in rat brain slices. *Brain Res.* 1997;769:71–85.
273. Kim Y, Spruston N. Target-specific output patterns are predicted by the distribution of regular-spiking and bursting pyramidal neurons in the subiculum. *Hippocampus.* 2012;22:693–706.
274. Wu CP, Huang HL, Asl MN, He JW, Gillis J, Skinner FK, Zhang L. Spontaneous rhythmic field potentials of isolated mouse hippocampal–subicular–entorhinal cortices in vitro. *J Physiol.* 2006;576:457–476.
275. Jackson J, Goutagny R, Williams S. Fast and Slow Gamma Rhythms Are Intrinsically and Independently Generated in the Subiculum. *J Neurosci.* 2011;31:12104–12117.
276. Colling SB, Stanford IM, Traub RD, Jefferys JGR. Limbic Gamma Rhythms. I. Phase-Locked Oscillations in Hippocampal CA1 and Subiculum. *J Neurophysiol.* 1998;80:155–161.
277. Stanford IM, Traub RD, Jefferys JGR. Limbic Gamma Rhythms. II. Synaptic and Intrinsic Mechanisms Underlying Spike Doublets in Oscillating Subicular Neurons. *J Neurophysiol.* 1998;80:162–171.
278. Gigg J, Finch DM, O'Mara SM. Responses of rat subicular neurons to convergent stimulation of lateral entorhinal cortex and CA1 in vivo. *Brain Res.* 2000;884:35–50.
279. Commins S, Anderson M, Gigg J, O'Mara SM. The effects of single and multiple episodes of theta patterned or high frequency stimulation on synaptic transmission from hippocampal area CA1 to the subiculum in rats. *Neurosci Lett.* 1999;270:99–102.
280. Commins S, Gigg J, Anderson M, O'Mara SM. Interaction between paired-pulse facilitation and long-term potentiation in the projection from hippocampal area CA1 to the subiculum. *NeuroReport.* 1998;9:4109.
281. Commins S, Gigg J, Anderson M, O'Mara SM. The projection from hippocampal area CA1 to the subiculum sustains long-term potentiation. *NeuroReport.* 1998;9:847.

282. Fidzinski P, Shor O, Behr J. Target-cell-specific bidirectional synaptic plasticity at hippocampal output synapses. *Eur J Neurosci*. 2008;27:1111–1118.
283. Fidzinski P, Wawra M, Dugladze T, Gloveli T, Heinemann U, Behr J. Low-frequency stimulation of the temporoammonic pathway induces heterosynaptic disinhibition in the subiculum. *Hippocampus*. 2011;21:733–743.
284. Fidzinski P, Wawra M, Bartsch J, Heinemann U, Behr J. High-frequency stimulation of the temporoammonic pathway induces input-specific long-term potentiation in subicular bursting cells. *Brain Res*. 2012;1430:1–7.
285. Sharp PE, Green C. Spatial correlates of firing patterns of single cells in the subiculum of the freely moving rat. *J Neurosci*. 1994;14:2339–2356.
286. Sharp PE. Subicular place cells expand or contract their spatial firing pattern to fit the size of the environment in an open field but not in the presence of barriers: Comparison with hippocampal place cells. *Behav Neurosci*. 1999;113:643–662.
287. O'Mara SM, Commins S, Anderson M. Synaptic plasticity in the hippocampal area CA1-subiculum projection: Implications for theories of memory. *Hippocampus*. 2000;10:447–456.
288. Taube JS, Muller RU. Comparisons of head direction cell activity in the postsubiculum and anterior thalamus of freely moving rats. *Hippocampus*. 1998;8:87–108.
289. Deadwyler SA, Hampson RE. Differential but Complementary Mnemonic Functions of the Hippocampus and Subiculum. *Neuron*. 2004;42:465–476.
290. Anderson MI, O'Mara SM. Responses of dorsal subicular neurons of rats during object exploration in an extended environment. *Exp Brain Res*. 2004;159:519–529.
291. Pare D, Dossi RC, Steriade M. Three types of inhibitory postsynaptic potentials generated by interneurons in the anterior thalamic complex of cat. *J Neurophysiol*. 1991;66:1190–1204.
292. Somogyi Gy, Hajdu F, Tömböl T. Ultrastructure of the anterior ventral and anterior medial nuclei of the cat thalamus. *Exp Brain Res*. 1978;31:417–431.
293. Sikes RW, Chronister RB, White LE. Origin of the direct hippocampus-anterior thalamic bundle in the rat: A combined horseradish peroxidase-Golgi analysis. *Exp Neurol*. 1977;57:379–395.
294. Swanson LW, Cowan WM. An autoradiographic study of the organization of the efferent connections of the hippocampal formation in the rat. *J Comp Neurol*. 1977;172:49–84.
295. Seki M, Zyo K. Anterior thalamic afferents from the mamillary body and the limbic cortex in the rat. *J Comp Neurol*. 1984;229:242–256.
296. Shibata H. Topographic organization of subcortical projections to the anterior thalamic nuclei in the rat. *J Comp Neurol*. 1992;323:117–127.

297. Amaral DG, Witter MP. The three-dimensional organization of the hippocampal formation: A review of anatomical data. *Neuroscience*. 1989;31:571–591.
298. Taube JS. Place cells recorded in the parasubiculum of freely moving rats. *Hippocampus*. 1995;5:569–583.
299. Jahnsen H, Llinás R. Ionic basis for the electro-responsiveness and oscillatory properties of guinea-pig thalamic neurones in vitro. *J Physiol*. 1984;349:227–247.
300. Pare D, Steriade M, Deschenes M, Oakson G. Physiological characteristics of anterior thalamic nuclei, a group devoid of inputs from reticular thalamic nucleus. *J Neurophysiol*. 1987;57:1669–1685.
301. Tsanov M, Wright N, Vann SD, Erichsen JT, Aggleton JP, O’Mara SM. Hippocampal inputs mediate theta-related plasticity in anterior thalamus. *Neuroscience*. 2011;187:52–62.
302. Vertes RP, Albo Z, Viana Di Prisco G. Theta-rhythmically firing neurons in the anterior thalamus: implications for mnemonic functions of Papez’s circuit. *Neuroscience*. 2001;104:619–625.
303. Albo Z, Prisco GVD, Vertes R. Anterior thalamic unit discharge profiles and coherence with hippocampal theta rhythm. *Thalamus Relat Syst*. 2003;2:133–144.
304. Tsanov M, Chah E, Vann SD, Reilly RB, Erichsen JT, Aggleton JP, O’Mara SM. Theta-Modulated Head Direction Cells in the Rat Anterior Thalamus. *J Neurosci*. 2011;31:9489–9502.
305. Blair HT, Cho J, Sharp PE. The Anterior Thalamic Head-Direction Signal Is Abolished by Bilateral But Not Unilateral Lesions of the Lateral Mammillary Nucleus. *J Neurosci*. 1999;19:6673–6683.
306. Zugaro MB, Tabuchi E, Fouquier C, Berthoz A, Wiener SI. Active Locomotion Increases Peak Firing Rates of Anterodorsal Thalamic Head Direction Cells. *J Neurophysiol*. 2001;86:692–702.
307. Vann SD, Aggleton JP. The mammillary bodies: two memory systems in one? *Nat Rev Neurosci*. 2004;5:35–44.
308. Aggleton JP, O’Mara SM, Vann SD, Wright NF, Tsanov M, Erichsen JT. Hippocampal–anterior thalamic pathways for memory: uncovering a network of direct and indirect actions. *Eur J Neurosci*. 2010;31:2292–2307.
309. Swanson LW, Cowan WM. Hippocampo-hypothalamic connections: origin in subicular cortex, not ammon’s horn. *Science*. 1975;189:303–304.
310. Allen GV, Hopkins DA. Mamillary body in the rat: Topography and synaptology of projections from the subicular complex, prefrontal cortex, and midbrain tegmentum. *J Comp Neurol*. 1989;286:311–336.
311. Allen GV, Hopkins DA. Topography and synaptology of mamillary body projections to the mesencephalon and pons in the rat. *J Comp Neurol*. 1990;301:214–231.

312. Gonzalo-Ruiz A, Morte L, Sanz JM. Glutamate/aspartate and leu-enkephalin immunoreactivity in mammillothalamic projection neurons of the rat. *Brain Res Bull.* 1998;47:565–574.
313. Bassett JP, Tullman ML, Taube JS. Lesions of the Tegmentomammillary Circuit in the Head Direction System Disrupt the Head Direction Signal in the Anterior Thalamus. *J Neurosci.* 2007;27:7564–7577.
314. Kocsis B, Vertes RP. Characterization of neurons of the supramammillary nucleus and mammillary body that discharge rhythmically with the hippocampal theta rhythm in the rat. *J Neurosci.* 1994;14:7040–7052.
315. Vertes RP, Kocsis B. Brainstem-diencephalo-septohippocampal systems controlling the theta rhythm of the hippocampus. *Neuroscience.* 1997;81:893–926.
316. Aggleton JP, Brown MW. Episodic memory, amnesia, and the hippocampal–anterior thalamic axis. *Behav Brain Sci.* 1999;22:425–444.
317. Ungiadze AA. [Responses of the gyrus cinguli to stimulation of the dorsal hippocampus in cats]. *Neirofiziologija Neurophysiol.* 1981;13:7–13.
318. Abbott LF, Varela JA, Sen K, Nelson SB. Synaptic Depression and Cortical Gain Control. *Science.* 1997;275:221–224.
319. Heschem S, Jahanshahi A, Schweimer JV, Mitchell SN, Carter G, Blokland A, Sharp T, Temel Y. Fornix deep brain stimulation enhances acetylcholine levels in the hippocampus. *Brain Struct Funct.* 2016;221:4281–4286.
320. Heschem S, Lim LW, Jahanshahi A, Steinbusch HWM, Prickaerts J, Blokland A, Temel Y. Deep brain stimulation of the fornical area enhances memory functions in experimental dementia: The role of stimulation parameters. *Brain Stimulat.* 2013;6:72–77.
321. Smith GS, Laxton AW, Tang-Wai DF, McAndrews MP, Diaconescu AO, Workman CI, Lozano AM. Increased Cerebral Metabolism After 1 Year of Deep Brain Stimulation in Alzheimer Disease. *Arch Neurol.* 2012;69:1141–1148.
322. Smith GS, Leon MJ de, George AE, Kluger A, Volkow ND, McRae T, Golomb J, Ferris SH, Reisberg B, Ciaravino J, Regina MEL. Topography of Cross-sectional and Longitudinal Glucose Metabolic Deficits in Alzheimer’s Disease: Pathophysiologic Implications. *Arch Neurol.* 1992;49:1142–1150.
323. Alexander GE, Chen K, Pietrini P, Rapoport SI, Reiman EM. Longitudinal PET Evaluation of Cerebral Metabolic Decline in Dementia: A Potential Outcome Measure in Alzheimer’s Disease Treatment Studies. *Am J Psychiatry.* 2002;159:738–745.
324. Laxton AW, Tang-Wai DF, McAndrews MP, Zumsteg D, Wennberg R, Keren R, Wherrett J, Naglie G, Hamani C, Smith GS, Lozano AM. A phase I trial of deep brain stimulation of memory circuits in Alzheimer’s disease. *Ann Neurol.* 2010;68:521–534.

325. Fontaine D, Deudon A, Lemaire JJ, Razzouk M, Viau P, Darcourt J, Robert P. Symptomatic Treatment of Memory Decline in Alzheimer's Disease by Deep Brain Stimulation: A Feasibility Study. *J Alzheimers Dis.* 2013;34:315–323.
326. Lozano AM, Fosdick L, Chakravarty MM, Leoutsakos J-M, Munro C, Oh E, Drake KE, Lyman CH, Rosenberg PB, Anderson WS, Tang-Wai DF, Pendergrass JC, Salloway S, Asaad WF, Ponce FA, Burke A, Sabbagh M, Wolk DA, Baltuch G, Okun MS, Foote KD, McAndrews MP, Giacobbe P, Targum SD, Lyketsos CG, Smith GS. A Phase II Study of Fornix Deep Brain Stimulation in Mild Alzheimer's Disease. *J Alzheimers Dis.* 2016;54:777–787.
327. Buzsáki G. Theta Oscillations in the Hippocampus. *Neuron.* 2002;33:325–340.
328. Hasselmo ME, Bodelón C, Wyble BP. A Proposed Function for Hippocampal Theta Rhythm: Separate Phases of Encoding and Retrieval Enhance Reversal of Prior Learning. *Neural Comput.* 2002;14:793–817.
329. Bland BH, Konopacki J, Kirk IJ, Oddie SD, Dickson CT. Discharge patterns of hippocampal theta-related cells in the caudal diencephalon of the urethan-anesthetized rat. *J Neurophysiol.* 1995;74:322–333.
330. Senova S, Chaillet A, Lozano AM. Fornical Closed-Loop Stimulation for Alzheimer's Disease. *Trends Neurosci.* 2018;41:418–428.