

Trauma and depressive symptomatology in middle-aged persons at high risk of dementia: the PREVENT Dementia Study

Karen Ritchie ,^{1,2,3} Isabelle Carrière,¹ Sarah Gregory,² Tam Watermeyer,^{2,4} Samuel Danso,² Li Su,⁵ Craig W Ritchie,² John T O'Brien ⁵

¹U1061 Neuropsychiatry, INSERM, University of Montpellier, Montpellier, France
²Centre for Dementia Prevention, Centre for Clinical Brain Sciences, University of Edinburgh, Edinburgh, UK
³Institut du Cerveau, Paris, France
⁴Department of Psychology, Northumbria University, Newcastle upon Tyne, UK
⁵Department of Psychiatry, University of Cambridge, Cambridge, UK

Correspondence to

Prof Karen Ritchie, U1061 Neuropsychiatry, INSERM University of Montpellier, Montpellier, France 34093; karen.ritchie@inserm.fr

KR and IC are joint first authors.

Received 11 May 2020
 Revised 23 September 2020
 Accepted 23 September 2020
 Published Online First 21 October 2020

ABSTRACT

Objective Depression and trauma are associated with changes in brain regions implicated in Alzheimer's disease. The present study examined associations between childhood trauma, depression, adult cognitive functioning and risk of dementia.

Methods Data from 378 participants in the PREVENT Dementia Study aged 40–59 years. Linear and logistic models were used to assess associations between childhood trauma, depression, dementia risk, cognitive test scores and hippocampal volume.

Results Childhood trauma was associated with depression and reduced hippocampal volume but not current cognitive function or dementia risk. Poorer performance on a delayed face/name recall task was associated with depression. Childhood trauma was associated with lower hippocampal volume however poorer cognitive performance was mediated by depression rather than structural brain differences.

Conclusion Depressive symptomatology may be associated with dementia risk via multiple pathways, and future studies should consider subtypes of depressive symptomatology when examining its relationship to dementia.

loss⁶ or white matter (WM) lesions⁷ accompanied by serotonergic deficiency and a dopamine–norepinephrine imbalance.⁸ Other studies have reported associations with tau accumulation and amyloid metabolism.^{9–12} Several epidemiological studies further suggest that depression onset occurs mostly in the few years preceding the diagnosis of AD dementia, and is therefore prodromal.¹² A comparison of familial AD mutation carriers with their non-carrier relatives found reduced depression scores in mutation carriers in the presymptomatic stages of the disease, but elevated levels in carriers once dementia symptoms emerged, suggesting depression may be a symptom of onset rather than a risk factor for AD.¹³

On the other hand a meta-analysis of prospective studies showed an inverse association with proximity to dementia, suggesting it to be a risk factor rather than a prodrome,¹⁴ and a postmortem study of persons with current major depression found no association between AD neuropathology and cognitive loss¹⁵ suggestive of independent aetiological processes.

Longer term prospective studies reaching back into the preclinical phases of AD, and the possible origin of the association, are lacking. A recent 10-year retrospective study of clinical and biomarker changes in persons with incident AD observed that depressive symptomatology was significantly higher 10 years before dementia diagnosis, remaining at a stable level to up to diagnosis independently of changes in other AD markers.¹⁶ This finding suggests the lack of a dose–response effect in the association between depression and biomarker changes, and that depression may pre-date brain changes in AD. Depression thus appears to be linked to, but not entirely explained by prodromal AD neurodegeneration. Moreover, depression may be the endpoint of multiple aetiological pathways implicating different underlying neuropathological processes and having different associations with AD onset.¹⁷ Research in this area should thus attempt to focus on more homogeneous subtypes of depression.

Neurobiological studies have demonstrated that childhood maltreatment may alter brain development by programming the glucocorticoid, noradrenergic and vasopressin stress response systems to over-react to new stressors,¹⁸ thus rendering the individual increasingly vulnerable to psychiatric

INTRODUCTION

Alzheimer's disease (AD) and depressive disorders were once considered to be mutually exclusive; the co-occurrence of depression and cognitive decline being termed 'pseudodementia'.¹ Depression is now known to be closely associated with AD, with a prevalence around 50% at diagnosis^{2,3} compared with 20%–25% for the general elderly population.⁴ A meta-analysis⁵ based on prospective population studies and adjusting for multiple confounders has estimated that late-life depression increases 1.65-fold the risk of future dementia, and also increases the risk of progression from mild cognitive impairment to dementia.

A central problem has been that the underlying mechanism which links the two pathologies is still uncertain, and controversy continues as to whether depression is a risk factor for AD or a prodromal feature of the disease itself. There is strong evidence that depressive symptoms are a direct result of AD-related brain changes. Structural and functional brain changes show depression in AD to be related to disruption of frontal-striatal and subcortical limbic pathways, either due to grey matter (GM)



© Author(s) (or their employer(s)) 2021. No commercial re-use. See rights and permissions. Published by BMJ.

To cite: Ritchie K, Carrière I, Gregory S, et al. *J Neurol Neurosurg Psychiatry* 2021;**92**:16–21.

disorder. These effects appear to be long-lasting¹⁹ inducing structural and functional changes, notably reduced development of the hippocampus and amygdala, and abnormal frontotemporal electrical activity.¹⁸ These brain structures having also been implicated in the aetiology of both depression and dementia, it is not surprising that child abuse has been associated with not only increased risk for lifetime depression^{20 21} but also lower cognitive scores.²² Several small clinical studies of young adult women with a history of physical and sexual abuse reported disorders of vigilance, memory and mathematical ability compared with controls.^{23–25}

Depression associated with childhood trauma may thus provide a clear-cut model of the association of a depression subtype with AD.

The present study is primarily designed to determine whether the association between childhood trauma and poorer cognitive performance is directly due to trauma-related hippocampal volume or mediated by depression. It further seeks to establish whether the impact on cognition may be greater in persons at genetic risk of dementia, thereby increasing brain burden in the face of possible pre-clinical changes. A secondary aim is to develop a putative model of the associations between trauma, depression, cognitive performance and dementia risk which may guide future longitudinal studies. The study uses data from a cohort of middle-aged persons in which around half have a family history of dementia in whom preclinical hippocampal changes have already been detected.²⁶

The following hypotheses are examined:

1. Middle-aged persons who have experienced childhood trauma will have higher rates of clinical depression than those who have not.
2. Middle-aged persons exposed to childhood trauma will have lower hippocampal volume.
3. Childhood trauma will be associated with current poorer cognitive function.
4. Cognitive performance will be poorest in persons with both a history of childhood trauma and a high dementia risk.

METHODS

Subjects were recruited from the PREVENT Dementia Study; a prospective population study enriched for dementia risk, with around half of the participants having a family history of dementia. The protocol has been described in detail elsewhere²⁷ (<http://bmjopen.bmj.com/cgi/content/full/bmjopen-2012-001893>). Briefly participants were recruited through the dementia register database, the Joint Dementia Research website and information about the study on the internet and in study presentations. The present study included 378 participants from west London, Oxford and Cambridge for whom baseline data were available and validated on all variables and of whom 203 have a parent with diagnosed AD, vascular or mixed dementia. Persons with a parent with diagnosed dementia were considered to be at increased risk of future dementia and current preclinical brain changes.

Trauma

Lifetime trauma was assessed using the Life Stressor Checklist-Revised,²⁸ a 30-item self-administered questionnaire which assesses age of occurrence, perceived level of danger and subsequent impact on everyday life in the past year. The Diagnostic and Statistical Manual of Mental Disorders, fourth edition (DSM IV)²⁹ criterion A for post-traumatic stress disorder (PTSD) was applied to individual responses to distinguish trauma from

milder transient stressors (for example moving house) and events with low perceived level of threat. Absence of information on the presence of subsequent intrusive symptoms did not allow us to make a formal diagnosis of PTSD. The items retained in relation to criterion A were death of someone close, physical and emotional abuse, sexual abuse, rape, family violence, adoption, separation from parents and neglect. For this study the scale was not formally scored, but rather the reporting of any of these items with a high associated perceived level of danger and subsequent impact on everyday life in the past year was classed as a traumatic event. The group was further divided into those experiencing these events before and after age 16. Within the total sample, 64 participants (16.9%) had been exposed to traumatic events and 54 (14.3%) before the age of 16.

Depression

The Center for Epidemiologic Studies-Depression Scale (CES-D)³⁰ was used to detect high levels of depressive symptomatology. Depression has been treated as a dichotomous variable with cases being defined as a score above the 16 cut-off point of the CES-D or current treatment with an antidepressant. Of the 378 participants examined, 64 (16.9%) had CES-D scores of 16 and over and a further 29 participants (7.7%) were receiving antidepressant treatment.

Cognitive testing

COGNITO is a computerised neuropsychological battery,³¹ designed to examine information processing across a wide range of cognitive functions in adults of all ages. Tests are administered using a tactile screen to capture information processing time as well as response accuracy.

Ten cognitive summary variables from the COGNITO battery were considered as dependent variables:

1. Working memory: the simultaneous presentation of auditory and visual attention tasks assessed by subtracting the time taken in milliseconds on this double task from the visual task alone.
2. Narrative recall: total number of correct elements on immediate recall of a story with a temporal progression requiring attention to macrostructure.
3. Description recall: total number of correct elements recalled of a description without thematic progression requiring attention to microstructure and recall of spatial location. The narrative and description recall are similar in terms of word frequency in the language and syntactic structure.
4. Implicit memory: difference in the number of steps in the progressive build-up of names on the screen required for recognition between names never seen and number of names previously learnt in an immediate recall task.
5. Name-face association: number of faces recognised after a delay from a series of 18 faces of which 9 have been previously shown with their corresponding names.
6. Form perception: number of correct answers in the matching of complex forms to a multiple-choice array.
7. Form perception speed: mean time taken in milliseconds for each trial.⁶
8. Phoneme comprehension: number of correct responses in the matching of a word with an image presented as part of a multiple-choice array including semantic, morphological and phonetic distractors.
9. Phoneme comprehension speed: mean time taken in milliseconds to perform.⁸

10. Verbal fluency: total sum of the number of words generated in 30s using both a semantic (vegetables) and phonemic (letter P) cue.

Imaging

Participants underwent multimodal 3T structural MRI on a single scanner including volumetric T1-weighted scans (176 slices, 1.0×1.0mm, 1.0mm slice thickness, TR=2300 ms, TE=2.98 ms, flip angle 9°). Brain tissue segmentation into GM, WM and cerebrospinal fluid was performed using the Gaussian mixture model in VBM toolbox of SPM8 (<http://www.fil.ion.ucl.ac.uk/spm>). The GM maps were then normalised using the DARTEL algorithm.³² Hippocampal region of interests (ROIs) were selected using AAL atlas in MNI space,^{33 34} and then inverse normalised back to each subject’s native space using the participant-specific diffeomorphic parameters estimated from the previous DARTEL procedure. The resulting ROIs were also masked using the thresholded GM probability maps (at threshold p>0.8) before the total hippocampal volume was calculated.

Statistical analyses

Cases of depression were first compared with non-cases using a X² test for categorical variables and Wilcoxon two-sample test

for continuous variables. The covariates associated with depression with a p<0.10 (except anxiety and educational level due to collinearity with socioeconomic class) were then entered in a logistic model with the sum of traumatic events.

A linear model adjusted for education, gender and age was used to test the association of continuous cognitive scores with depression or having experienced at least one traumatic event before the age of 16 as well as with hippocampal volume. For two non-continuous cognitive scores (form matching and phoneme comprehension), a logistic model was used after dichotomisation (poor performance being defined as <7 and <9 correct responses, respectively). A linear model adjusted for intra-cranial volume was also used to test the association of hippocampal volume with depression and having experienced at least one traumatic event. This latter model on traumatic events was further adjusted for education, gender, age and depression. All the analyses were carried out using SAS, V.9.4.

RESULTS

Trauma and depression

The association between depression and socioeconomic and clinical characteristics is given in [table 1](#). The range of the total

Table 1 The association between depressive symptomatology (CES-D) and clinical and socioeconomic characteristics

	Depressive symptomatology		P value*
	CES-D <16 and not treated (n=285)	CES-D ≥16 or treated (n=93)	
	n (%)	n (%)	
Sex, female	197 (69.1)	66 (71.0)	0.74
Marital status			0.02
Married or living as a couple	223 (78.3)	61 (66.3)	
Educational level			0.06
College/university	204 (71.8)	57 (61.3)	
Living alone	37 (13.0)	17 (18.3)	0.2
Socioeconomic class, n=358			0.04
Higher managerial, administrative occupations	75 (27.6)	14 (16.3)	
Lower managerial, administrative occupations	108 (39.7)	31 (36.0)	
Intermediate occupations	31 (11.4)	17 (19.8)	
Employers, own account, technical, routine occupation	30 (11.0)	16 (18.6)	
Never worked or long-term unemployed	28 (10.3)	8 (9.3)	
BMI (kg/m ²)			0.09
Normal (<25)	115 (40.3)	26 (28.0)	
Overweight (25–30)	106 (37.2)	40 (43.0)	
Obese (≥30)	64 (22.5)	27 (29.0)	
Hypertension (treated or SBP >140)	44 (15.4)	22 (23.7)	0.07
Cardiovascular disease (declared)	24 (8.4)	10 (10.8)	0.5
Diabetes (treated or declared or glycaemia >7 mmol/L), n=377	8 (2.8)	8 (8.6)	0.02
Anxiety symptoms (Spielberger state scale >31)	71 (24.9)	51 (54.8)	<0.0001
Parental history of dementia (n=377)	151 (53.2)	52(55.9)	0.64
Traumatic events at any age (at least 1 event)	39 (13.7)	25 (26.9)	0.003
Traumatic events after 16 years only (at least 1 event)	6 (2.1)	4 (4.3)	0.25
Traumatic events at 16 years or younger (at least 1 early event)	33 (11.6)	21 (22.6)	0.009
	Median (IQR)	Median (IQR)	P value†
Age (years)	53 (47–56)	53 (49–56)	0.76
Hippocampal grey matter volume (mm ³)	3689 (3525–3951)	3681 (3428–3949)	0.66
Intracranial volume (cm ³), n=193	1387 (1299–1477)	1354 (1268–1451)	0.22

*X² test.

†Wilcoxon two-sample test.

‡With parental history of dementia.

BMI, body mass index; CES-D, Center for Epidemiologic Studies-Depression Scale; SBP, systolic blood pressure.

Table 2 Adjusted association between childhood trauma and depression (n=356)

	OR*	95% CI	P value
Marital status: married or living as a couple	0.59	0.34–1.03	0.06
Socioeconomic class			0.03
Higher managerial, administrative occupations	1		
Lower managerial, administrative occupations	1.57	0.76–3.20	
Intermediate occupations, employers, own account, technical, routine occupation	2.94	1.40–6.17	
Never worked or long-term unemployed	1.56	0.57–4.30	
BMI: overweight or obese	1.62	0.91–2.87	0.1
Hypertension (treated or SBP>140)	1.19	0.62–2.31	0.6
Diabetes (treated or declared or glycaemia >7 mmol/L)	2.03	0.58–7.18	0.27
Traumatic events before age 16 (at least one)	2.52	1.33–4.80	0.005

*Adjusted for other variables in the table.

BMI, body mass index; SBP, systolic blood pressure.

sum of traumatic events was 0-5 for childhood events and 0-2 for events after 16 years.

Of the 54 participants exposed to childhood trauma, 21 (38.9%) had a CES-D score of 16 and over or were currently being treated for depression. The association between childhood trauma and depressive symptomatology is significant ($p=0.009$). A multivariate logistic model with depression as the dependent variable was used to examine the association between childhood traumatic events (at least one) adjusted for marital status, socioeconomic status, hypertension, diabetes and body mass index (table 2). Adjustment was not made for anxiety given the high correlation between performance on the depression and anxiety scales due to overlap in items and also probably high rates of mixed anxiety and depression in participants.

Hippocampal volume

Structural imaging data were available for the west London centre only (n=193, see online supplemental table 1 for subsample description). Table 3 shows the associations adjusted for intracranial volume between hippocampal volume as the dependent variable and depression, traumatic events (at least one) before and after the age of 16. Three linear models are presented showing the independent associations of depression, trauma after 16 years and childhood trauma with hippocampal volume.

In the subsample with an MRI examination, 21 participants were treated with antidepressants. In order to control for any possible effect of treatment on brain volume, we also conducted the analyses excluding participants undergoing treatment. The result did not change and the association with depression adjusted for ICV remained non-significant ($p=0.23$).

Table 3 Associations between hippocampal volume as the dependent variable and depression or traumatic events (separate linear models)

Independent variables	Hippocampal grey matter volume (mm ³)	
	β (SE)*	P value
Depressive symptomatology (CES-D ≥ 16 or treated)	44.47 (33.65)	0.19
Traumatic events after 16 years (at least one event)	60.25 (81.73)	0.46
Traumatic events before 16 years (at least one event)	-90.52 (38.20)	0.02

*Adjusted for total intracranial volume.

CES-D, Center for Epidemiologic Studies-Depression Scale;

The association with trauma before age 16 remained significant when it was adjusted for ICV, education level, gender and age (β (SE)=-93.4 (39.0); $p=0.02$) as well as when depression was further added to the model (β (SE)=-99.4 (39.0); $p=0.01$). The relationship with hippocampal volume is not altered with adjustment by depression.

Cognitive performance in middle age

The association between current cognitive performance based on subscores from COGNITO and traumatic events before age 16 was examined using a multivariate model adjusted by education level, gender and age (table 4).

The number of correct answers for form matching and phoneme comprehension was not statistically associated with traumatic events before 16 years, the adjusted OR (95% CI) of poor performances was equal to 0.82 (0.46–1.47) and 0.94 (0.50–1.76), respectively.

A significant association was found between depression and delayed recall of names ($p=0.03$) but not on the other cognitive measures (online supplemental tables 2 and 3). No significant association was found between hippocampal volume and cognitive measures (online supplemental table 4).

Risk of dementia

In persons with a family history of dementia, depression prevalence was 25.6% compared with 23.6% in those without a family history ($p=0.64$). A total of 11.3% of persons with a family history of dementia reported being exposed to childhood trauma compared with 17.8% without ($p=0.07$).

No interaction effect was found between parental history of dementia and depression in relation to cognition. The interaction for name recall is $p=0.54$.

Given the complexity of the observed associations we constructed an association graph (figure 1) designed to facilitate understanding and provide a hypothetical basis for future research.

DISCUSSION

This study aimed to determine whether persons with a history of childhood trauma may be at increased risk of both depression and loss of hippocampal volume in midlife, which may in turn be associated with cognitive performance in persons at increased risk for dementia.

The estimated prevalence of clinically significant depression in this study, based on CES-D scores (16.9%) and current antidepressant treatment (7.7%) was observed to be very high (24.6%) compared with 5.1% estimated for a similar age group in western countries using DSM IV/Composite International Diagnostic Interview criteria.³³ The rate remains high even if treated depression is excluded on the basis that antidepressants may have in some cases been prescribed for reasons other than depression. Although different measures of depression have been used, previous studies show high concordance between DSM major depression and the CES-D.³⁵ While increased rates of depression might have been expected within this cohort enriched for dementia risk, depression prevalence was not observed in this midlife group to be associated with a family history of dementia. One-third of the cohort were exposed to a traumatic event, and of these 18% experienced the trauma before the age of 16. We found an association between history of trauma and both clinical-level depression and volume loss in the hippocampus. Lower hippocampal volume was observed to be specifically related to childhood trauma and not with trauma experienced at

Table 4 Association between childhood trauma and current cognitive performance

Dependent variables	Exposure to traumatic event		P value
	Yes Mean (SEM)* N=54	No Mean (SEM)* N=323	
Working memory: time difference between double task and simple task (milliseconds)	-414.3 (431.9)	25.2 (192.9)	0.33
Form matching: mean time for correct responses (milliseconds)	5781.9 (214.3)	5713.4 (95.7)	0.76
Phoneme comprehension: mean time for correct responses (milliseconds)	1556.7 (41.7)	1500.9 (18.64)	0.20
Name-face associations: number of names correctly recalled	4.71 (0.28)	5.05 (0.13)	0.24
Verbal fluency: number of correct answers	27.80 (0.89)	27.78 (0.39)	0.98
Narrative recall: number of correct answers	12.73 (0.61)	12.79 (0.27)	0.92
Descriptive recall: number of correct answers	11.70 (0.58)	12.42 (0.26)	0.24
Implicit memory: difference between learnt and new items	1.20 (0.10)	1.03 (0.05)	0.13

*Adjusted for education level, gender and age.

other ages. This finding is consistent with previous observations that trauma has its greatest effect on brain development through alterations in Hypothalamic–Pituitary–Adrenal (HPA) axis functioning when occurring before full brain maturation.^{18 19} It also underscores the need for future research to consider the context and timing of traumatic incidents along developmental trajectories.

In this midlife cohort we observed that severe childhood trauma is associated in midlife with both higher rates of clinical depression and lower hippocampal volume. For those persons who have experienced childhood trauma who develop dementia in the future, it may be postulated that lower volume will add to the burden of hippocampal loss associated with the dementia. We did not at this point, however, observe any association with current cognitive performance or increased dementia risk (family history), suggesting that the direction of causality is probably not persons with a family history of dementia being more vulnerable to early-life trauma.

Investigations of the interaction between adverse childhood experiences with genetic risk factors for AD in regard to later-life cognitive deterioration have provided mixed findings.^{22 36} A study of childhood abuse and cognition in bipolar disorder has suggested the possibility of a gene-environment interaction implicating genes known to exert a neurotrophic effect in response to cellular injury³⁷ notably Apolipoprotein E (ApoE). Polymorphisms on the ApoE gene may thus induce compensatory neurite outgrowth and synaptogenesis following neuronal injury³⁸ and thus modulate the neurobiological impact of environmental trauma in children while having an adverse effect in later life. Furthermore HPA axis vulnerability to trauma may further be modulated by genetic and epigenetic factors.^{39 40} Together these observations suggest a far more complex aetiological model than we have been able to explore here due to lack of statistical power.




Figure 1 Association graph. NS, non-significant (dotted arrow).

Overall our findings would be consistent with the idea that depression may in some cases be an indicator of underlying HPA axis dysfunction or altered sensitivity due to early-life trauma which has led to associated alterations to the hippocampus. While this cannot be seen in this midlife cohort to be related to either cognitive functioning or dementia risk, hippocampal volume loss may add to the burden of later dementia-related pathology. The aetiology of depression appears in this case to be independent of that of dementia, as previously suggested by O’Brien *et al*¹⁵ in relation to postmortem studies of AD neuropathology in depressed elderly.

Our aetiological model based on cross-sectional data suggests that childhood trauma is associated with both higher rates of depression and lower hippocampal volume. Poorer performance on an episodic memory test is associated with trauma but mediated by depression rather than hippocampal volume. Thus, the relationship of depression to dementia risk in persons experiencing childhood trauma appears to be neither a risk factor nor a prodromal feature of the disease but that of a comorbidity which could potentially add to brain burden.

It is important to note, however, that only 11.3% of persons at high AD risk experienced severe early-life trauma. Thus, even if our hypothesis is correct, it will only concern a small proportion of dementia cases. There has been an underlying assumption in most previous studies of depression and AD that depression is a single clinical entity³⁻⁶ despite well-documented variability in age of onset, precipitating factors, clinical profiles, genetic risk, associated pathologies and treatment response. Our findings do not preclude the possibility of other depression subtypes being linked to AD by alternative mechanisms. The importance of this study lies in the demonstration that depressive symptomatology may be present in persons at high risk for dementia via multiple pathways, and suggest that future studies should consider more carefully the origins of depressive symptomatology when examining its relationship to dementia.

The strengths of the present study are the younger age of the dementia-risk cohort and the identification of a relatively homogeneous depression subgroup. On the other hand the data rely on subjective recall of traumatic events and lack of longitudinal data. Future studies from PREVENT will be able to further explore the hypotheses suggested in this study with a larger cohort, longitudinal data, direct measures of HPA axis functioning and AD biomarkers with a view to the identification of novel targets for therapeutic intervention in adults with a history of childhood maltreatment at high risk of dementia.

Twitter Craig W Ritchie @Craig_ritchie68

Acknowledgements The PREVENT Dementia Programme has been financed by a research grant from the UK charity, the Alzheimer's Society and Philanthropic Donations. Genotyping was carried out by Lee Murphy, Edinburgh Clinical Research Facility, University of Edinburgh.

Contributors KR designed the study and wrote the original draft of the manuscript, JTOB, LS, SD, TW, CWR and SG commented on and added to the discussion sections. IC assisted in the design of the study, carried out the statistical analyses and the writing of results.

Funding The Prevent Dementia Study has been funded by the UK Alzheimer Society Grants no. 178, 264.

Competing interests None declared.

Patient consent for publication Not required.

Ethics approval NHS Research Ethics Committee London Camberwell St-Giles (REC reference: 12/LO/1023).

Provenance and peer review Not commissioned; externally peer reviewed.

Data availability statement Data are available upon reasonable request. Data are made available to all researchers subject to approval by the Prevent Dementia Steering Group.

Supplemental material This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

ORCID iDs

Karen Ritchie <http://orcid.org/0000-0002-0688-8982>

John T O'Brien <http://orcid.org/0000-0002-0837-5080>

REFERENCES

- Kral VA. The relationship between senile dementia (Alzheimer type) and depression. *Can J Psychiatry* 1983;28:304–6.
- Engedal K, Barca ML, Laks J, et al. Depression in Alzheimer's disease: specificity of depressive symptoms using three different clinical criteria. *Int J Geriatr Psychiatry* 2011;26:944–51.
- Lyketsos CG, Olin J. Depression in Alzheimer's disease: overview and treatment. *Biol Psychiatry* 2002;52:243–52.
- Starkstein SE, Jorge R, Mizrahi R, et al. The construct of minor and major depression in Alzheimer's disease. *Am J Psychiatry* 2005;162:2086–93.
- Mourao RJ, Mansur G, Malloy-Diniz LF, et al. Depressive symptoms increase the risk of progression to dementia in subjects with mild cognitive impairment: systematic review and meta-analysis. *Int J Geriatr Psychiatry* 2016;31:905–11.
- Lebedev AV, Beyer MK, Fritze F, et al. Cortical changes associated with depression and antidepressant use in Alzheimer and Lewy body dementia: an MRI surface-based morphometric study. *Am J Geriatr Psychiatry* 2014;22:4–13.
- O'Brien J, Perry R, Barber R, et al. The association between white matter lesions on magnetic resonance imaging and noncognitive symptoms. *Ann N Y Acad Sci* 2000;903:482–9.
- Tagariello P, Girardi P, Amore M. Depression and apathy in dementia: same syndrome or different constructs? A critical review. *Arch Gerontol Geriatr* 2009;49:246–9.
- Qiu WQ, Sun X, Selkoe DJ, et al. Depression is associated with low plasma Abeta42 independently of cardiovascular disease in the homebound elderly. *Int J Geriatr Psychiatry* 2007;22:536–42.
- Namekawa Y, Baba H, Maeshima H, et al. Heterogeneity of elderly depression: increased risk of Alzheimer's disease and Aβ protein metabolism. *Prog Neuropsychopharmacol Biol Psychiatry* 2013;43:203–8.
- Sun X, Steffens DC, Au R, et al. Amyloid-Associated depression. *Arch Gen Psychiatry* 2008;65:542–50.
- Chi S, Yu J-T, Tan M-S, et al. Depression in Alzheimer's disease: epidemiology, mechanisms, and management. *J Alzheimers Dis* 2014;42:739–55.
- Ringman JM, Liang L-J, Zhou Y, et al. Early behavioural changes in familial Alzheimer's disease in the dominantly inherited Alzheimer network. *Brain* 2015;138:1036–45.
- Owby RL, Crocco E, Acevedo A, et al. Depression and risk for Alzheimer disease: systematic review, meta-analysis, and meta-regression analysis. *Arch Gen Psychiatry* 2006;63:530–8.
- O'Brien J, Thomas A, Ballard C, et al. Cognitive impairment in depression is not associated with neuropathologic evidence of increased vascular or Alzheimer-type pathology. *Biol Psychiatry* 2001;49:130–6.
- Ritchie K, Carrière I, Berr C, et al. The clinical picture of Alzheimer's disease in the decade before diagnosis: clinical and biomarker trajectories. *J Clin Psychiatry* 2016;77:e305–11.
- Malhi GS, Parker GB, Greenwood J. Structural and functional models of depression: from sub-types to substrates. *Acta Psychiatr Scand* 2005;111:94–105.
- Teicher MH, Andersen SL, Polcari A, et al. Developmental neurobiology of childhood stress and trauma. *Psychiatr Clin North Am* 2002;25:397–426.
- Heim C, Nemeroff CB. The role of childhood trauma in the neurobiology of mood and anxiety disorders: preclinical and clinical studies. *Biol Psychiatry* 2001;49:1023–39.
- Bernet CZ, Stein MB. Relationship of childhood maltreatment to the onset and course of major depression in adulthood. *Depress Anxiety* 1999;9:169–74.
- Kaufman J, Charney D. Effects of early stress on brain structure and function: implications for understanding the relationship between child maltreatment and depression. *Dev Psychopathol* 2001;13:451–71.
- Ritchie K, Jaussett I, Stewart R, et al. Adverse childhood environment and late-life cognitive functioning. *Int J Geriatr Psychiatry* 2011;26:503–10.
- Navalta CP, Polcari A, Webster DM, et al. Effects of childhood sexual abuse on neuropsychological and cognitive function in college women. *J Neuropsychiatry Clin Neurosci* 2006;18:45–53.
- Bremner JD, Vermetten E, Afzal N, et al. Deficits in verbal declarative memory function in women with childhood sexual abuse-related posttraumatic stress disorder. *J Nerv Ment Dis* 2004;192:643–9.
- Bremner JD, Randall P, Scott TM, et al. Deficits in short-term memory in adult survivors of childhood abuse. *Psychiatry Res* 1995;59:97–107.
- McKeever A, Paris AF, Cullen J, et al. Hippocampal Subfield volumes in middle-aged adults at risk of dementia. *J Alzheimers Dis* 2020;75:1211–8.
- Ritchie CW, Ritchie K. The prevent study: a prospective cohort study to identify mid-life biomarkers of late-onset Alzheimer's disease. *BMJ Open* 2012;2:e001893.
- Wolfe J, Kimerling R. Gender issues in the measurement of post-traumatic stress disorder. In: Wilson JP, Keane TM, eds. *Assessing psychological trauma and PTSD*. New York: Guilford Press, 1997.
- American Psychiatric Association. *Diagnostic and statistical manual of mental disorders*. Washington: American Psychiatric Association, 2000.
- Radloff LS. The CES-D scale: a self-report depression scale for research in the general population. *Appl Psychol Measurement* 1977;1:385–401.
- Ritchie K, de Roquefeuil G, Ritchie CW, et al. COGNITO: computerized assessment of ageing-related changes in cognitive processing. *J Psychol Psychother* 2014.
- Ashburner J. A fast diffeomorphic image registration algorithm. *Neuroimage* 2007;38:95–113.
- Tzourio-Mazoyer N, Landeau B, Papathanassiou D, et al. Automated anatomical labeling of activations in SPM using a macroscopic anatomical parcellation of the MNI MRI single-subject brain. *Neuroimage* 2002;15:273–89.
- Kessler RC, Birnbaum HG, Shahly V, et al. Age differences in the prevalence and comorbidity of DSM-IV major depressive episodes: results from the who world mental health survey initiative. *Depress Anxiety* 2010;27:351–64.
- Sjöberg L, Karlsson B, Atti A-R, et al. Prevalence of depression: comparisons of different depression definitions in population-based samples of older adults. *J Affect Disord* 2017;221:123–31.
- Moceri VM, Kukull WA, Emanuel I, et al. Using census data and birth certificates to reconstruct the early-life socioeconomic environment and the relation to the development of Alzheimer's disease. *Epidemiology* 2001;12:383–9.
- Savitz J, van der Merwe L, Stein DJ, et al. Genotype and childhood sexual trauma moderate neurocognitive performance: a possible role for brain-derived neurotrophic factor and apolipoprotein E variants. *Biol Psychiatry* 2007;62:391–9.
- Poirier J. Apolipoprotein E in animal models of CNS injury and in Alzheimer's disease. *Trends Neurosci* 1994;17:525–30.
- Ancelin M-L, Scali J, Norton J, et al. The effect of an adverse psychological environment on salivary cortisol levels in the elderly differs by 5-HTTLPR genotype. *Neurobiol Stress* 2017;7:38–46.
- Wrigglesworth J, Ancelin M-L, Ritchie K, et al. Association between DNA methylation of the KITLG gene and cortisol levels under stress: a replication study. *Stress* 2019;22:162–8.

Table S1: Comparison between participants with MRI imaging and those without MRI imaging

	Brain MRI		P-value ^a
	No	Yes	
	(n=185)	(n=193)	
	n (%)	n (%)	
Sex, female	128 (69.2)	135 (70.0)	0.87
Marital status			0.30
Married or living as a couple	135 (73.0)	149 (77.6)	
Educational level			0.002
College / university	141 (76.6)	120 (62.2)	
Living alone	27 (14.6)	27 (14.0)	0.87
Socio-economic class, n=358			0.007
- Higher managerial, administrative occupations	47 (27.5)	42 (22.5)	
- Lower managerial, administrative occupations	73 (42.7)	66 (35.3)	
- Intermediate occupations	21 (12.3)	27 (14.4)	
- Employers, own account, technical, routine occupation	23 (13.4)	23 (12.3)	
- Never worked or long-term unemployed	7 (4.1)	29 (15.5)	
Depressive symptomatology (CES-D \geq 16 or treated)	49 (26.5)	44 (22.8)	0.41
Parental history of dementia, n=377	108 (58.7)	95 (49.2)	0.07
BMI (kg/m ²)			0.09
Normal (<25)	75 (40.5)	66 (34.2)	
Overweighted (25-30)	61 (33.0)	85 (44.0)	
Obese (\geq 30)	49 (26.5)	42 (21.8)	
Hypertension (treated or SBP>140)	38 (20.5)	28 (14.5)	0.12
Cardiovascular disease (declared)	12 (6.5)	22 (11.4)	0.10
Diabetes (treated or declared or glycemia>7mmol/l), n=377	8 (4.4)	8 (4.2)	0.92

Anxiety symptoms (Spielberger state scale >31)	60 (32.4)	62 (32.1)	0.95
Traumatic events before 16 years (at least one)	23 (12.4)	31 (16.1)	0.31
	Median (IQR)	Median (IQR)	P-value ^b
Age (years)	53 (48-56)	53 (48-56)	0.72

Table S2. Association between depression and current cognitive performance

Dependent variables	Depressive symptomatology			Depressive symptomatology		
	CES-D<16 and not treated (n=285)	CES-D≥16 or treated (n=93)	P- value	CES-D<16 and not treated (n=284)	CES-D≥16 or treated (n=93)	P- value
	Mean(SEM) ^a	Mean (SEM) ^a		Mean(SEM) ^b	Mean (SEM) ^b	
Working memory: time difference between double task and simple task (milliseconds)	-129.6 (180.4)	242.0 (315.9)	0.31	-121.3 (206.7)	214.4 (327.0)	0.36
Form matching: mean time for correct responses (milliseconds)	5713.0 (89.5)	5851.3 (156.6)	0.44	5685.1 (102.5)	5828.5 (162.1)	0.43
Phoneme comprehension: mean time for correct responses (milliseconds)	1498.8 (17.7)	1488.5 (31.0)	0.75	1511.1 (20.0)	1500.7 (31.7)	0.77
Name-face associations: number of names correctly recalled	5.5 (0.13)	4.9 (0.22)	0.03	5.1 (0.13)	4.6 (0.21)	0.03
Verbal fluency: number of correct answers	28.8 (0.38)	27.8 (0.66)	0.18	28.0 (0.42)	27.2 (0.67)	0.26
Narrative recall: number of correct answers	13.3 (0.26)	12.8 (0.45)	0.31	12.9 (0.29)	12.5 (0.46)	0.46
Descriptive recall: number of correct answers	13.1 (0.26)	12.3 (0.45)	0.14	12.5 (0.28)	11.9 (0.44)	0.22

Implicit Memory: difference between learnt and new items	1.02 (0.04)	0.99 (0.08)	0.72	1.07 (0.05)	1.02 (0.08)	0.58
----------------------------------------------------------	-------------	-------------	------	-------------	-------------	------

^a Non adjusted.

^b Adjusted for education level, gender and age.

Table S3. Association between depression and current categorical cognitive performances

	Depressive symptomatology		P- value ^a	P- value ^b
	CES-D<16 and not treated (n=285) n (%)	CES-D≥16 or treated (n=93) n (%)		
Form matching, <7 correct responses	144 (50.5)	54 (58.1)	0.21	0.19
Phoneme comprehension, <9 correct responses	85 (29.8)	35 (37.6)	0.16	0.18

^a Non adjusted, Chi squared test.

^b Adjusted for education level, gender and age, logistic regression.

Table S4. Associations between hippocampal volume as the dependent variable and cognitive scores (separate linear models), n=193

Dependent variables	Hippocampal gray matter volume (Adjusted cognitive change for one cm ³)	
	β (SE) ^a	p-value ^a
Linear model		
Working memory: time difference between double task and simple task (milliseconds)	490 (1188)	0.68
Form matching: mean time for correct responses (milliseconds)	-0.80 (536)	0.99
Phoneme comprehension: mean time for correct responses (milliseconds)	40.37 (110)	0.72
Name-face associations: number of names correctly recalled	0.17 (0.76)	0.82
Verbal fluency: number of correct answers	-0.26 (2.27)	0.91
Narrative recall: number of correct answers	0.40 (1.71)	0.81
Descriptive recall: number of correct answers	0.56 (1.47)	0.70
Implicit Memory: difference between learnt and new items	-0.12 (0.32)	0.71
Logistic regression	OR (95%IC)	p-value ^a
Form matching, <7 correct responses	0.55 (0.13; 2.39)	0.43
Phoneme comprehension, <9 correct responses	2.82 (0.60;13.33)	0.19

^a Adjusted for total intracranial volume, education, gender and age

