



Blood-based high sensitivity measurements of beta-amyloid and phosphorylated tau as biomarkers of Alzheimer's disease: a focused review on recent advances

Joyce R. Chong,^{1,2} Nicholas J. Ashton,^{3,4,5,6} Thomas K. Karikari,^{6,7} Tomotaka Tanaka,^{1,8,9} Michael Schöll,^{3,6,10} Henrik Zetterberg ,^{6,10,11,12} Kaj Blennow,^{6,12} Christopher P. Chen,^{1,2} Mitchell K.P. Lai ^{1,2}

► Additional supplemental material is published online only. To view, please visit the journal online (<http://dx.doi.org/10.1136/jnnp-2021-327370>).

For numbered affiliations see end of article.

Correspondence to

Dr Mitchell K.P. Lai, Department of Pharmacology, National University of Singapore, Unit 09-01, 14 Medical Drive, Kent Ridge 117599, Singapore; Mitchell.lai@dementia-research.org

Received 13 June 2021
Accepted 30 August 2021
Published Online First 11 September 2021

ABSTRACT

Discovery and development of clinically useful biomarkers for Alzheimer's disease (AD) and related dementias have been the focus of recent research efforts. While cerebrospinal fluid and positron emission tomography or MRI-based neuroimaging markers have made the in vivo detection of AD pathology and its consequences possible, the high cost and invasiveness have limited their widespread use in the clinical setting. On the other hand, advances in potentially more accessible blood-based biomarkers had been impeded by lack of sensitivity in detecting changes in markers of the hallmarks of AD, including amyloid- β (A β) peptides and phosphorylated tau (P-tau). More recently, however, emerging technologies with superior sensitivity and specificity for measuring A β and P-tau have reported high concordances with AD severity. In this focused review, we describe several emerging technologies, including immunoprecipitation-mass spectrometry (IP-MS), single molecule array and Meso Scale Discovery immunoassay platforms, and appraise the current literature arising from their use to identify plaques, tangles and other AD-associated pathology. While there is potential clinical utility in adopting these technologies, we also highlight the further studies needed to establish A β and P-tau as blood-based biomarkers for AD, including validation with existing large sample sets, new independent cohorts from diverse backgrounds as well as population-based longitudinal studies. In conclusion, the availability of sensitive and reliable measurements of A β peptides and P-tau species in blood holds promise for the diagnosis, prognosis and outcome assessments in clinical trials for AD.

memory deficits as well as non-amnestic cognitive symptoms, including impairments in language, visuospatial and executive function that severely impact activities of daily living. Neuropathologically, AD is characterised by progressive synaptic dysfunction and neuronal loss in brain regions that are essential for higher cognitive functions,³ culminating in brain atrophy and clinical symptoms in patients. The hallmarks of AD include cortical extracellular amyloid- β (A β) plaques composed of highly aggregated, fibrillar 40-amino-acid to 42-amino-acid A β peptides (A β 40 and A β 42), as well as intracellular neurofibrillary tangles (NFTs) and dystrophic neurites surrounding the plaques, both composed of paired helical filament-forming, abnormally hyperphosphorylated tau protein aggregates.² As neurodegeneration is thought to be well underway by the time patients exhibit clinical symptoms,⁴ these core pathophysiological features have been targets for AD biomarker development—a key area in dementia research that holds promise for early detection of the disease which in turn helps improve the drug development process for AD (see later). While *postmortem* neuropathological evaluation remains the gold standard for confirmatory diagnosis of AD pathology, the core AD diagnostic biomarkers in living individuals with positron emission tomography (PET assessment of in vivo A β and tau cortical burden) or cerebrospinal fluid (CSF measurements of A β 42, tau and P-tau) analyses have gained acceptance^{5,6} but remain relatively inaccessible.

APPLICATIONS AND CHALLENGES OF BLOOD-BASED AD BIOMARKERS

Shift towards a biological definition of AD

Historically, AD was conceived of as a clinical-pathological construct, such that cognitive symptoms defined the presence of AD.⁷ However, approximately 30% of individuals clinically diagnosed as AD do not display significant AD neuropathological changes at autopsy or by in-vivo imaging.⁸ The National Institute on Aging and Alzheimer's Association (NIA-AA) research framework has recently suggested that AD should be defined as a biological construct using biomarkers

INTRODUCTION

There are over 50 million people worldwide suffering from dementia, a number which is projected to triple by 2050.¹ The high prevalence of dementia is accompanied by a massive social and economic burden; the current annual cost of dementia is estimated at US\$1 trillion and is set to double by 2030.¹ The most common cause of dementia is Alzheimer's disease (AD), which accounts for approximately 60%–80% of all dementia cases.² AD is characterised by progressive



© Author(s) (or their employer(s)) 2021. No commercial re-use. See rights and permissions. Published by BMJ.

To cite: Chong JR, Ashton NJ, Karikari TK, et al. *J Neurol Neurosurg Psychiatry* 2021;**92**:1231–1241.

that are characteristic of AD pathophysiology, such as A β and phosphorylated tau (P-tau).^{8,9} This biological definition of AD is potentially useful in both research and clinical care settings. In the former, it facilitates efforts to understand the sequence of events leading to cognitive impairment that is associated with AD neuropathology. In the latter, it allows clinicians to confirm the aetiological diagnosis with greater certainty and provide biomarker-guided targeted therapies when they are available.⁷ Furthermore, it improves early identification (screening) of older people at risk of developing AD, providing access to important patient cohorts crucial for the clinical evaluation of promising therapeutic strategies of those with significant burden of pathological proteins.

Recruitment of appropriate trial subjects for disease-modifying therapies

Given that current AD treatments provide only symptomatic relief, there has been a focus on clinical trials of disease-modifying therapies (DMT), targeting specific AD pathologies, in particular amyloidosis, but recently also tau pathology. In DMT trials, it is imperative to show that an intervention modifies the target pathophysiological process and improves clinical symptoms.⁸ Disappointingly, past DMT trials have largely reported no or minimal cognitive benefits.¹⁰ A possible explanation is the inclusion of participants lacking the target pathology since recruitment was based mainly on clinical assessment. Another possibility is the inclusion of individuals too advanced in the disease process, with a clinical course which may be difficult to modify regardless of the efficacy of the drug candidates. Thus, it is necessary to conduct biomarker-driven subject selection in DMT trials to facilitate target engagement and reliably assess the efficacy of interventions.^{9–11} An example of such a trial is a phase II clinical study of BAN2401 (also called lecanemab) in early AD subjects with positive biomarkers for brain amyloid pathology.¹² BAN2401 is a humanised monoclonal antibody that selectively binds to, neutralises and eliminates soluble toxic A β aggregates. BAN2401 showed a significant, dose-dependent reduction in PET A β such that approximately 80% of participants demonstrated reduced A β aggregate densities and were reclassified from A β positive to negative (based on amyloid PET visual read) at 18 months with the highest tested dose. Importantly, participants also showed slower cognitive decline.^{13,14} Given these promising results, a global multi-centre phase III clinical study of BAN2401 (Clarity AD) has been initiated. Other examples include the phase III trials, ENGAGE and EMERGE, which aimed to assess the cognitive and functional impacts of the aggregated A β -binding monoclonal antibody drug aducanumab in individuals with mild cognitive impairment (MCI) due to early AD as determined by a positive PET A β scan. In a report at the end of 2019, it was noted that at the highest dose, participants in EMERGE had a significant reduction in the rate of cognitive decline using a standardised neuropsychological test (Clinical Dementia Rating-Sum of Boxes, CDR-SB).¹⁵ Similarly, in ENGAGE, a subgroup of participants who received the highest dose at least 10 times declined more slowly on the CDR-SB. In substudies, aducanumab which has been approved by the United States Food and Drug Administration as of June 2021, caused a dose-dependent reduction in brain amyloid burden. Overall, these clinical trials point to the importance of trial recruitment with supporting biomarker evidence to ensure that the appropriate participants are enrolled, and to provide evidence of the efficacy of the tested drugs in targeting AD pathology.

Current challenges in the use of AD biomarkers

While AD biomarkers measured by PET or CSF are highly indicative of AD pathophysiology, the challenges involved in their usage, including invasiveness of procedures, high cost, limited accessibility to scanners and cyclotrons, and limited utility as a screening tool, have impeded their widespread application in the clinical and research settings.^{3,7} In contrast, blood-based biomarkers would be a readily accepted and practical approach if they had sensitivity and specificity comparable to neuroimaging and CSF markers.^{3,9} However, most early studies on blood A β 42, A β 40 and A β 42/A β 40 ratio using conventional ELISA or multi-analyte Luminex immunoassays reported little or no difference between AD and control groups.¹⁶ Furthermore, the majority of these early studies were based on comparing blood A β in clinically diagnosed patients with AD with cognitively unimpaired controls which, given the uncertain value of clinical assessments in assessing pathological burden, might have underestimated and confounded the diagnostic value of plasma A β .^{9,16} The specific limitations of conventional blood A β measurements include the low abundance of A β in peripheral blood which may result from filtering effects of the blood–brain barrier, dilution in the large plasma volume, rapid metabolism and clearance, and adhesion to other plasma proteins leading to epitope masking and analytical interference. As a consequence, early measurements of A β concentrations in blood were much lower than those in CSF.^{3,7,9} This low abundance impeded reliable, quantitative measurements of peripheral A β in earlier studies using conventional immunoassays with their inherent limits in sensitivity.

Furthermore, measurement of soluble A β in the peripheral blood may not be reflective of brain amyloid plaque burden, as indicated by the poor correlation between blood and CSF A β levels.^{17,18} This limitation may have been compounded by early studies using assays which measured only soluble A β in the blood and could not recognise aggregated isoforms, leading to under-detection of A β oligomers (A β Os) that are widely postulated to be the major neurotoxic forms of A β .^{19–21} The ability to measure specific A β isoforms such as A β Os could therefore be more pathophysiologically relevant to brain amyloid burden.

Emerging technologies for blood-based amyloid measurements

In recent years, there has been a growing impetus for the development of new platforms and immunoassays for measuring peripheral A β . These developments helped address the major challenges listed earlier by having improved sensitivity compared with conventional assays, and/or the ability to measure A β variants that may be more relevant to AD pathology.⁹ The following section aims to summarise these emerging techniques or technologies, with brief descriptions of the assay principles followed by appraisals of their ability to identify individuals with significant brain A β burden (A β +) compared with those without (A β –, as confirmed by brain PET or CSF analyses). A summary of the results is provided in [table 1](#) (with detailed cohort information given in online supplemental table S1). Group comparisons between A β – and A β – are presented in [table 1](#). To assess the diagnostic performance of blood A β in detecting elevated A β , area under the receiver operating characteristic curve (AUC) values were given where reported ([table 1](#)). For associations between blood A β and continuous variables such as the standardised uptake value ratio (SUVR) and CSF A β levels, correlation or regression results was considered (online supplemental table

Table 1 Association between blood amyloid and brain amyloid pathology

Result	Cohort details	Technology type	AUC (Aβ+ vs Aβ-)	Ref
↓ Plasma Aβ42 in Aβ+ group vs Aβ- group*	22 Aβ-, 40 Aβ+ (PET)	IP-MS	0.81	28
	Cohort 1 (NCGG): 71 Aβ-, 50 Aβ+ (PET)	IP-MS	Cohort 1: 0.87 (0.91†)	29
	Cohort 2 (AIBL): 115 Aβ-, 137 Aβ+ (PET)		Cohort 2: 0.72 (0.80†)	
			Cohort 1 and 2: 0.79	
	359 Aβ-, 82 Aβ+ (PET)	IP-MS	0.74 (0.79‡)	30
	46 Aβ-, 23 Aβ+ (PET)	Simoa	0.66	25
	191 Aβ-, 57 Aβ+ (CSF)	Simoa	0.66	25
	359 Aβ-, 82 Aβ+ (PET)	Simoa	0.59 (0.71‡)	30
↓ Plasma Aβ42 in Aβ+ CU, Aβ+ SCD, Aβ+ MCI, Aβ+ AD vs Aβ- CU, Aβ- MCI	CU (200 Aβ-, 74 Aβ+), 60 Aβ+ SCD, 121 Aβ+ MCI, 53 Aβ+ AD (CSF)	Simoa		18
	CU (366 Aβ-, 147 Aβ+), MCI (108 Aβ-, 157 Aβ+), 64 Aβ+ AD (CSF)	Elecsys		35
↑ Plasma exosome bound Aβ42 in Aβ+ group vs Aβ- group*	45 Aβ-, 23 Aβ+ (PET)	APEX	0.99	40 and 41
No sig. difference in plasma Aβ42 between Aβ+ vs Aβ- groups*	10 Aβ-, 10 Aβ+ (PET)	MSD		37
↓ Plasma Aβ42/Aβ40 ratio in Aβ+ group vs Aβ- group*	22 Aβ-, 40 Aβ+ (PET)	IP-MS	0.80	28
	23 Aβ-, 18 Aβ+ (PET or CSF)	IP-MS	0.89	31
	115 Aβ-, 43 Aβ+ (PET)	IP-MS	0.88 (0.94§)	22
	359 Aβ-, 82 Aβ+ (PET)	IP-MS	0.82 (0.84‡)	30
	161 Aβ- and 38 Aβ+ (PET)	Simoa	0.79 (0.81§)	23
	191 Aβ-, 57 Aβ+ (CSF)	Simoa	0.77 (0.83§)	25
	359 Aβ-, 82 Aβ+ (PET)	Simoa	0.62 (0.73‡)	30
	45 Aβ-, 23 Aβ+ (PET)	Simoa	0.82	41
	474 Aβ-, 368 Aβ+ (CSF)	Elecsys	0.77 (0.80¶, 0.85**)	35
	161 Aβ- and 38 Aβ+ (PET)	ELISA	0.78 (0.78§)	23
↓ Plasma Aβ42/Aβ40 ratio in Aβ+ CU, Aβ+ SCD, Aβ+ MCI, Aβ+ AD vs Aβ- CU, Aβ- MCI	CU (200 Aβ-, 74 Aβ+), 60 Aβ+ SCD, 121 Aβ+ MCI, 53 Aβ+ AD (CSF)	Simoa		18
	CU (366 Aβ-, 147 Aβ+), MCI (108 Aβ-, 157 Aβ+), 64 Aβ+ AD (CSF)	Elecsys		35
No sig. difference in plasma Aβ42/Aβ40 ratio between Aβ+ vs Aβ- groups*	46 Aβ-, 23 Aβ+ (PET)	Simoa	0.68 (0.79§)	25
	32 Aβ-, 8 Aβ+ (PET)	Simoa		87
	10 Aβ-, 10 Aβ+ (PET)	MSD		37
↑ Plasma Aβ40/Aβ42 ratio in Aβ+ group vs Aβ- group*	Cohort 1 (NCGG): 71 Aβ-, 50 Aβ+ (PET)	IP-MS	Cohort 1: 0.97 (0.98†)	29
	Cohort 2 (AIBL): 115 Aβ-, 137 Aβ+ (PET)		Cohort 2: 0.84 (0.85†)	
			Cohort 1 and 2: 0.89	
↑ Plasma APP669-711/Aβ42 ratio in Aβ+ group vs Aβ- group*	22 Aβ-, 40 Aβ+ (PET)	IP-MS	0.97	28
	Cohort 1: 71 Aβ-, 50 Aβ+ (PET)	IP-MS	Cohort 1: 0.92 (0.93†)	29
	Cohort 2: 115 Aβ-, 137 Aβ+ (PET)		Cohort 2: 0.83 (0.85†)	
			Cohort 1 and 2: 0.86	
↑ Plasma composite biomarker in Aβ+ group vs Aβ- group* (composite=average of the normalised values of APP669-711/Aβ42 ratio and Aβ40/Aβ42 ratio)	Cohort 1: 71 Aβ-, 50 Aβ+ (PET)	IP-MS	Cohort 1: 0.97 (0.97†)	29
	Cohort 2: 115 Aβ-, 137 Aβ+ (PET)		Cohort 2: 0.88 (0.89†)	
			Cohort 1 and 2: 0.91	
	359 Aβ-, 82 Aβ+ (PET)	IP-MS	0.82 (0.84‡)	30

*For comparison between Aβ+ and Aβ- groups, participants of different clinical diagnoses are included.

†AUC after adjusted for age, gender, Apolipoprotein E genotype (APOE), clinical category.

‡AUC after adjusted for age, gender, APOE.

§AUC after adjusted for age, APOE.

¶AUC for inclusion of plasma Aβ42 and Aβ40 as separate predictors in logistic regression.

**AUC for inclusion of plasma Aβ42, Aβ40 and APOE as separate predictors in logistic regression.

††AUC after adjusted for age, gender.

Aβ, amyloid-β; AD, Alzheimer's disease; AIBL, Australian Imaging, Biomarker and Lifestyle Study of Ageing; AUC, area under the receiver operating characteristic curve; CSF, cerebrospinal fluid; CU, Cognitively unimpaired; IP-MS, immunoprecipitation-mass spectrometry; MCI, mild cognitive impairment; MSD, Meso Scale Discovery; NCGG, Japanese National Center for Geriatrics and Gerontology; PET, positron emission tomography; SCD, Subjective cognitive decline; Simoa, single molecule array.

S1). Taken together, these data show blood Aβ biomarkers demonstrate improved discrimination performance when participants were classified into amyloid positive or negative based on PET or CSF amyloid measurements, compared with classifications based on clinical assessments. Nevertheless, blood Aβ typically showed much lower fold change (10%–15%) in plasma compared with CSF (40%–50%), likely due

in part to a large proportion of plasma Aβ coming from peripheral sources of the peptide.²²

Next generation ELISAs

As mentioned, early generation ELISAs were largely unable to accurately detect and measure AD-associated changes in blood

A β . However, significant improvements have been made in newer ELISAs to enhance their performance. For instance, the EUROIMMUN ELISA study²³ employed C-terminal and N-terminal antibodies, while most previous ELISAs did not, and consequently detected different A β fragments. Furthermore, improvements in assay design and conjugation method may also have resulted in increased sensitivities, thus enabling the detection of more subtle variations in biomarker profiles than attainable with early generation ELISAs. These improvements likely led to similar performance between the EUROIMMUN ELISA and the ultrasensitive single molecule array (Simoa) ('Amyblood') assay (see later) in a head-to-head comparison, where the accuracy of detecting PET A β + in non-demented participants through measurements of plasma A β 42/A β 40 ratio was similar, and plasma A β 42/A β 40 as well as A β 42/t-tau ratios measured by both methods correlated to the same extent with amyloid PET and CSF A β 42/t-tau.²³ Nevertheless, more head-to-head comparisons among the different available ELISAs are needed.

Simoa immunoassay

The Simoa immunoassay is a digital ELISA, allowing concentrations to be determined digitally rather than by measurement of the total analogue signal (Quanterix, USA). Briefly, after formation of the immunocomplex on paramagnetic beads, the beads are transferred to thousands of femtoliter-sized wells, each sized to hold only one bead. By confining the fluorophores generated by individual enzymes to extremely small volumes (approximately 40 fL), a high local concentration of fluorescent products is achieved. Therefore, a readable signal is detected even if only a single sandwich complex is present on the bead. The analyser counts the number of wells containing an enzyme-labelled bead and wells containing a bead. The ratio between the counts provides the average enzyme per bead number, and the concentration of the target analyte in the sample is derived from a standard curve.

The first version of the Simoa assay for blood A β was published in 2011.²⁴ Using this assay, two large, independent studies, Janelidze *et al*¹⁸ and Verberk *et al*,²⁵ reported no or weak correlations between the plasma A β biomarkers and established measurements such as SUVR, CSF A β 42 or CSF A β 42/A β 40 ratio ($r=0.16$ to 0.38). In distinguishing between PET A β + and A β - subjects, Janelidze *et al* reported a modest AUC of 0.60 – 0.62 for the plasma A β biomarkers, while Verberk *et al* reported numerically higher AUCs of 0.66 – 0.79 . Likewise, in differentiating between CSF A β + and A β - subjects, Janelidze *et al* reported AUCs of 0.66 – 0.68 for the plasma A β biomarkers, while Verberk *et al* reported AUCs of 0.66 – 0.83 . The varying reported ranges may be due to differences in the cohorts used, adjustments of AUCs (for age, gender and/or apolipoprotein E genotype) or the combination of detector-capture antibody pairs used. Janelidze *et al* used the 'first-generation' singleplex from Quanterix which employed N-terminal antibody 6E10, which binds to the RHD sequence located at residues 5–7 of the A β peptide, as the capture antibody and isoform-specific C-terminal antibodies, namely H31L21 and 2G3 for A β 42 and A β 40, respectively, as the detector antibodies. However, such antibody pair combination may result in lower assay specificity.²³ In contrast, Verberk *et al* used the Neurology 3-Plex which swapped the antibodies that are used to capture and detect plasma A β .²⁵ Matching the antibody orientation of Verberk *et al*, a small-scale study by Thijssen *et al*²⁶ selected the C-terminal antibodies 21F12 and 2G3 as capture antibodies for A β 42 and A β 40, respectively, and

N-terminal antibody 3D6 that binds to the first five residues of the A β peptide, as the detector antibody. When compared with the commercially available assay—which used an N-terminal antibody to capture both peptides, followed by 21F12 or 2G3 for detection of A β 42 and A β 40, respectively—the prototype assay performed better in discriminating between A β + and A β - subjects (AUC 0.95 vs 0.85) and produced a stronger correlation between plasma A β 42/A β 40 ratio and CSF A β 42 ($r=0.71$ vs $r=0.53$) (online supplemental table S1). Overall, these studies indicated that while Simoa provides superior sensitivity, the combination of detector-capture antibody pairs should be carefully considered to maximise assay performance.

A major advantage of the Simoa immunoassay is its increased sensitivity when compared with conventional immunoassays, measuring proteins at femtomolar concentration and allowing detection of low abundance targets such as peripheral A β . Moreover, the assay may be performed in a fully automated setup, increasing efficiency and output, and minimising variability of results common in a manual setup. The scalability of this platform implies that it is highly adaptable to clinical care and clinical trial settings. However, as discussed earlier, further optimisation to the assays to validate its sensitivity, specificity and consistency is warranted. Besides the diagnostic utility, Simrén *et al* showed that longitudinal change in Simoa A β 42/A β 40 measures were sensitive to grey matter loss in cognitively unimpaired individuals,²⁷ suggesting the potential prognostic utility of Simoa measured A β .

Immunoprecipitation-mass spectrometry

This technique involves the isolation and enrichment of A β peptides from plasma by immunoprecipitation (IP) using a specific antibody, followed by identification and relative quantification of the individual A β isoforms including A β 42, A β 40, A β 38 or APP669-711 with highly sensitive mass spectrometry (MS).^{28–32} While the principle underlying this approach is similar for the studies, there are key differences in methodologies. Kaneko *et al*²⁸ and Nakamura *et al*²⁹ employed matrix-assisted laser desorption/ionisation-time-of-flight MS (MALDI-TOF/MS), whereas Ovod *et al*³¹ and Keshavan *et al*³⁰ used liquid chromatography tandem mass spectrometry MS (LC-MS/MS). Pannee *et al*³² used MALDI-TOF/MS A β profiling followed by LC-MS/MS to confirm the identities of the A β peptides. In comparison with LC-MS/MS which generates multiple mass spectra for each fragmented peptide and compares these spectra to a database for protein identification, in MALDI-TOF/MS only one MS spectrum is generated, compiling the masses of all peptides. As such, MALDI-TOF/MS provides a quicker analysis and higher throughput, making it more feasible for large-scale studies. However, MALDI-TOF/MS is less accurate and reliable in protein identification due to the lack of true sequence dependence of data since only a mass spectrum is generated for all peptides.

Next, the studies have selected different antibodies and number of rounds of IP, which may affect the overall specificity and sensitivity of the assay. In addition to A β 42 and A β 40, Kaneko *et al* and Nakamura *et al* measured APP669-711 (A β 3–40) while Pannee *et al* and Ovod *et al* measured A β 38 (APP672-709). The purpose of these measurements was to correct for inter-individual variability in the overall metabolic production of A β .³³ Similar to A β 40, both APP669-711 and A β 38 are thought to increase due to higher overall A β production, but are unaffected by AD pathology.³³ Thus, the ratio of each individual isoforms and A β 42 may serve as a better predictor of

brain amyloid burden than A β 42 alone. For instance, CSF A β 42/A β 38 ratio was reported to be better than CSF A β 42 alone at predicting PET A β positivity, with a performance that is comparable to CSF A β 42/A β 40 ratio.³⁴

The published AUC results suggested that plasma APP669-711/A β 42 ratio (AUC \geq 0.86), A β 42/A β 40 ratio (AUC \geq 0.80) or A β 40/A β 42 ratio (AUC \geq 0.87) was able to discriminate between A β + and A β - subjects. In their study, Nakamura *et al* evaluated the classification ability (A β + vs A β -) of a composite biomarker—an average of the normalised values of APP669-711/A β 42 ratio and A β 40/A β 42 ratio. The composite biomarker showed the highest classification ability (AUC \geq 0.91) among the A β biomarkers. A later study by Keshavan *et al* measured and compared the ability of three blood-based candidates, namely LC-MS measures of plasma A β and Simoa-measures of plasma A β and tau phosphorylated at threonine 181 (P-tau181), to detect A β PET status in a population-based cohort of non-dementia individuals.³⁰ In this study, the LC-MS plasma composite biomarker (AUC=0.82) showed the highest AUC among other plasma biomarkers, including LC-MS A β 42 (AUC=0.74), Simoa A β 42 (AUC=0.59) and A β 42/A β 40 ratio (AUC=0.62). The plasma LC-MS A β 42/A β 40 ratio showed an AUC of 0.82. Although the composite biomarker showed the highest classification ability among the A β biomarkers, there was reservation regarding the interpretation of the data. A composite biomarker should be derived from two independent markers but APP669-711/A β 42 ratio and A β 40/A β 42 ratio are not independent—A β 42 was accounted for two times in the formula and being the most impactful on the analyses, it could potentially skew the overall results and conclusion.

A major advantage of IP-MS is the consistency of the Receiver Operating Characteristic (ROC) results across different studies, despite differences in methodology. All studies reported high AUCs for the combined ratios of A β 42 with A β 40 or APP669-711, indicating the superior ability of these ratios to detect elevated brain A β . Furthermore, this approach allows for the detection of multiple A β -related peptides in human plasma such as APP669-711, for which there are currently no commercially available antibody-based assay kits. Nonetheless, there are limitations to IP-MS which may reduce feasibility for widespread clinical use, such as complexity of the procedures, long processing times and inconsistency in analytical procedures across different diagnostic labs. Development of an automated assay system to standardise the analytic factors and increase throughput is needed to support its usage in a larger setting.

Elecsys immunoassays

The Elecsys immunoassay is an electrochemiluminescence (ECL) method using a sandwich principle performed in a fully automated setup (Roche Diagnostics). First, samples are mixed with biotinylated antibodies specific to the target A β isoforms and ruthenium-labelled detection antibodies, resulting in the formation of a sandwich complex. Next, streptavidin-coated magnetic beads are added which bind to the biotinylated antibodies. For measurement, the reaction mixture is aspirated into a measuring cell where the beads are magnetically captured onto the surface of an electrode. To start the reaction, voltage is applied to the electrode, leading to emission of photons from the ruthenium complex, which is measured by a photomultiplier. The analyte concentration is determined via a calibration curve.

Using CSF A β 42/A β 40 ratio as a marker of A β status against which plasma A β was compared, Palmqvist *et al* reported a decrease in Elecsys-measured plasma A β 42/A β 40 ratio in A β +

subjects,³⁵ and there was also a marked decrease in plasma A β 42 and A β 40, especially in AD dementia subjects.³⁵ The plasma A β 42, A β 40 and A β 42/A β 40 ratio predicted A β positivity with an AUC of 0.71, 0.54 and 0.77, respectively. When using plasma A β 42 and A β 40 as separate predictors in a logistic regression, the AUC improved slightly to 0.80, and further increased to 0.85 after adding APOE genotype into the model. As APOE4 allele carriers have significantly lower CSF A β 42 levels, the combination of APOE genotype and blood-based biomarkers could further support the diagnosis of AD.³⁶ In the study, a moderate positive correlation ($r=0.48$) was reported between CSF and plasma A β 42/A β 40 ratios. The inclusion of APOE genotype in a combined model (A β 42 and A β 40) significantly improved the AUC, suggesting that future biomarkers studies may consider the inclusion of AD risk factors (eg, age, gender, APOE) to improve prediction.^{25,29} The current results are promising and warrant further validation in independent cohorts to determine the utility of Elecsys-measured plasma A β for the detection of elevated brain A β .

Meso Scale Discovery platform

Similar to the Elecsys immunoassay, the Meso Scale Discovery (MSD) platform is an ECL-based immunoassay (Meso Scale Diagnostics) which allows multiplexing. The MSD plate has a working electrode surface where specific capture antibodies are immobilised on independent and well-defined spots. Depending on the capture antibody immobilised on each spot, multiple target analytes may be captured from the sample. Furthermore, both techniques use different ECL labels and antibody pairs, which may impact on their sensitivity and specificity. Plates containing samples, capture antibody and detection antibody conjugated with MSD SULFO-TAG labels are loaded into the MSD instrument where a voltage is applied to the plate electrodes, causing the captured labels to emit light. Multiple excitation cycles amplify signals to increase the sensitivity of the assay and analyte concentrations are determined via a calibration curve.

Using PET for A β status, Vogelgsang *et al* reported no significant difference in plasma A β 42 and A β 42/A β 40 ratio between A β + and A β - groups.³⁷ The study also showed a moderate but positive correlation between CSF and plasma A β 42/A β 40 ratios ($r=0.43$), comparable to results reported for Elecsys (see earlier). This is unsurprising given the similar underlying assay principles. Currently, there is little evidence supporting the use of MSD platform in the measurement of plasma A β as a blood-based biomarker. Since the study cohort was small ($n=41$), future studies using a larger cohort may provide more indication on the potential use of this platform.

Immunomagnetic reduction

Immunomagnetic reduction (IMR) quantifies the concentrations of target proteins/peptides in a sample by measuring the percentage reduction in the alternating current magnetic susceptibility of the IMR reagent caused by the binding of the antibody-coated magnetic nanobeads with the targets (MagQu, Taiwan ROC). When the IMR reagent which contains the antibody-coated magnetic beads is mixed with the sample, the target analytes bind with the antibodies and part of the magnetic beads become enlarged, leading to a reduction in the oscillation speed and magnetic signal of the beads. The reduction percentage is referred to as the IMR signal. Sample analyte concentration is then calculated based on the established relationship between protein standard concentration and IMR signal.

In contrast to other immunoassays, IMR-measured plasma A β 42 was increased in AD dementia.³⁸ A possible explanation is that other immunoassays are mostly based on sandwich ELISA which relies on the binding of both capture and detection antibodies to measure plasma A β . Since A β is frequently bound to plasma proteins, this may induce a potential stereoscopic obstacle for two antibodies to associate with one A β molecule simultaneously, leading to partial loss of signal. In comparison, the IMR method uses a single antibody to capture A β molecule in the plasma. Therefore, the IMR method has a higher possibility of capturing and detecting the target A β molecules in various conformations, such as isolated, complex or oligomeric forms.

Two IMR-based studies have reported on the association of plasma A β with brain A β burden, with Tzen *et al*³⁹ showing a significant association between IMR-measured plasma A β 42/A β 40 ratio and PET SUVR ($\beta=0.65$) while Teunissen *et al*³⁸ reported a negative correlation between plasma A β 42 and CSF A β 42 ($r=-0.35$). Future studies may need to compare AUC performances between PET and CSF markers, as well as their associations with IMR within the same cohort. Based on the current results, there is a lack of consistent findings to support diagnostic utility for this approach.

Amplified plasmonic exosome platform

While the above-mentioned techniques and platforms mainly addressed A β sensitivity issues, amplified plasmonic exosome (APEX) and the subsequently described methods have the potential to specifically measure different A β aggregation species. APEX is based on the finding that exosomes can bind to extracellular A β 42 proteins via glycoproteins or glycolipids on the exosomal plasma membrane. In their study, Lim *et al* established that exosomes have a higher binding affinity to larger A β 42 aggregates which also have a stronger propensity to form aggregated, fibrillar structures.⁴⁰ Briefly, after binding the target to the APEX nanosensor via the capture antibody, insoluble optical deposits are formed on the sensor through in-situ enzymatic amplification. This deposition enhances the surface plasmon resonance signal, as represented by a greater spectral shift. Through modifications in the sensor's design and fabrication (eg, patterned silicon nitride membrane, double-layered nanostructure, back illumination), APEX is designed to achieve nanoscale detection. To measure exosome-bound A β 42 from plasma, A β 42 is directly enriched from plasma via the A β 42 capture antibody on the APEX nanosensor, and the relative amount of CD63, an established exosomal marker associated with the captured A β 42, is measured via the CD63 detection antibody. Measurements are made relative to a sample-matched negative control, where the same sample was incubated over a control sensor functionalised with IgG isotype control antibody.

In their small cohort of 72 subjects, Lim *et al* reported a strong positive correlation between the level of plasma exosome-bound A β 42 and SUVR ($r=0.95$). Using the same cohort, Tanaka *et al*⁴¹ performed the first head-to-head comparison study between APEX and Simoa and showed that plasma exosome-bound A β 42 (AUC=0.99) outperformed Simoa-measured A β 42/A β 40 ratio (AUC=0.82) and A β 42 alone (AUC=0.78) in distinguishing between PET A β + and A β - subjects. The superior performance of the plasma exosome-bound A β 42 necessitates further validation in larger independent cohorts to establish APEX as a biomarker platform.

Multimer detection system

Multimer detection system (MDS) is a sandwich ELISA that preferentially detected oligomers over monomers.⁴² A unique epitope exists in the A β monomer, with multiple copies of this epitope found in the multimer. Thus, if antibodies targeting the unique/overlapping epitope were used for both capturing and detecting antibodies, the monomer would only be occupied by one of the antibodies and no signal would be produced. In contrast, multiple copies of the unique epitope in a multimer would allow binding of both the capturing and detecting antibodies to produce detectable signals.

In an earlier study, An *et al*⁴³ reported that MDS was unable to discriminate crude A β O levels in the plasma of patients with AD from controls due to low concentrations. However, spiking the plasma samples with synthetic A β 42 resulted in significant increases in A β O levels in the patients with AD, but not in controls.⁴³ A possible explanation is that oligomerisation of A β is influenced by potential factors in plasma of patients with AD, which may be absent or in lower concentrations in controls. Adopting a similar concept to measure plasma A β O by MDS, Wang *et al* spiked synthetic A β 42 peptide and used epitope-overlapping A β antibodies at the N-terminus 3-4 of A β .⁴² They reported a moderate correlation between plasma A β O levels and PIB PET SUVR ($r=0.43$) or CSF A β 42 levels ($r=-0.44$). The present study samples were small ($n=50$). Furthermore, the current long incubation time (144 hours) needs to be reduced to be feasible for clinical use.

Interdigitated microelectrode sensor system

Kim *et al* introduced the use of an interdigitated microelectrode (IME) sensor system, together with 4-(2-hydroxyethyl)-1-piperazinepropanesulfonic acid (EPPS)—a molecule which aids in converting aggregated A β into its monomeric form; for the measurement of plasma A β in the heterogenous and monomerised states.⁴⁴ Briefly, the chip's sensing zone was coated with anti-A β antibody to capture the A β in plasma. Two microchannels were incorporated on top of the chip to simultaneously load two different plasma samples—the EPPS treated and non-treated samples. The interaction between plasma A β and anti-A β antibodies was measured using an impedance measurement system, which was constructed to cancel noise signals from parasitic capacitance of IME and subsequently amplify the low-level signals. The impedance change increases as the concentration of A β increases. To mitigate interindividual and intraindividual variations in A β levels, a self-standard ratio was calculated by dividing the concentration of homogenous A β monomers (EPPS treated) by that of heterogenous A β (non-treated).

Using blood collected from two clinical institutes, Kim *et al* reported a positive correlation between the self-standard ratio and PET SUVR in both cohorts ($r=0.551$ and 0.414 , respectively). Similar to the majority of the other studies, more results, including AUC, are needed to establish the utility of this method.

Blood-derived P-tau: a promising marker of AD pathophysiology

Besides A β , there have been concerted efforts to accurately measure another established CSF biomarker, namely P-tau, in blood. These attempts are driven by the development of ultra-sensitive technologies enhanced by automation and improved throughput, such as the Simoa platform (see earlier). Following initial analytical challenges that hampered successful application of blood P-tau assays, recent advances in the understanding of tau biochemical processing in the brain as well as the subsequent

release of soluble P-tau into biofluids have enabled unprecedented rapid development of P-tau biomarkers in blood. First, a blood-based alternative of the established CSF marker of tau pathology, P-tau181, was developed by focusing on fragments containing the N-terminal portion of the protein that appears to be released into blood more abundantly than the mid-region forms that conventional CSF assays detect. Blood P-tau181 assessment methods from different sources, mainly the University of Gothenburg and Eli Lilly (using Simoa and MSD technologies, respectively), have shown excellent analytical and diagnostic performances in several landmark publications (tables 2 and 3, with detailed cohort information given in online supplemental table S2). Next, to be developed were P-tau biomarkers focusing on other phosphorylated sites, for example, P-tau217⁴⁵ and P-tau231⁴⁶ which have shown comparable diagnostic utility and, in some cases, superior predictive abilities particularly in the preclinical stages of AD (tables 2 and 3). Furthermore, IP-MS technology-based multiplexed assays have been developed that can measure multiple P-tau biomarkers in the same plasma samples simultaneously, including various tau species phosphorylated at residues apart from threonine-181.⁴⁷

Summary of the recent blood P-tau181 findings

Most current studies employed Simoa or MSD for the measurement of blood P-tau181, with results broadly similar across the two platforms. Several studies showed that blood P-tau181 concentrations were increased in MCI due to AD and AD dementia, but not in non-AD dementias including frontotemporal dementia (FTD) and vascular dementia (VaD).^{48–54} Specifically, blood P-tau181 was consistently elevated in the Aβ+ subjects (eg, Aβ+ MCI and Aβ+ AD).^{30 48 51 52 54–56} In a recent study, it was reported that the earliest increases in plasma P-tau181 occurred shortly before PET and CSF Aβ markers reached abnormal levels, with changes accelerated as the severity of Aβ pathology increased.⁵⁷ Moreover, plasma P-tau181 reached abnormal levels only after PET and CSF Aβ biomarkers reached relatively advanced abnormality levels—it was estimated that plasma P-tau181 reached abnormal levels approximately 6.5 and 5.7 years after CSF and PET Aβ measures.⁵⁷

Next, plasma P-tau181 was also shown to be able to differentiate AD from non-AD neurodegenerative diseases (AUCs=0.82–0.94).^{48 51–53} Plasma P-tau181 detected cross-sectional elevated brain Aβ (Aβ+) in combined CU, MCI and dementia subjects (all subjects; AUCs=0.72–0.95),^{45 47 48 51–54} as well as in subgroups of non-demented (CU+MCI; AUCs=0.71–0.81),^{30 51 52 54} CU (AUCs=0.67–0.86)^{30 46–48 53–55} and MCI subjects (AUCs=0.68–0.94).^{46–48 51 53–55} The range of AUCs observed may be due to differences in the definitions of Aβ positivity (which could be related to the different PET Aβ tracers or CSF cut-offs used in the different studies) and the cohorts in terms of prevalence of tau pathology, which is also highly associated with plasma P-tau181. Some of the above-mentioned studies further demonstrated P-tau181 associations, cross-sectionally and longitudinally, with NFT burden,^{48 52–55 57} grey matter atrophy,^{27 58} hippocampal atrophy,^{48 51 55} cortical atrophy brain,⁵⁹ metabolic dysfunction^{56 59} and cognitive impairment.^{27 48 51 55 60} As illustrated in the majority of these studies, besides performing the association analyses in the whole cohort, it may be important to stratify the subjects by their cognitive (or diagnostic) and Aβ status as the observed associations could be driven by a particular subgroup, such as the Aβ+ subjects. This may not be surprising given that blood P-tau181 likely reflects the pathophysiological processes specifically associated with AD pathology.

Table 2 Comparisons of blood phosphorylated tau (P-tau) among diagnostic groups stratified by Aβ status

Result	Cohort	Technology type	P-tau isoform	Ref
↑ Blood P-tau in Aβ+ AD or neuropathologically confirmed AD vs non-AD neurodegenerative diseases	TRIAD and BioFINDER-2	Simoa	P-tau181	48
		Simoa	P-tau181	49
	Neuropathology cohort	Simoa	P-tau181	50
		Simoa	P-tau181	51
	BioFINDER (cohort 1) and a neuropathology cohort	MSD	P-tau181	52
	UCSF Memory, Aging Center and ARTFL (cohort 1), and a neuropathology cohort	MSD	P-tau181	53
↑ Blood P-tau in Aβ+ AD vs Aβ– AD	BioFINDER-2 and a neuropathology cohort	MSD	P-tau217	45
	TRIAD and a neuropathology cohort	Simoa	P-tau231	46
↑ Blood P-tau in Aβ+ AD vs Aβ+ MCI	TRIAD and BioFINDER-2	Simoa	P-tau181	48
	BioFINDER-2	MSD	P-tau217	45
	TRIAD	Simoa	P-tau231	46
↑ Blood P-tau in Aβ+ AD vs Aβ– MCI	TRIAD and BioFINDER-2	Simoa	P-tau181	48
		Simoa	P-tau181	55
	BioFINDER-2	MSD	P-tau217	45
	TRIAD	Simoa	P-tau231	46
↑ Blood P-tau in Aβ+ AD vs Aβ+ CU and Aβ– CU	TRIAD and BioFINDER-2	Simoa	P-tau181	48
		Simoa	P-tau181	55
		MSD	P-tau181	54
	BioFINDER (cohort 1)	MSD	P-tau181	52
	BioFINDER-2	MSD	P-tau217	45
↑ Blood P-tau in Aβ+ MCI vs Aβ– MCI	TRIAD	Simoa	P-tau231	46
	BioFINDER-2	Simoa	P-tau181	48
		Simoa	P-tau181	51
		Simoa	P-tau181	55
	BioFINDER (cohort 2)	MSD	P-tau181	52
	BioFINDER-2	MSD	P-tau217	45
↑ Blood P-tau in Aβ+ MCI vs Aβ+ CU	TRIAD	Simoa	P-tau231	46
		Simoa	P-tau181	55
	BioFINDER (cohort 1 and 2)	MSD	P-tau181	52
		Simoa	P-tau181	55
		MSD	P-tau181	52
↑ Blood P-tau in Aβ+ MCI vs Aβ– CU	TRIAD and BioFINDER-2	Simoa	P-tau181	48
		Simoa	P-tau181	51
		Simoa	P-tau181	55
	BioFINDER (cohort 1 and 2)	MSD	P-tau181	52
		MSD	P-tau181	54
	Validation cohort	IP-MS	P-tau181	47
	BioFINDER-2	MSD	P-tau217	45
	Validation cohort	IP-MS	P-tau217	47
	TRIAD	Simoa	P-tau231	46
		Simoa	P-tau181	48
↑ Blood P-tau in Aβ+ CU vs Aβ– CU	TRIAD and BioFINDER-2	Simoa	P-tau181	48
		Simoa	P-tau181	55
	BioFINDER (cohort 1 and 2)	MSD	P-tau181	52
	BioFINDER-2	MSD	P-tau217	45
	Validation cohort	IP-MS	P-tau217	47
	TRIAD	Simoa	P-tau231	46

Continued

Table 2 Continued

Result	Cohort	Technology type	P-tau isoform	Ref
↑ Blood P-tau in Aβ+ CU vs Aβ- MCI	BioFINDER-2	Simoa	P-tau181	48
	BioFINDER (cohort 2)	MSD	P-tau181	52
	BioFINDER-2	MSD	P-tau217	45
	TRIAD	Simoa	P-tau231	46

Aβ, amyloid-β; AD, Alzheimer's disease; CU, Cognitively unimpaired; IP-MS, immunoprecipitation-mass spectrometry; MCI, mild cognitive impairment; MSD, Meso Scale Discovery; Simoa, single molecule array; UCSF, University of California at San Francisco.

Table 3 AUCs of blood phosphorylated tau (P-tau) in distinguishing between Aβ+ and Aβ- participants

	AUC	Remarks	Technology	Ref
P-tau181	0.88	CU young adults+CU elderly +MCI+AD+FTD	Simoa	48
	0.84	CU+MCI (CIND)+AD+VaD	Simoa	51
	0.76	CU+MCI+AD+non-AD	Simoa	45
	0.80	CU+MCI+AD	MSD	54
	0.79	CU+MCI+AD+non-AD	MSD	52
	0.91	CU+MCI +AD+FTLD	MSD	53
	0.82	CU+AD	MSD	87
	0.95	CU young +CU elderly +MCI+AD (discovery cohort)	IP-MS	47
	0.72	CU+MCI+AD (validation cohort)		
	0.77	CU	Simoa	48
	0.70	CU	Simoa	55
	0.77	CU	Simoa	46
	0.72	CU	Simoa	30
	0.70	CU	MSD	54
	0.86	CU	MSD	53
	0.67	CU (validation cohort)	IP-MS	47
	0.74	MCI	Simoa	48
	0.80	MCI	Simoa	55
	0.82	MCI (CIND)	Simoa	51
	0.75	MCI	Simoa	46
	0.85	MCI	MSD	54
	0.94	MCI	MSD	53
	0.68	MCI (validation cohort)	IP-MS	47
	0.80	CU+MCI (CIND)	Simoa	51
	0.71	CU+MCI+prior neurological conditions	Simoa	30
	0.75	CU+MCI	MSD	54
	0.81	CU+MCI	MSD	52
P-tau217	0.87	CU+MCI+AD+non-AD	MSD	45
	0.84	CU+AD	MSD	87
	0.99	CU young +CU elderly +MCI+ AD (discovery cohort)	IP-MS	47
	0.93	CU+MCI+AD (validation cohort)		
	0.86	CU (validation cohort)		
	0.93	MCI (validation cohort)		
P-tau231	0.83	CU	Simoa	46
	0.80	MCI	Simoa	

Aβ, amyloid-β; AD, Alzheimer's disease; AUC, area under the receiver operating characteristic curve; CIND, Cognitively impaired no dementia; CU, Cognitively unimpaired; FTD, Frontotemporal dementia; FTLD, Frontotemporal lobar degeneration; IP-MS, immunoprecipitation-mass spectrometry; MCI, mild cognitive impairment; MSD, Meso Scale Discovery; Simoa, single molecule array; VaD, Vascular dementia.

Given associations between blood P-tau181 and AD pathology, the P-tau biomarker may be useful in research studies exploring disease pathogenesis, such as genetic risk factors for AD. For instance, a study showed that polygenic risk scores (PRSs) for AD including APOE (APOE PRSs) was associated with plasma P-tau181 independent of diagnostic and Aβ pathology status. In contrast, polygenic risk for AD beyond APOE (non-APOE PRSs) was associated with plasma P-tau181 only in MCI and PET Aβ+ subjects, suggesting that having MCI and being Aβ+ is the most vulnerable combination (in comparison with other status such as CU or Aβ- status).⁶¹ These results give insight into the relationship between genetic risk for AD and P-tau181, which warrants further studies.

Comparison of biomarker performance between P-tau species

A few papers have evaluated P-tau181 and P-tau217 in the very early phases of MCI/dementia. A study examining familial AD found significantly higher P-tau217 concentrations 20 years before the expected year of onset (EYO) of symptoms in individuals diagnosed with MCI,⁴⁵ while another reported significant increases in P-tau181, 16 years before EYO in symptomatic Familial AD mutation carriers (both MCI and AD).⁶² More recently, head-to-head comparisons of N-terminal-directed P-tau181 and P-tau217 biomarkers in CSF^{63,64} showed similar performances; however, both were superior to conventional P-tau181 measured on mid-region fragments. These findings support the idea that both the measured plasma P-tau biomarkers are increased in the preclinical phase of FAD. Further, studies evaluating these P-tau biomarkers in later stages of disease showed tight associations between biomarker levels in plasma samples taken during life and AD neuropathology assessed post-mortem, with AUC values for differentiating AD from non-AD neurodegenerative disorders at 0.89 for P-tau217,⁴⁵ 0.97 for P-tau181⁵⁰ and 0.997 for P-tau231⁴⁶ (online supplemental table S2). These findings further support the hypothesis that increased P-tau specifically reflect AD pathology. However, further studies directly comparing these tau biomarkers in plasma using the same assay technology in the same cohort are needed to ascertain whether P-tau181, P-tau217 and P-tau231 substantially differ in diagnostic utility in different phases of the AD continuum. In asymptomatic AD, P-tau231 shows increases in individuals with subthreshold levels of PET-measured Aβ burden in the first quartile, while CSF P-tau217 and plasma P-tau181 were increased in the third or fourth quartile.⁴⁶ Finally, a few studies have demonstrated that longitudinal increases of blood P-tau levels were associated with longitudinal brain atrophy and cognitive decline, particularly in the patients with AD, and may differentiate MCI converters from non-converters,^{27,46,48,65} suggesting potential utility in longer-term monitoring in therapeutic trials.

Multi-marker approach towards AD biomarkers

Since AD is pathologically characterised by amyloid plaques, NFTs as well as neurodegeneration, a comprehensive ATN (amyloid, tau, neurodegeneration) classification system has been advanced to better account for the complex pathophysiological processes central to AD pathogenesis.⁸ To the extent that blood-based biomarkers accurately reflects brain changes, a corresponding combination of biomarkers may yield superior utility as well. For example, using data from an Asian cohort of patients with AD with concomitant cerebrovascular diseases (CeVDs), it has been reported that combining Simoa P-tau181 with Aβ42 measures yielded better AUCs for amyloid positivity and hippocampal atrophy than any single

marker evaluated (P-tau181, total tau, A β 40 and A β 42).⁵¹ An earlier study reported similar improved performance of combining IMR measurements of A β and tau in identifying AD in both prodromal and dementia phases.⁶⁶ However, as various P-tau species seem to be associated with both amyloid and tau pathologies,⁶⁷ they could potentially be used together as biomarkers to detect A and T pathophysiology, especially in settings where IP-MS plasma A β measurements are not feasible, or when plasma A β measures cannot distinguish A+ and A- cases. Nonetheless, the fact that different P-tau markers appear to be altered at different stages of AD pathology could be a source of complications.

Finally, amyloid and tau pathologies are also associated with various pathophysiological conditions, including synaptic dysfunction, neuroaxonal damage and injury, neuroinflammation and oxidative stress⁶⁸; as well as with concomitant CeVDs which may be relatively frequent in Asian and other less-studied populations and interact additively or synergistically with AD in worsening cognitive functions.^{69–71} It would therefore be of interest to investigate if combining A β and tau markers with those for synaptic, neuroaxonal injury, inflammatory, oxidative stress and endothelial/vascular injury in a multi-marker panel may yield improved clinical utility and further insights into disease mechanisms. A potential blood biomarker candidate is neurofilament-L (NfL), which shows good correlation with CSF NfL and reflects severity of axonal degeneration and injury.^{72–73} NfL is a disease-unspecific marker, as elevations are observed in several neurodegenerative disorders including AD,⁷⁴ thus limiting its differential diagnostic potential. Nonetheless, studies have reported on the utility of blood NfL for predicting and monitoring longitudinal changes in cognition,^{65–75–77} brain atrophy^{65–75–79} and hypometabolism.^{65–73–75–77} Unlike P-tau181, which was associated with cognitive decline and prospective neurodegeneration that are characteristic for AD, the associations of blood NfL were not AD specific.⁶⁵ Furthermore, both blood biomarkers were independently associated with longitudinal changes in cognition and AD-typical neurodegeneration.⁶⁵ Taken together, the combination of blood NfL to the multi-marker panel may aid as a screening tool to identify individuals with ongoing neurodegeneration and improve prediction of disease progression in AD, as well as monitor treatment response in disease-modifying trials (eg, downstream effects of the tested drugs on neurodegeneration).^{8–72} Besides NfL, another potential blood biomarker is glial fibrillar acidic protein (GFAP), which showed early increases with rising brain A β deposition,⁸⁰ was positively associated with A β PET in cognitively normal individuals,^{80–81} and demonstrated good diagnostic performance in identifying elevated brain amyloid among non-demented elderly (AUC=0.75–0.80).^{80–82–84} Notably, the diagnostic performance improved with the inclusion of plasma A β 42/A β 40 ratio and other AD risk factors such as age, gender and/or APOE ϵ 4 (AUC increased from 0.84 to 0.92).^{82–84} Altogether, these results indicate that blood GFAP may be an early biomarker of reactive astrogliosis associated with A β pathology in the prodementia phase. Moreover, higher baseline blood GFAP levels in non-demented elderly were associated with progression to dementia and steeper rates of cognitive decline,^{84–85} with effect sizes greater than NfL,⁸⁵ implying that GFAP could be a better prognostic marker for incident dementia than NfL. More head-to-head comparisons of these promising blood biomarkers of AD pathology (P-tau and A β), axonal injury (NfL) and reactive astrogliosis (GFAP) should be performed to evaluate their diagnostic and prognostic capabilities, alone and in combination.

ETHICAL AND CONSUMER ISSUES ASSOCIATED WITH BLOOD BIOMARKERS

In the near future, the increasing ability to receive accurate diagnostic and prognostic information with the ease of a blood test will revolutionise AD research and care. Given the concerns surrounding the risk of developing AD and related dementias, such testing may also be the basis of profitable direct-to-consumer (DTC) tests which may come with significant ethical and social issues, as highlighted in.⁸⁶ Briefly, DTC blood testing may raise concerns about confidentiality and discrimination, as when a DTC testing company sells consumers' data to third parties without informed consent. In regions where laws offer limited protections against biomarker-based discrimination, insurers could use the positive blood test result to refuse an individual of disability or long-term care insurance for prodromal individuals, and companies may deny opportunities for employment or promotion. Next, in societies that lack proper awareness of AD, DTC blood testing may amplify AD-associated stigma and discrimination even in the absence of cognitive symptoms. Therefore, it is necessary to communicate the facts and educate the public to dispel misconceptions about the disease. Furthermore, policies to protect individuals with positive DTC test results from discrimination and stigmatisation should be implemented. Before these issues are addressed, it is advisable that blood testing for dementia is restricted by regulation to the clinical setting.

Concluding remarks

Recent advances in the development of ultrasensitive, high throughput analytical technologies and platforms have enabled the discovery and potential clinical application of promising biomarkers that reflect AD brain pathology in blood samples. These biomarkers will be critical in both diagnostic and prognostic assessments for AD as well as in longitudinal monitoring and therapeutic trials. However, there is a need for further research to (1) directly compare the performance of different biomarker platforms; (2) validate initial findings in larger cohorts from diverse backgrounds and (3) evaluate the utility of multi-marker panels, in order to fully realise the potential of blood-based biomarkers for AD and associated conditions.

Author affiliations

¹Memory, Aging and Cognition Centre, National University Health Systems, Singapore

²Department of Pharmacology, Yong Loo Lin School of Medicine, National University of Singapore, Singapore

³Wallenberg Centre for Molecular and Translational Medicine, University of Gothenburg, Gothenburg, Sweden

⁴Psychology and Neuroscience, King's College London, Institute of Psychiatry, Maurice Wohl Institute Clinical Neuroscience Institute, London, UK

⁵NIHR Biomedical Research Centre for Mental Health and Biomedical Research Unit for Dementia, South London and Maudsley NHS Foundation, London, UK

⁶Department of Psychiatry and Neurochemistry, Institute of Neuroscience and Physiology, The Sahlgrenska Academy at the University of Gothenburg, Mölndal, Sweden

⁷Department of Psychiatry, University of Pittsburgh, Pittsburgh, Pennsylvania, USA

⁸Department of Neurology, National Cerebral and Cardiovascular Center, Suita, Osaka, Japan

⁹Clinical Imaging Research Centre, Yong Loo Lin School of Medicine, National University of Singapore, Singapore

¹⁰Department of Neurodegenerative Disease, UCL Queen Square Institute of Neurology, University College London, London, UK

¹¹Clinical Neurochemistry Laboratory, Sahlgrenska University Hospital, Gothenburg, Sweden

¹²UK Dementia Research Institute at UCL, University College London, London, UK

Contributors JCR, NA, TKK, TT, MS, HZ, KB, CL-HC and ML contributed to the conception and structure of the review, drafted the text and prepared the tables.

Funding This work is supported by the National Medical Research Council of Singapore (NMRC/CSA-SI/007/2016). TKK was funded by the BrightFocus Foundation (#A2020812F), the International Society for Neurochemistry's Career Development Grant, the Swedish Alzheimer Foundation (Alzheimerfonden; #AF-930627), the Swedish Brain Foundation (Hjärnfonden; #FO2020-0240), the Swedish Dementia Foundation (Demensförbundet), the Swedish Parkinson Foundation (Parkinsonfonden), Gamla Tjänarinnor Foundation, the Aina (Ann) Wallströms and Mary-Ann Sjöbloms Foundation, the Agneta Prytz-Folkes & Gösta Folkes Foundation (#2020-00124), the Gun and Bertil Stohnes Foundation and the Anna Lisa and Brother Björnsson's Foundation. MS is supported by the Knut and Alice Wallenberg Foundation (Wallenberg Centre for Molecular and Translational Medicine Fellow; KAW 2014.0363), the Swedish Research Council (#2017-02869), and the Swedish state under the agreement between the Swedish government and the County Councils, the ALF-agreement (#ALFGBG-813971) (all paid to the institution). HZ is a Wallenberg Scholar supported by grants from the Swedish Research Council (#2018-02532), the European Research Council (#681712), Swedish State Support for Clinical Research (#ALFGBG-720931), the Alzheimer Drug Discovery Foundation (ADDF), USA (#201809-2016862), the AD Strategic Fund and the Alzheimer's Association (#ADSF-21-831376-C, #ADSF-21-831381-C and #ADSF-21-831377-C), the Olav Thon Foundation, the Erling-Persson Family Foundation, Stiftelsen för Gamla Tjänarinnor, Hjärnfonden, Sweden (#FO2019-0228), the European Union's Horizon 2020 research and innovation programme under the Marie Skłodowska-Curie grant agreement No 860197 (MIRIADE), and the UK Dementia Research Institute at UCL.

Competing interests HZ has served at scientific advisory boards for Alector, Eisai, Denali, Roche Diagnostics, Wave, Samumed, Siemens Healthineers, Pinteon Therapeutics, Nervgen, AZTherapies and CogRx, has given lectures in symposia sponsored by Celectricon, Fujirebio, Abecore and Biogen, and is a co-founder of Brain Biomarker Solutions in Gothenburg AB (BBS), which is a part of the GU Ventures Incubator Programme. KB has served as a consultant, at advisory boards, or at data monitoring committees for Abcam, Axon, Biogen, JOMDD/Shimadzu, Julius Clinical, Lilly, MagQu, Novartis, Prothena, Roche Diagnostics and Siemens Healthineers, and is a co-founder of Brain Biomarker Solutions in Gothenburg AB (BBS), which is a part of the GU Ventures Incubator Programme.

Patient consent for publication Not required.

Provenance and peer review Commissioned; externally peer reviewed.

Supplemental material This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

ORCID iDs

Henrik Zetterberg <http://orcid.org/0000-0003-3930-4354>
Mitchell K.P. Lai <http://orcid.org/0000-0001-7685-1424>

REFERENCES

- World Alzheimer Report. *The state of the art of dementia research: new frontiers*. Alzheimer's Disease International, 2018.
- Alzheimer's Association. 2018 Alzheimer's disease facts and figures. *Alzheimers Dement* 2018;14:367–429.
- Ashton NJ, Schöll M, Heurling K, et al. Update on biomarkers for amyloid pathology in Alzheimer's disease. *Biomark Med* 2018;12:799–812.
- Jack CR, Knopman DS, Jagust WJ, et al. Hypothetical model of dynamic biomarkers of the Alzheimer's pathological cascade. *Lancet Neurol* 2010;9:119–28.
- Shaw LM, Arias J, Blennow K, et al. Appropriate use criteria for lumbar puncture and cerebrospinal fluid testing in the diagnosis of Alzheimer's disease. *Alzheimers Dement* 2018;14:1505–21.
- Krishnadas N, Villemagne VL, Doré V, et al. Advances in brain amyloid imaging. *Semin Nucl Med* 2021;51:241–52.
- Hampel H, O'Bryant SE, Molinuevo JL, et al. Blood-based biomarkers for Alzheimer disease: mapping the road to the clinic. *Nat Rev Neurol* 2018;14:639–52.
- Jack CR, Bennett DA, Blennow K, et al. NIA-AA research framework: toward a biological definition of Alzheimer's disease. *Alzheimers Dement* 2018;14:535–62.
- Bateman RJ, Blennow K, Doody R, et al. Plasma biomarkers of AD emerging as essential tools for drug development: an EU/US CTAD Task force report. *J Prev Alzheimers Dis* 2019;6:169–73.
- Tolar M, Abushakra S, Hey JA, et al. Aducanumab, gantenerumab, BAN2401, and ALZ-801—the first wave of amyloid-targeting drugs for Alzheimer's disease with potential for near term approval. *Alzheimers Res Ther* 2020;12:95.
- Mintun MA, Lo AC, Duggan Evans C, et al. Donanemab in early Alzheimer's disease. *N Engl J Med* 2021;384:1691–704.
- Swanson CJ, Zhang Y, Dhadda S, et al. A randomized, double-blind, phase 2B proof-of-concept clinical trial in early Alzheimer's disease with lecanemab, an anti-A β protofibril antibody. *Alzheimers Res Ther* 2021;13:80.
- Swanson CJ, Zhang Y, Dhadda S, Wang J. Clinical and biomarker updates from BAN2401 Study 201 in early AD. Clinical Trials on Alzheimer's Disease Conference: 24–27 October 2018; Barcelona.
- Swanson C, Zhang Y, Dhadda S, Wang J. The GAP study of BAN2401 study 201 in early AD. Persistence of BAN2401-mediated amyloid reductions post-treatment: a preliminary comparison of amyloid status between the core phase of BAN2401-g000-201 and baseline of the open-label extension phase in subjects with early Alzheimer's disease. Clinical Trials on Alzheimer's Disease conference: 4–7 December 2019; San Diego.
- Budd-Haeberlein S, VHC, Tian Y, Chalkias S. EMERGE and ENGAGE topline results: two phase 3 studies to evaluate aducanumab in patients with early Alzheimer's disease. Clinical Trials on Alzheimer's Disease Conference: 4–7 December 2019; San Diego.
- Toledo JB, Shaw LM, Trojanowski JQ. Plasma amyloid beta measurements - a desired but elusive Alzheimer's disease biomarker. *Alzheimers Res Ther* 2013;5:8.
- Hansson O, Zetterberg H, Vanmechelen E, et al. Evaluation of plasma A β 40 and A β 42 as predictors of conversion to Alzheimer's disease in patients with mild cognitive impairment. *Neurobiol Aging* 2010;31:357–67.
- Janelidze S, Stomrud E, Palmqvist S, et al. Plasma β -amyloid in Alzheimer's disease and vascular disease. *Sci Rep* 2016;6:26801.
- Larson ME, Lesné SE. Soluble A β oligomer production and toxicity. *J Neurochem* 2012;120:125–39.
- Sakono M, Zako T. Amyloid oligomers: formation and toxicity of Abeta oligomers. *Febs J* 2010;277:1348–58.
- Lesné SE, Sherman MA, Grant M, et al. Brain amyloid- β oligomers in ageing and Alzheimer's disease. *Brain* 2013;136:1383–98.
- Schindler SE, Bollinger JG, Ovod V, et al. High-Precision plasma β -amyloid 42/40 predicts current and future brain amyloidosis. *Neurology* 2019;93:e1647–59.
- De Meyer S, Schaeverbeke JM, Verberk IMW, et al. Comparison of ELISA- and SIMOA-based quantification of plasma A β ratios for early detection of cerebral amyloidosis. *Alzheimers Res Ther* 2020;12:162.
- Zetterberg H, Mörtberg E, Song L, et al. Hypoxia due to cardiac arrest induces a time-dependent increase in serum amyloid β levels in humans. *PLoS One* 2011;6:e28263.
- Verberk IMW, Slot RE, Verfaillie SCJ, et al. Plasma amyloid as Prescanner for the earliest Alzheimer pathological changes. *Ann Neurol* 2018;84:648–58.
- Thijssen E, Verberk IMW, Vanderstichele HM. O3-09-06: a prototype SIMOA assay quantifying plasma amyloid beta 1-42 and 1-40 isoforms can differentiate participants with AD from healthy control subjects. *Alzheimer's & Dementia* 2018;14:P1039.
- Simrén J, Leuzy A, Karikari TK, et al. The diagnostic and prognostic capabilities of plasma biomarkers in Alzheimer's disease. *Alzheimers Dement* 2021;17:1145–56.
- Kaneko N, Nakamura A, Washimi Y, et al. Novel plasma biomarker surrogating cerebral amyloid deposition. *Proc Jpn Acad Ser B Phys Biol Sci* 2014;90:353–64.
- Nakamura A, Kaneko N, Villemagne VL, et al. High performance plasma amyloid- β biomarkers for Alzheimer's disease. *Nature* 2018;554:249–54.
- Keshavan A, Pannee J, Karikari TK, et al. Population-based blood screening for preclinical Alzheimer's disease in a British birth cohort at age 70. *Brain* 2021;144:434–49.
- Ovod V, Ramsey KN, Mawuenyega KG, et al. Amyloid β concentrations and stable isotope labeling kinetics of human plasma specific to central nervous system amyloidosis. *Alzheimers Dement* 2017;13:841–9.
- Pannee J, Törnqvist U, Westerlund A, et al. The amyloid- β degradation pattern in plasma—a possible tool for clinical trials in Alzheimer's disease. *Neurosci Lett* 2014;573:7–12.
- Janelidze S, Zetterberg H, Mattsson N, et al. CSF A β 42/A β 40 and A β 42/A β 38 ratios: better diagnostic markers of Alzheimer disease. *Ann Clin Transl Neurol* 2016;3:154–65.
- Molinie JL, Ayton S, Batrla R, et al. Current state of Alzheimer's fluid biomarkers. *Acta Neuropathol* 2018;136:821–53.
- Palmqvist S, Janelidze S, Stomrud E, et al. Performance of fully automated plasma assays as screening tests for Alzheimer disease-related β -amyloid status. *JAMA Neurol* 2019;76:e191632.
- Henriksen K, O'Bryant SE, Hampel H, et al. The future of blood-based biomarkers for Alzheimer's disease. *Alzheimers Dement* 2014;10:115–31.
- Vogelsgang J, Shahpasand-Kroner H, Vogelsgang R, et al. Multiplex immunoassay measurement of amyloid- β_{42} to amyloid- β_{40} ratio in plasma discriminates between dementia due to Alzheimer's disease and dementia not due to Alzheimer's disease. *Exp Brain Res* 2018;236:1241–50.
- Teunissen CE, Chiu M-J, Yang C-C, et al. Plasma amyloid- β (A β 42) correlates with cerebrospinal fluid A β 42 in Alzheimer's disease. *J Alzheimers Dis* 2018;62:1857–63.
- Tzen K-Y, Yang S-Y, Chen T-F, et al. Plasma A β but not tau is related to brain PIB retention in early Alzheimer's disease. *ACS Chem Neurosci* 2014;5:830–6.
- Lim CZJ, Zhang Y, Chen Y, et al. Subtyping of circulating exosome-bound amyloid β reflects brain plaque deposition. *Nat Commun* 2019;10:1144.

- 41 Tanaka Tet *et al.* Head-To-Head comparison of amplified plasmonic exosome A β 42 platform and single-molecule array immunoassay in a memory clinic cohort. *Eur J Neurol* 2020.
- 42 Wang MJ, Yi S, Han J-Y, *et al.* Oligomeric forms of amyloid- β protein in plasma as a potential blood-based biomarker for Alzheimer's disease. *Alzheimers Res Ther* 2017;9:98.
- 43 An SSA *et al.* Dynamic changes of oligomeric amyloid β levels in plasma induced by spiked synthetic A β 42. *Alzheimer's Res Ther* 2017;9:86.
- 44 Kim Y, Yoo YK, Kim HY, *et al.* Comparative analyses of plasma amyloid- β levels in heterogeneous and monomerized states by interdigitated microelectrode sensor system. *Sci Adv* 2019;5:eaav1388.
- 45 Palmqvist S, Janelidze S, Quiroz YT, *et al.* Discriminative accuracy of plasma Phospho-tau217 for Alzheimer disease vs other neurodegenerative disorders. *JAMA* 2020;324:772–81.
- 46 Ashton NJ, Pascoal TA, Karikari TK, *et al.* Plasma p-tau231: a new biomarker for incipient Alzheimer's disease pathology. *Acta Neuropathol* 2021;141:709–24.
- 47 Barthélemy NR, Horie K, Sato C, *et al.* Blood plasma phosphorylated-tau isoforms track CNS change in Alzheimer's disease. *J Exp Med* 2020;217.
- 48 Karikari TK, Pascoal TA, Ashton NJ, *et al.* Blood phosphorylated tau 181 as a biomarker for Alzheimer's disease: a diagnostic performance and prediction modelling study using data from four prospective cohorts. *Lancet Neurol* 2020;19:422–33.
- 49 Benussi A, Karikari TK, Ashton N, *et al.* Diagnostic and prognostic value of serum NfL and p-Tau₁₈₁ in frontotemporal lobar degeneration. *J Neurol Neurosurg Psychiatry* 2020;91:960–7.
- 50 Lantero Rodriguez J, Karikari TK, Suárez-Calvet M, *et al.* Plasma p-tau181 accurately predicts Alzheimer's disease pathology at least 8 years prior to post-mortem and improves the clinical characterisation of cognitive decline. *Acta Neuropathol* 2020;140:267–78.
- 51 Chong JR, Ashton NJ, Karikari TK. Plasma p-Tau181 to A β 42 ratio is associated with brain amyloid burden and hippocampal atrophy in an Asian cohort of Alzheimer's disease patients with concomitant cerebrovascular disease. *Alzheimers Dement* 2021 doi:10.1002/alz.12332
- 52 Janelidze S, Mattsson N, Palmqvist S, *et al.* Plasma p-Tau181 in Alzheimer's disease: relationship to other biomarkers, differential diagnosis, neuropathology and longitudinal progression to Alzheimer's dementia. *Nat Med* 2020;26:379–86.
- 53 Thijssen EH, La Joie R, Wolf A, *et al.* Diagnostic value of plasma phosphorylated tau 181 in Alzheimer's disease and frontotemporal lobar degeneration. *Nat Med* 2020;26:387–97.
- 54 Mielke MM, Hagen CE, Xu J, *et al.* Plasma phospho-tau181 increases with Alzheimer's disease clinical severity and is associated with tau- and amyloid-positron emission tomography. *Alzheimer's & Dementia* 2018;14:989–97.
- 55 Karikari TK, Benedet AL, Ashton NJ. Diagnostic performance and prediction of clinical progression of plasma phospho-tau181 in the Alzheimer's Disease Neuroimaging Initiative. *Mol Psychiatry* 2020.
- 56 Lussier FZ, Benedet AL, Theriault J, *et al.* Plasma levels of phosphorylated tau 181 are associated with cerebral metabolic dysfunction in cognitively impaired and amyloid-positive individuals. *Brain Commun* 2021;3:fcab073.
- 57 Moscoso A, Grothe MJ, Ashton NJ, *et al.* Time course of phosphorylated-tau181 in blood across the Alzheimer's disease spectrum. *Brain* 2021;144:325–39.
- 58 Tissot C, L Benedet A, Theriault J, *et al.* Plasma pTau181 predicts cortical brain atrophy in aging and Alzheimer's disease. *Alzheimers Res Ther* 2021;13:69.
- 59 Hansson O, Cullen N, Zetterberg H, *et al.* Plasma phosphorylated tau181 and neurodegeneration in Alzheimer's disease. *Ann Clin Transl Neurol* 2021;8:259–65.
- 60 Clark C, Lewczuk P, Kornhuber J, *et al.* Plasma neurofilament light and phosphorylated tau 181 as biomarkers of Alzheimer's disease pathology and clinical disease progression. *Alzheimers Res Ther* 2021;13:65.
- 61 Zettergren A, Lord J, Ashton NJ, *et al.* Association between polygenic risk score of Alzheimer's disease and plasma phosphorylated tau in individuals from the Alzheimer's disease neuroimaging initiative. *Alzheimers Res Ther* 2021;13:17.
- 62 O'Connor A, Karikari TK, Poole T. Plasma phospho-tau181 in presymptomatic and symptomatic familial Alzheimer's disease: a longitudinal cohort study. *Mol Psychiatry* 2020;34.
- 63 Karikari TK, Emeršič A, Vrillon A, *et al.* Head-to-head comparison of clinical performance of CSF phospho-tau T181 and T217 biomarkers for Alzheimer's disease diagnosis. *Alzheimers Dement* 2021;17:755–67.
- 64 Suárez-Calvet M, Karikari TK, Ashton NJ, *et al.* Novel tau biomarkers phosphorylated at T181, T217 or T231 rise in the initial stages of the preclinical Alzheimer's continuum when only subtle changes in A β pathology are detected. *EMBO Mol Med* 2020;12:e12921.
- 65 Moscoso A, Grothe MJ, Ashton NJ, *et al.* Longitudinal associations of blood phosphorylated Tau181 and neurofilament light chain with neurodegeneration in Alzheimer disease. *JAMA Neurol* 2021;78:396–406.
- 66 Chiu M-J, Yang S-Y, Horng H-E, *et al.* Combined plasma biomarkers for diagnosing mild cognition impairment and Alzheimer's disease. *ACS Chem Neurosci* 2013;4:1530–6.
- 67 Mattsson-Carlgren N, Janelidze S, Bateman RJ, *et al.* Soluble P-tau217 reflects amyloid and tau pathology and mediates the association of amyloid with tau. *EMBO Mol Med* 2021;13:e14022.
- 68 Pereira C, Agostinho P, Moreira PI, *et al.* Alzheimer's disease-associated neurotoxic mechanisms and neuroprotective strategies. *Curr Drug Targets CNS Neurol Disord* 2005;4:383–403.
- 69 Chen C, Homma A, Mok VCT, *et al.* Alzheimer's disease with cerebrovascular disease: current status in the Asia-Pacific region. *J Intern Med* 2016;280:359–74.
- 70 Attems J, Jellinger KA. The overlap between vascular disease and Alzheimer's disease—lessons from pathology. *BMC Med* 2014;12:206.
- 71 Lam BYK, Yiu B, Ampil E, *et al.* High burden of cerebral white matter lesion in 9 Asian cities. *Sci Rep* 2021;11:11587.
- 72 Gaetani L, Blennow K, Calabresi P, *et al.* Neurofilament light chain as a biomarker in neurological disorders. *J Neurol Neurosurg Psychiatry* 2019;90:870–81.
- 73 Benedet AL, Ashton NJ, Pascoal TA, *et al.* Plasma neurofilament light associates with Alzheimer's disease metabolic decline in amyloid-positive individuals. *Alzheimers Dement* 2019;11:679–89.
- 74 Ashton NJ, Janelidze S, Al Khleifat A, *et al.* A multicentre validation study of the diagnostic value of plasma neurofilament light. *Nat Commun* 2021;12:3400.
- 75 Mielke MM, Syrjanen JA, Blennow K, *et al.* Plasma and CSF neurofilament light: relation to longitudinal neuroimaging and cognitive measures. *Neurology* 2019;93:e252–60.
- 76 Mattsson N, Andreasson U, Zetterberg H, *et al.* Association of plasma neurofilament light with neurodegeneration in patients with Alzheimer disease. *JAMA Neurol* 2017;74:557–66.
- 77 Mattsson N, Cullen NC, Andreasson U, *et al.* Association between longitudinal plasma neurofilament light and neurodegeneration in patients with Alzheimer disease. *JAMA Neurol* 2019;76:791–9.
- 78 Weston PSJ, Poole T, Ryan NS, *et al.* Serum neurofilament light in familial Alzheimer disease: a marker of early neurodegeneration. *Neurology* 2017;89:2167–75.
- 79 Benedet AL, Leuzys A, Pascoal TA, *et al.* Stage-specific links between plasma neurofilament light and imaging biomarkers of Alzheimer's disease. *Brain* 2020;143:3793–804.
- 80 Pereira JB, Janelidze S, Smith R, *et al.* Plasma GFAP is an early marker of amyloid- β but not tau pathology in Alzheimer's disease. *Brain* 2021.
- 81 Asken BM, Elahi FM, La Joie R, *et al.* Plasma glial fibrillary acidic protein levels differ along the spectra of amyloid burden and clinical disease stage. *J Alzheimers Dis* 2020;78:265–76.
- 82 Verberk IMW, Thijssen E, Koelewijn J, *et al.* Combination of plasma amyloid beta_(1–42/1–40) and glial fibrillary acidic protein strongly associates with cerebral amyloid pathology. *Alzheimers Res Ther* 2020;12:118.
- 83 Chatterjee P, Pedrini S, Stoops E, *et al.* Plasma glial fibrillary acidic protein is elevated in cognitively normal older adults at risk of Alzheimer's disease. *Transl Psychiatry* 2021;11:27.
- 84 Cicognola C, Janelidze S, Hertz J, *et al.* Plasma glial fibrillary acidic protein detects Alzheimer pathology and predicts future conversion to Alzheimer dementia in patients with mild cognitive impairment. *Alzheimers Res Ther* 2021;13:68.
- 85 Verberk IMW, Laarhuis MB, van den Bosch KA, *et al.* Serum markers glial fibrillary acidic protein and neurofilament light for prognosis and monitoring in cognitively normal older people: a prospective memory clinic-based cohort study. *Lancet Healthy Longev* 2021;2:e87–95.
- 86 Largent EA, Wexler A, Karlawish J. The future is P-Tau-Anticipating direct-to-consumer Alzheimer disease blood tests. *JAMA Neurol* 2021;78:379–80.
- 87 Brickman AM, Manly JJ, Honig LS, *et al.* Plasma p-Tau181, p-tau217, and other blood-based Alzheimer's disease biomarkers in a multi-ethnic, community study. *Alzheimers Dement* 2021;17:1353–64.