

Supplemental table 1: Differences in terms of baseline characteristics between patients with and without MRI

	Without MRI (n=30)	With MRI (n=144)	ASD (%)
Clinical characteristics			
Age, years, mean \pm SD	70.4 \pm 14.1	63.4 \pm 13.1	49.0
Female gender	20 (66.7)	46 (31.9)	74.1
Previous stroke	4 (13.3)	10 (6.9)	21.3
Atrial fibrillation	6 (20.0)	19 (13.2)	18.4
Arterial Hypertension	18 (60.0)	89 (61.8)	3.7
Diabetes mellitus	4 (13.3)	22 (15.3)	5.6
Dyslipidemia	13 (43.3)	43 (29.9)	28.2
Smoking	5 (16.7)	33 (22.9)	15.7
Heavy alcohol intake	5 (16.7)	52 (36.1)	45.2
Treatment			
Antiplatelet	9 (30.0)	29 (20.1)	22.9
Anticoagulants	6 (20.0)	18 (12.5)	20.4
Anti-hypertensive	20 (66.7)	79 (54.9)	24.4
Statins	11 (36.7)	28 (19.4)	39.1
Stroke characteristics			
NIHSS, median (IQR)	7 (4 to 10)	6 (3 to 11)	12.5
ICH volume ml, median (IQR)	3.4 (0.7 to 15.9)	5.8 (2.0 to 16.7)	32.9

Data are n (%) or median (IQR) unless otherwise indicated. Abbreviations: CI=confidence interval; ASD = Absolute standard difference; mRS=modified Rankin scale; NIHSS= National Institutes of Health Stroke Scale; ICH: intracerebral hemorrhage