

eTable 3. Baseline Characteristics of Patients with LGII Antibody Encephalitis Stratified by Type of Acute Immunotherapy

	Patients (n = 118)	Untreated (n = 12)	Multiple IT (n = 36)	Monotherapy (n = 70)	P Value
Demographics					
Age-at-symptom onset, median (range)	66 (17-87)	61 (47-79)	65 (36-80)	66 (17-87)	.73
Male sex	78/118 (66%)	7/12 (58%)	20/36 (56%)	51/70 (73%)	.17
Clinical characteristics at presentation					
Memory loss	103/118 (87%)	10/12 (83%)	33/36 (92%)	60/70 (86%)	.65
mRS, median (range)	3 (0-5)	3 (1-3)	3 (2-5)	3 (0-5)	.29
Kokmen STMS scale score, median (range)	32 (18-38)	37 (22-38)	30.5 (18-38)	33 (24-38)	.32
Seizures (any type)	104/118 (88%)	8/12 (67%)	32/36 (89%)	64/70 (91%)	.05
Faciobrachial dystonic seizures	62/118 (53%)	3/12 (25%)	21/36 (58%)	38/70 (54%)	.14
Pilomotor	13/118 (11%)	0/12 (0%)	5/36 (14%)	8/70 (11%)	.56

Abbreviations: IT, immunotherapy; mRS, modified Rankin Scale; STMS, Short Test of Mental Status.