

Online Table 1. Proportion of patients eligible for and receiving intervention for acute stroke in SLSR of 2005-2010^{¥§}

Values are number receiving intervention/number eligible for intervention (%) unless stated otherwise

Variable	Socioeconomic Deprivation (Carstairs score – quintile)				P value
	Deprived (2 nd -5 th Q)		Least deprived (1 st Q)		
	N=946		N=244		
	n	%	n	%	
No with ischaemic stroke	818		192		
Thrombolysis	98/783	12.5	23/189	12.2	0.897
Aspirin started at any time in acute phase [†]	635/746	85.1	151/172	87.8	0.368
Aspirin started by 48 hours after stroke	472/523	90.3	112/124	90.3	0.980
No with failed swallow screen [‡]	288		69		
Fed by enteral route [‡]	149/280	53.2	36/68	52.9	0.968
Intravenous fluids *	521/954	54.5	125/223	56.1	0.674
No requiring physiotherapy or occupational therapy *	796		183		
Physiotherapy or occupational therapy received, n(%)	640/764	83.8	154/176	87.5	0.218
No requiring speech and language therapy, *n(%)	666		163		
Speech and language therapy received	423/625	67.7	108/156	69.2	0.710

[¥]data available from 2005. [§]All data analysis excluding patients with SAH. [†]within first seven days of stroke. [‡] Nasogastric or percutaneous endoscopic gastrostomy. *Analysis limited to patients admitted to hospital.