

Supplementary Material**Table of Contents**

Pages 2 - 4	<i>Supplementary Table 1: key observational studies on anticoagulation timing</i>
Page 5	<i>Supplementary references</i>

Supplementary Table 1: Key observational studies of outcomes after AF-associated acute stroke according to anticoagulation timing

Study	Population and OAC type	Timing groups	Data on stroke severity, size and HT	Follow-up period	Ischaemic outcomes	Bleeding outcomes
NOACISP (1)	155 participants 100% DOAC	≤7d: n = 100 >7d: n = 55	NIHSS by timing group: ≤7d: 3 (1-7) >7d: 7 (3-14)	3-6m (mean 140d)	IS annualised rate: ≤7d: 5.1% >7d: 9.3%	sICH annualised rate: ≤7d: 0.0% >7d: 0.0%
RAF-NOACs (2)	1127 participants 100% DOAC	0-2d: n = 153 3-14d: n = 710 >14d: n = 264	Mean NIHSS: 7.7 (6.2) Infarct size ^a : 40.9% small, 33.3% medium, 22.1% large Any HT at baseline: 9.1%	90d	IS, TIA, SE, sICH, and major extracranial bleeding: - 0-2d: 12.4% - 3-14d: 2.1% - >14d: 9.1%	
CROMIS-2 (3)	1355 participants 37% DOAC 63% VKA	0-4d: n = 358 >4d: n = 977	NIHSS by timing group: 0-4d: 2 (1-4) >4d: 6 (3-11)	90d	Ischaemic stroke: 0-4d: 0.8% >4d: 1.6%	sICH: 0-4d: 0.0% >4d: 0.2%
RELAXED (4)	1305 participants 100% DOAC	<3d: n = 584 3-7d: n = 435 8-14d: n = 198 >14d: n = 88	Median NIHSS: 8 (3-17) Median initiation by NIHSS: <4: 2.8d 4-13: 3.5d >14: 4.6d Median initiation by size:	90d	Ischaemic stroke: <3d: 1.5% 3-7d: 2.5% 8-14d: 2.0% >14d: 6.8%	sICH: <3d: 0.2% 3-7d: 0.9% 8-14d: 0.0% >14d: 0.0%

			<p>- <4cm³ (34.0%): 2.9d</p> <p>- 4 – 22.5cm³ (32.5%): 2.9d</p> <p>- >22.5cm³ (33.4%): 5.8d</p>			
SAMURAI-NVAF (5)	499 participants 100% DOAC	<p>≤72h: n = 223</p> <p>>72h: n = 276</p>	<p>NIHSS by timing group:</p> <p>≤72h: 3 (1-8)</p> <p>>72h: 5 (2-13.5)</p> <p>Infarct size ^b by timing group:</p> <p>≤72h: 36.4% small, 55.3% medium, 8.3% large</p> <p>>72h: 25.3% small, 55.5% medium, 19.2% large</p>	90d	<p>IS/TIA:</p> <p>≤72h: 9.4%</p> <p>>72h: 10.5%</p>	<p>sICH:</p> <p>≤72h: 0.5%</p> <p>>72h: 1.5%</p>
IAC (6)	1289 participants 67.6% DOAC 31.0% VKA	<p>0-3d: n = 617</p> <p>4-14d: n = 535</p> <p>>14d: n = 137</p>	<p>NIHSS by timing group:</p> <p>0-3d: 5 (1-9)</p> <p>4-14d: 10 (4-16)</p> <p>>14d: 15 (9-21)</p> <p>Infarcts >60cm³ by timing group:</p> <p>0-3d: 5.9%</p> <p>4-14d: 17.4%</p> <p>>14d: 32.8%</p> <p>Any (HI/PH) HT at baseline:</p> <p>0-3d: 8.6%</p> <p>4-14d: 18.7%</p> <p>>14d: 48.9%</p>	90d	<p>IS/TIA/SE:</p> <p>0-3d: 10.3%</p> <p>4-14d: 9.7%</p> <p>>14d: 10.2%</p>	<p>sICH:</p> <p>0-3d: 0.3%</p> <p>4-14d: 1.7%</p> <p>>14d: 2.9%</p>

ALVO (7)	686 participants 100% DOAC Acute LVO or stenosis	<48h: n = 263 ≥48h: n = 423	NIHSS by timing group: <48h: 12 (7-18) ≥48h: 14 (6-20)	90d	IS: <48h: 3.5% ≥48h: 0.5%	sICH: <48h: 1.1% ≥48h: 1.0%
-------------	---	--------------------------------	--	-----	---------------------------------	-----------------------------------

DOAC: direct oral anticoagulant; IS: ischaemic stroke; LVO: large vessel occlusion; OAC: oral anticoagulant; SE: systemic embolism; sICH: symptomatic intracranial haemorrhage; TIA: transient ischaemic attack; VKA: vitamin K antagonist

^a small: <1.5cm³; medium: MCA deep or superficial branch, internal borderzone, superficial ACA or PCA branch; large: complete ACA, MCA, or PCA territory, two superficial or one superficial and one deep MCA branches, >1 arterial territory, lesion >1.5cm³ in brainstem or cerebellum; ^b small: <1.5cm maximum diameter; large:>1/3rd ACA, MCA, or PCA territory, or cerebellar hemisphere; medium: all others.

Supplementary References

1. Seiffge DJ, Traenka C, Polymeris A, Hert L, Peters N, Lyrer P, et al. Early start of DOAC after ischemic stroke: Risk of intracranial hemorrhage and recurrent events. *Neurology*. 2016 Nov 1;87(18):1856–62.
2. Paciaroni M, Agnelli G, Falocci N, Tsivgoulis G, Vadikolias K, Liantinioti C, et al. Early Recurrence and Major Bleeding in Patients With Acute Ischemic Stroke and Atrial Fibrillation Treated With Non-Vitamin-K Oral Anticoagulants (RAF-NOACs) Study. *J Am Heart Assoc*. 2017 Nov 29;6(12).
3. Wilson D, Ambler G, Banerjee G, Shakeshaft C, Cohen H, Yousry TA, et al. Early versus late anticoagulation for ischaemic stroke associated with atrial fibrillation: multicentre cohort study. *J Neurol Neurosurg Psychiatry*. 2019 Mar;90(3):320–5.
4. Yasaka M, Minematsu K, Toyoda K, Mori E, Hirano T, Hamasaki T, et al. Rivaroxaban administration after acute ischemic stroke: The RELAXED study. *PLoS One*. 2019;14(2):e0212354.
5. Mizoguchi T, Tanaka K, Toyoda K, Yoshimura S, Itabashi R, Takagi M, et al. Early Initiation of Direct Oral Anticoagulants After Onset of Stroke and Short- and Long-Term Outcomes of Patients With Nonvalvular Atrial Fibrillation. *Stroke*. 2020 Mar;51(3):883–91.
6. Yaghi S, Trivedi T, Henninger N, Giles J, Liu A, Nagy M, et al. Anticoagulation Timing in Cardioembolic Stroke and Recurrent Event Risk. *Ann Neurol*. 2020;88(4):807–16.
7. Yoshimura S, Uchida K, Sakai N, Imamura H, Yamagami H, Tanaka K, et al. Safety of Early Administration of Apixaban on Clinical Outcomes in Patients with Acute Large Vessel Occlusion. *Transl Stroke Res*. 2020 Aug 7;