CATAPLEXY

By

S. A. KINNIER WILSON, LONDON

THE peculiar phenomenon now usually described as cataplexy is no clinical novelty. Its occurrence in association with, or relation to, the syndrome of narcolepsy is well known; it is sometimes, though erroneously, taken to form an integral part of that syndrome, and is therefore rather apt to be regarded as an abnormal symptom which was unknown until narcolepsy began, as it has of late, to occupy neurological attention more than ever. But while cataplexy as a symptom has features which can be at once detected if it is at all pronounced, minor degrees of the same phenomenon are less easy to recognize, while its physiology and possible biological meaning, as well as its underlying mechanism, are still obscure.

In this brief communication I should like to illustrate the clinical aspects of cataplexy by reference to several fresh cases that have come under my notice, and to make some attempt at understanding its 'meaning' and at discovering the anatomophysiological basis of its remarkable manifestations.

Some years ago I published a fairly full account of an attack of cataplexy which I had the good fortune to observe from beginning to end as it occurred in a narcoleptic patient; during that attack I was able to prove that the tendon-reflexes were abolished for the time being, and that an extensor plantar response was present. The essence of the phenomenon was its reversibility; within a few minutes there was a complete return to the normal status quo ante. One or two further examples may now be cited.

Case I.—A male patient, age 36, has suffered from characteristic attacks of narcolepsy for about six years. In addition, he has developed cataplectic fits during the course of the last four years. These are induced by stimuli of an emotional kind, though of diverse qualities.

He had been sitting in my consulting-room for about an hour and was, possibly, rather fatigued, though not, so far as I could see, excited or anxious; nor had I at that moment been making him laugh. It was about seven o'clock in the evening and he had had a tiring day. However this may be, he suddenly said, 'it's coming on now,' in a slightly indistinct voice, and at the same moment his head fell forward on his breast in a series of little jerking movements. His legs sprawled out in front, but he maintained his seat in the arm-chair, nor did his hands let go of the arms of the chair. There was no alteration of colour, and, as I observed, no particular change in the pulse-rate. Nevertheless he was immobile and speechless for the time. I at once found complete abolition of knee-jerks and ankle-jerks, and on taking off his socks was unable to elicit any plantar response, either flexor or extensor, though his feet were not cold. A few seconds later, he moved his right foot to and fro with a series of slowish movements, limited in range. I asked him what he was doing, but he made no reply. The tonus of the musculature was reduced to one of profound limpness.

About two minutes later, he shook his head from side to side and muttered in an almost unintelligible voice, that he was 'coming round' and that the thing was 'passing off.' A moment later he sat up, smiled, and told me in a louder voice that 'it was over.' I now found that his knee-jerks were sluggish, and that his plantars were in feeble flexion. He said, in answer to a question, that he had moved his right foot intentionally, it being the sole part of his whole musculature that he could move. He was inwardly struggling to 'break the spell' of immobility—in other words, he was in possession of his conscious faculties and was making every effort to innervate his muscles—yet the only outward, visible, sign was a feeble little shaking movement of the right foot.

While this conversation between us was going on, he once more suddenly relapsed into the cataplectic state, and the same phenomena appeared again, just as before. Again there was recovery, and yet again there was relapse. Thus a succession of cataplectic fits—or, at least, a series of partial attacks and partial recoveries—took place after the first one. The time taken from beginning to end, when at length the patient emerged to a normal state without further cataplectic manifestations, was 14 minutes.

The interesting feature of this case, here briefly reported, consists in the occurrence of a *status cataplecticus*, of which, so far as is known to me, no instance has hitherto been put on record. Perhaps the phase might also be described as *serial cataplexy*; whichever term be preferred, the fact remains that the clinical happening brings cataplexy, by another of its aspects, into line with certain aspects of epilepsy—and this, in my opinion, has its importance.

A second feature is the abolition of the plantar reflex, which would seem to indicate that reflectivity as a whole, both cutaneous and deep, is in abeyance during cataplexy; and that while the flexor response is absent an extensor response may not take its place.

Case II.—A male, age 58, has had characteristic attacks of both narcolepsy and cataplexy for at least 18 years, i.e., since 1914 (incidentally, they began before epidemic encephalitis as known at present made its appearance in England).

During the course of my examination in the consulting-room, without any particular affective stimulus so far as was obvious to me, except that perhaps I had 'stimulated' him by remarking that I wished I could see an attack and that he was not going to 'oblige' me, his wife suddenly said, 'He's going to have one now'; and as I looked at him his head sank forward and downward, his eyes closed, little flickers appeared round the corners of the mouth. The resemblance to some types of petit mal was very close indeed, but there was no loss of colour, and the body and limbs were immobile. I spoke to him but he made no answer. His knee-jerks, previously active, were at this moment lost. I did not have time to take off his shoes to try the plantar responses, for he lifted his head and said in an indistinct voice that 'there was nothing the matter.' The total duration was but a few seconds, certainly not more than 30; and it appeared that all his cataplectic attacks are similarly short.

Of especial interest in this case, aside from the fact that had I not known its nature the superficially close likeness to petit mal might have been misleading, are features which the patient himself narrated in connexion with the cataplectic attacks. In the first place, they are brought on by 'anything

unusual or unexpected,' any experience whose emotional content is definite enough but which varies widely in degree. They were first noticed when he was fishing ; this sport he only took up in adult life, having never fished as a boy. Whenever he hooked a fish he at once 'went helpless,' his legs giving way underneath him, and his grasp on the rod relaxing. So constantly did this—to him unforeseen and inexplicable—phenomenon recur that finally he gave up fishing altogether. The attack, secondly, is accompanied by definite physical sensations : he has a peculiar feeling at the pit of the stomach, 'like a thrill' ; he has 'a catching at the heart,' sometimes amounting to palpitation ; and, more definite still, it is invariably attended by marked dryness of the mouth and throat—a *xerostomia* which he seeks to overcome by moving tongue and lips, yet the outward expression of this effort, as we have seen, is no more than a few flickers which might pass for those of *petit mal*.

It is generally admitted that one of the most distinctive characteristics of cataplexy is its occurrence as a sequel to some or other emotional stimulus, of which possibly the most common is one connected with the funny side of things and with the act of laughter. Hence it becomes of interest for our subject to look a little more closely at what actually occurs when laughter grips our sides. There is, first of all, involvement of facial and respiratory musculatures, which is familiar to all ; but if laughter is overwhelming, more escapes from control ; the person concerned 'shakes' with laughter, he is 'convulsed' with laughter, he cannot maintain his erect position but may roll on the floor. That is to say, laughter lets loose a series of diffuse, even convulsive movements—and yet the subject is fully conscious, exactly as is the cataplectic despite outward appearances. Again, however, the converse may take place ; instead of being the plaything of movement, the opposite condition affects his frame ; he is 'helpless' from laughing, he is 'weak' with laughter, he 'nearly dies' with it. Surely, then, we have here an instance of a double effect, or double possibility—either 'convulsion' or 'paralysis.' The affective stimulus either excites or inhibits, or inhibition succeeds excitation ; and for some reason at present insufficiently understood, in the case of cataplexy the stimulus seems invariably to inhibit.

Further, the avalanche may extend beyond the skeletal to the nonstriated musculature ; visceral accompaniments are obvious enough in the case of laughter—the skin glows and reddens, the eyes sparkle, increased glandular secretion occurs. Analogous or similar phenomena may be in evidence when the subject is seized with weeping as the result of an appropriate emotional stimulus ; his body is 'shaken' with sobs, he too is 'weak' from the experience, and in certain cases the escape involves even decontrol of visceral apparatus, just as in cases of excessive laughter.

Brissaud recounts the history of one of his patients which is apposite in this connexion.

'Une dame lui annonce que son petit chien vient de mourir. "Vraiment (dit-il) ; oh ! que c'est regrettable !" Est-ce le dernier mot "regrettable" qui, par une sorte

de suggestion, réveille en lui une série d'idées tristes ? Toujours est-il qu'il se produit comme un déclenchement de tous les ressorts expressifs ; la simple tristesse de la physionomie entraîne les larmes ; après les larmes, les sanglots ; bien plus, ce malheureux homme n'est plus maître de ses sphincters.'

Thus at the bidding of imperious emotion released mechanisms come into action, and both bladder and rectum may void their content. That such occurrences are not confined to persons actually suffering from some nervous affection but may be developed in others who are 'normal,' there is abundant testimony from the colloquial speech of different races of mankind. The significant point which emerges for our discussion is, that there must be radiation of the original affective impulse from an emotional 'centre' to a number of somatic and visceral 'centres' in physiological relation therewith.

At this point it becomes desirable to note that varieties of the same series of events are to be found in some patients who also suffer from epilepsy, particularly if that epilepsy is conditioned by stimuli of a psychical—that is to say, affective—kind.

Case III.—A male patient, age 28, has suffered from epileptic fits, both major and minor, since he was 14 years old. The major attacks, which occur but rarely, are always preceded by a vivid dream, of an unusually terrifying sort ; he imagines in his dream that he is about to be hanged, or shot, for some misdemeanour—sometimes both hanged and shot (!) ; the origin of this nightmare has been successfully traced but need not here be detailed. Sometimes the dream antedates the fit by 24 hours, sometimes much less, but the fit itself only occurs in sleep, even sleep of a few minutes after a midday meal or for any other reason.

The minor attacks are so brief as to pass almost unnoticed, and do not appear to be activated by any discoverable stimulus.

In addition, however, the patient has observed for years that under any circumstances of excitement, pleasant or unpleasant, under the influence of unusual emotional events, he always loses control of his bowel. So much has this been the case that it has been regarded as a sort of 'family joke.' When out hunting rats with a terrier, when shooting pheasants, he has had repeatedly to 'retire' in view of the imminent evacuation of the rectal contents. This has sometimes happened three times of a morning, when engaged in sport. Similarly, when once called upon to make a speech unexpectedly at a social gathering, he had to leave the room.

The clinical features of this peculiar case are very instructive ; an emotional experience reverberates in his dreams and precipitates his epilepsy ; affective excitants also cause decontrol of visceral mechanisms. On the other hand, however, he has never had any attack recalling those of full-blown cataplexy.

Evidence which need not here be recapitulated has been gradually collecting in favour of the belief that the optic thalamus, in particular the palæothalamus, is related to afferent impulses laden with emotional tone ; and that the expression of the emotions, in movement of skeletal muscle and activation of neurosympathetic mechanisms, is initiated therefrom. To exclude the cortex however from the succession of events is impossible, for the impulses, coloured with affect, are appreciated in consciousness and have

therefore presumably reached the percipient cortex. Thalamofugal impulses are capable of acting on the cortex, in a plus or minus direction. They stimulate or inhibit, according to circumstances. One of the proofs of this is, that in cataplexy the patient may be unable to speak. No doubt the muscles of phonation and articulation are temporarily paralyzed in cataplexy, just like other musculatures; but when a person is 'speechless' he is something more than 'inarticulate.' The cataplectic may be 'deprived of speech' though the attack has not radiated through the whole body.

Case IV.—A case in point is that of a narcoleptic and cataplectic subject who once saw approaching him on a thronged street a friend whom he had not seen for many years. At the very moment of pleasant anticipation, just when he was walking forward to greet him, the emotion of the moment overpowered him; 'rooted to the spot,' he could not hold out his hand, could not say a single word; the friend passed without observing him and was lost in the crowd; by the time my patient had recovered, he had disappeared.

It is, I submit, plausible that inhibition of the cortex had taken place as a result of stimuli reaching it via the thalamus. But we must hasten also to note that this postulated inhibition is local and restricted. For observation and the statements of our patients convince us that during the average fit of cataplexy consciousness is preserved and therefore the cortex as a whole has not undergone abrogation of its functions. There is a block somewhere between 'volition' and 'innervation'—just as there is in hysteria. Where precisely this occurs, in an anatomophysiological sense, is peculiarly difficult to determine; it appears to me more feasible to suggest that it occurs at the middle (rolandic) level than anywhere else. We know that the 'will to innervate' is maintained, for sometimes cataplectic patients say that they 'struggle inwardly' to move, though nothing is effected.

Yet the fact must not be lost sight of, that the musculature is, for the time being, emptied of tone, and that the reflexes are in abeyance. Ordinarily, when lesions of the upper motor neurones develop slowly, lower automatisms come into increased activity, so that the reflexes are exaggerated and involuntary spasmodic movements may occur. But in the case of sudden shock-like lesions of the upper neurones, the lower neurones are simultaneously affected, and reflectivity may be entirely lost. Hence the flaccidity and the reflex loss of the limbs cannot—in the case of cataplexy, which is as fulminant as any shock—be taken by itself to prove that the site of the physiological disorder is at a low (spinal) level. The truth may well be, that instantaneous inhibition radiates through upper and lower motor neurones alike, universally.

It is striking to think that this can and does occur with such rapidity because of emotional experience, by no means always profound—sometimes trifling or almost unrecognizable; but it supplies an excellent illustration of the reaction of the organism *as a whole* to affective stimuli.

If, then, we may suppose that the motor characteristics of cataplexy are the visible expression of total and immediate paralysis of the skeletal

musculature by inhibition of its nerve-cells of origin—a single massive motor unit—a ‘motor ganglion,’ acted on powerfully by sensory excitation, we must also seek to ascertain whether the hypothesis will account for the atony of the muscles. That tonus must be represented at the cortical level has long been considered by myself a rational inference from various clinical data at our disposal ; and at the Berne Congress of 1931 I made a brief communication urging the need to accept this view if these clinical data are to be reasonably explained. At the same Congress Sir Charles Sherrington tentatively expressed the conclusion to which he was inclined to turn, viz., that tonus and movement are one and the same. ‘Tone follows movement like a shadow.’ Should this theory receive support, it will explain the motor phenomena of cataplexy without much difficulty ; for we should then affirm that inhibition of the unified ‘motor ganglion’ from the sensory side simultaneously entails inhibition of tonus throughout the musculature innervated from that cortical ‘ganglion.’

In addition, however, to thalamofugal action on the motor cortex, thalamohypothalamic radiation is equally characteristic of the cataplectic attack. Immediately preceding the latter, or with it, symptoms occur which are referable to visceral centres situated round the ventricular system, reaching from the floor and sides of the third to the floor of the fourth ; the patient ‘loses his breath,’ feels ‘hot’ or ‘cold,’ has a sensation at the pit of the stomach, his heart palpitates, and so forth. During the fit itself there may be pallor, perspiration, etc. The xerostomia of one of my cases is also classifiable in this neurovisceral series (it is interesting to note that the symptom has been observed in thalamic lesions apart altogether from its appearing in connexion with cataplexy). Although the main incidence of the attack is on skeletal musculature, these visceral concomitants merit attention, since they indicate spread downwards as well. Further proof of this is provided by the observed fact that sometimes cataplectic and narcoleptic seizures are interchangeable, sleep taking the place of the motor phenomena, and there is good reason to believe that sleep is related to the function of some periventricular ‘centre.’

The emotion which is so profound as to cause relaxation of sphincters does not seem to characterize cataplexy ; nevertheless we have the record of one of Redlich’s cases, in which hearty laughing led to an attack of powerlessness which culminated in loss of both vesical and rectal control. Extensive radiation of the kind, however unusual, may therefore be found on occasion ; and in this respect we find an epileptic analogy worthy of note.

The ‘storm centre’ of the cataplectic seizure would thus appear to be thalamic, with immediate spread to cortex in part and to hypothalamus and ventricular floors in part.

It is desirable, in conclusion, to take a still wider view of the phenomena under discussion. Have they a biological meaning ? To quote from a previous article of my own : ‘Is it conceivable that we have here through disease

stumbled upon a biologically old automatism, a defence-reaction in which immobility is of advantage to the frightened animal, "struck all of a heap"? Is cataplexy akin to the Totstellreflex, where the animal is "frightened out of its life"? The cartoonist of the War depicted the helpless, paralysed feeling of the soldier as the shell approached him by sketching enormous weights chained to his feet. He could not move for fear. Emotion inhibited movement. In certain circumstances it must have meant safety to the animal to lie motionless, as still as the dead. Some species of animal react thus with comparative regularity. From this aspect at least, the phenomena of the affective cataplectic fit fall into line with reactions phylogenetically ancient, and are capable of psychological and teleological explanation.

But the difficulty remains: in clinical cases cataplexy results more commonly from activation on the side of laughter, and has indeed been observed when joy or gladness has provided the excitant. It is not easy to see how the hilarious mood should cause an immobility of biological advantage to the person concerned. Yet, once again, everyday parlance has the expression, 'I nearly died of laughing.' It would seem, perhaps, that the mechanism once utilized by fear to produce motionlessness can also be activated by an emotion of opposite quality, and that what has a biological meaning in one set of circumstances has little or none in another. The question, however, needs closer consideration than can be allotted to it in this place.

It is also worth remarking that the matter of unexpectedness has much to do with the onset of the cataplectic attack. If the subject is prepared beforehand, it may never evolve. Forewarned is forearmed in a physiological as well as in a military sense. The cortex can exert a corticifugal, cortico-thalamic inhibitory influence before thalamocortical impulses get their start. This fact of clinical observation has often been confirmed; and it is of interest because it is not paralleled to anything like the same extent in the case of epilepsy. In the case of the latter, cortical inhibition counts almost for nothing. May we infer that corticothalamic inhibition has greater momentum than corticocortical inhibition?

Cataplexy is a physiological experiment evolving before the clinician's eyes, and giving him a chance of insight into the working of neural mechanisms more revealing than that provided by epilepsy. He should take every opportunity of studying it minutely, for he has this great advantage over the experimentalist, that the phenomena of cataplexy are at present incapable of being reproduced in the laboratory.